

Homework Problems

Section 1.1

Problem 4

- a. $\lceil 0.763 \rceil = \boxed{1}$
- b. $2\lceil 0.6 \rceil - \lceil 1.2 \rceil = 1 - 2 = \boxed{-1}$
- c. $\lceil 1.1 \rceil + \lceil 3.3 \rceil = 2 + 4 = \boxed{6}$
- d. $\lceil \sqrt{3} \rceil - \lfloor \sqrt{3} \rfloor = 2 - 1 = \boxed{1}$
- e. $\lceil -73 \rceil - \lfloor -73 \rfloor = -73 - (-73) = -73 + 73 = \boxed{0}$

Problem 8

How many multiples of 10 are there between the following pairs of numbers?

This can be solved for any pair n, m such that $m \leq n$ using the formula:

$$\left\lfloor \frac{n}{10} \right\rfloor - \left\lfloor \frac{m-1}{10} \right\rfloor$$

- a. $m = 1, n = 80$

$$\left\lfloor \frac{80}{10} \right\rfloor - \left\lfloor \frac{1-1}{10} \right\rfloor = 8 - 0 = \boxed{8}$$

- b. $m = 0, n = 100$

$$\left\lfloor \frac{100}{10} \right\rfloor - \left\lfloor \frac{0-1}{10} \right\rfloor = 10 - (-1) = \boxed{11}$$

Note that 0 is a multiple of every integer including 10, so it is included in this count.

- c. $m = 9, n = 2967$

$$\left\lfloor \frac{2967}{10} \right\rfloor - \left\lfloor \frac{9-1}{10} \right\rfloor = 296 - 0 = \boxed{296}$$

- d. $m = -6, n = 34$

$$\left\lfloor \frac{34}{10} \right\rfloor - \left\lfloor \frac{-6-1}{10} \right\rfloor = 3 - (-1) = \boxed{4}$$

- e. $m = 10^4, n = 10^5$

$$\left\lfloor \frac{10^5}{10} \right\rfloor - \left\lfloor \frac{10^4-1}{10} \right\rfloor = 10000 - 999 = \boxed{9001}$$

Problem 18

a. Fact 4 states that between any number n and k there are $\lfloor \frac{n}{k} \rfloor$ multiples of k between 1 and n . Therefore, in the case $k = 1$, there are n positive integers between 1 and n . In effect, the count of numbers between 1 and any positive integer n is simply n . This is obvious as any given count of a collection of size n is the value of n elements.

b. Given fact 4 once more, if $k > n$, then any two given numbers of k and n will result in $\lfloor \frac{n}{k} \rfloor = 0$ multiples of k between 1 and n . This is again obvious as a number greater than another given number cannot evenly divide it into an integer value ≥ 1 .

Problem 19

a.

$$\text{Let } x = 0.5, y = 0.5. \text{ Then } \lfloor 0.5 \rfloor + \lfloor 0.5 \rfloor < \lfloor 0.5 + 0.5 \rfloor \\ 0 + 0 < \lfloor 1 \rfloor \rightarrow 0 < 1$$

Therefore the inequality holds for $y = 0.5, x = 0.5$

b.

$$\text{Let } x = 0, y = 0. \text{ Then } \lfloor 0 \rfloor + \lfloor 0 \rfloor = \lfloor 0 + 0 \rfloor \\ 0 + 0 = \lfloor 0 \rfloor \rightarrow 0 = 0$$

Therefore the equality holds for $y = 0, x = 0$

c. All floor operations round down to the nearest integer. Any given pair, x, y must either contain integers or non-integers. If both are integers, then the equation $\lfloor x \rfloor + \lfloor y \rfloor$ simplifies always to $x + y$. In any case for x, y that at least one is a non-integer, the floor operator will reduce that value such that $\lfloor x \rfloor < x$ and/or $\lfloor y \rfloor < y$. Therefore, $\lfloor x \rfloor + \lfloor y \rfloor \leq x + y$.

We know that $\lfloor x + y \rfloor$ is the largest integer less than or equal to $x + y$ and that $\lfloor x \rfloor + \lfloor y \rfloor$ is an integer.

Given these facts, $\lfloor x \rfloor + \lfloor y \rfloor$ must also always be less than or equal to $\lfloor x + y \rfloor$.

Section 1.2

Problem 12

No, two different even integers cannot be relatively prime. We define a given integer n to be even if $2|n$. Any given pair of even integers n, m can be expressed as $n = 2k$ and $m = 2j$ for some integers k, j . The factor 2 in these equations would be a common factor that is not 1, so the two even integers cannot be relatively prime.

Problem 14

a. We can expand the expression lm into the expression adl . We can then very clearly see that both d and adl share the common factor d . Therefore, in any case where l is an integer, ml is evenly divided by d , thus $d \mid ml$.

b. In both cases, we can factor out d , which inherently defines the operations as factors of d .

$$m + n = ad + bd = d(a + b)$$

$$m - n = ad - bd = d(a - b)$$

c. We can see that $17m - 72n = 17ad - 72bd = d(17a - 72b)$. As both a and b are integers, so must $17a - 72b$. Therefore, d divides $17m - 72n$.

Problem 16 b.

The factors of 36 are: 1, 2, 3, 4, 6, 9, 12, 18, 36. All numbers not sharing these factors are therefore relatively prime (except for sharing 1). Those are: 1, 5, 7, 11, 13, 17, 19, 23, 25, 29, 31, and 35.

Section 1.3

Problem 2

List the elements ($\{\}$) in the following sets:

a.

$$\left\{ \frac{1}{n} : n = 1, 2, 3, 4 \right\} = \left\{ 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4} \right\}$$

b.

$$\{n^2 - n : n = 0, 1, 2, 3, 4\} = \{0, 0, 2, 6, 12\}$$

c.

$$\left\{ \frac{1}{2^2} : n \in \mathbb{P}, n \text{ is even and } n < 11 \right\} = \left\{ \frac{1}{4}, \frac{1}{16}, \frac{1}{36}, \frac{1}{64}, \frac{1}{100} \right\}$$

d.

$$\{2 + (-1)^n : n \in \mathbb{N}\} = \{3, 1, 3, 1, \dots\} \text{ or simply } \{3, 1\}$$

Problem 6

Repeat Exercise 4 for the following sets (determine the following sets):

a.

$$\{n \in \mathbb{N} : n|12\} = \{1, 2, 3, 4, 6, 12\}$$

b.

$$\{n \in \mathbb{N} : n^2 + 1 = 0\} = \emptyset$$

as all values of n result in $n^2 + 1 > 0$

c.

$$\left\{ n \in \mathbb{N} : \left\lfloor \frac{n}{3} \right\rfloor = 8 \right\} = \{24, 25, 26\}$$

d.

$$\left\{ n \in \mathbb{N} : \left\lceil \frac{n}{2} \right\rceil = 8 \right\} = \{15, 16\}$$

Problem 8

How many elements are in the following sets? Write ∞ if the set is infinite.

a.

$$\{n \in \mathbb{N} : n^2 = 2\} : \boxed{\emptyset} \text{ as no integer } k \text{ exists such that}$$

$$\sqrt{2} = k \text{ and } k \in \mathbb{N} \text{ as } \sqrt{2} \approx 1.41$$

b.

$$\{n \in \mathbb{Z} : 0 \leq n \leq 73\} : (73 - 0) + 1 = \boxed{74}$$

c.

$$\begin{aligned} \{n \in \mathbb{Z} : 5 \leq |n| \leq 73\} &: (73 - 5) + 1 - (5 - 73) + 1 \\ &= 68 - (-68) + 2 = 136 + 1 = \boxed{138} \end{aligned}$$

d.

$$\{n \in \mathbb{Z} : 5 < n < 73\} : 73 - 5 - 1 = \boxed{67}$$

e.

$$\{n \in \mathbb{Z} : n \text{ is even and } |n| \leq 73\} :$$

Using the formula defined in worksheet 1 to determine

$$\text{all even numbers between } m \text{ and } n : \left\lfloor \frac{n}{2} \right\rfloor - \left\lfloor \frac{m-1}{2} \right\rfloor$$

$$\left\lfloor \frac{73}{2} \right\rfloor - \left\lfloor \frac{0-1}{2} \right\rfloor = 36 - 0 = 36$$

$|n|$ includes negative even integers as well, thus $36 * 2 = 72$

Including 0 as an even integer gives a final total of $72 + 1 = \boxed{73}$

f.

$\{x \in \mathbb{Q} : 0 \leq x \leq 73\} : \boxed{\infty}$, as there are infinite rational numbers between any two given integers.

g.

$\{x \in \mathbb{Q} : x^2 = 2\} : \boxed{0}$ as there is no number defined as

$m \in \mathbb{Z}, n \in \mathbb{Z}$ such that $\left(\frac{m}{n}\right)^2 = 2$ or $\frac{m}{n} = \sqrt{2}$, because $\sqrt{2}$ is irrational.

h.

$\{x \in \mathbb{R} : x^2 = 2\} : \boxed{2}$, we defined before that the number $\sqrt{2}$ satisfies this condition.

\mathbb{R} includes irrational numbers whereas \mathbb{Q} does not, allowing for $\sqrt{2}$.

Additionally, $-\sqrt{2}$ also satisfies this condition as $(-\sqrt{2})^2 = 2$, making the set be $\{-\sqrt{2}, \sqrt{2}\}$