## MATH-253: HW1

Due on 1/22/2024

Prof. Oleksandr Bobrovnikov (Sasha), Spring 2024, 1/22/2024

**Kaleb Burris** 

## 2.1

a) 
$$\overrightarrow{RP} = \langle -1 - (-3), 3 - 7 \rangle = \underline{\langle 2, -4 \rangle}$$

b) 
$$\overrightarrow{PQ} = \underline{2\vec{i} - 4\vec{j}}$$

a) 
$$\overrightarrow{PQ}=\langle 2,2\rangle, \overrightarrow{PR}=-\overrightarrow{RP}=\langle -2,4\rangle.$$
 Therefore:

$$\begin{aligned} 2\overrightarrow{PQ} - 2\overrightarrow{PR} &= 2 \cdot \langle 2, 2 \rangle - 2 \cdot (-2, 4) \\ &= \langle 4, 4 \rangle - (-4, 8) \\ &= \underline{\langle 8, -4 \rangle} \end{aligned}$$

b) 
$$8\vec{i} - 4\vec{j}$$

$$\|\overrightarrow{PQ}\| = \sqrt{2^2 + 2^2} = \sqrt{8} = 2\sqrt{2}$$

a) 
$$\langle 1, 1 \rangle$$

b) 
$$\vec{i} + \vec{j}$$

$$\vec{v} = \langle 2 - (-1), 1 - (-3) \rangle = \langle 3, 4 \rangle$$
$$\|\langle v \rangle\| = \sqrt{3^2 + 4^2} = \sqrt{9 + 16} = \sqrt{25} = 5$$
$$/3 \quad 4 \setminus 3$$

$$\vec{v} = \langle x - 1, 0 - 1 \rangle = \langle x - 1, -1 \rangle$$
 
$$\|\vec{v}\| = \sqrt{10} = \sqrt{(x - 1)^2 + (-1)^2} = \sqrt{x^2 - 2x + 2}$$
 
$$10 = x^2 - 2x + 2$$
 
$$0 = x^2 - 2x - 8$$
 
$$0 = (x + 2) \cdot (x - 4)$$
 
$$x = 2, -4$$
 
$$x \text{ must be } -4 : Q(-2, 0)$$

a) 
$$\vec{a} = \langle -2, 4 \rangle \quad \vec{b} = \langle -2, 2 \rangle$$

$$-3\langle -2, 4 \rangle + \langle -2, 2 \rangle - 4i + j = \langle 6, 12 \rangle + \langle -2, 2 \rangle + \langle -4, 0 \rangle + \langle 0, 1 \rangle$$

$$= \langle 0, 15 \rangle; \quad \|\langle 0, 15 \rangle\| = \underline{15}$$

$$||v|| = 3, u = \langle -2, 5 \rangle$$

$$\sqrt{(-2)^2 + 5^2} = \sqrt{29}$$

$$3\left\langle \frac{-2}{\sqrt{29}}, \frac{5}{\sqrt{29}} \right\rangle = \left\langle \frac{-6}{\sqrt{29}}, \frac{15}{\sqrt{29}} \right\rangle$$

$$\vec{u} = 6\langle \cos(60^{\circ}), \sin(60^{\circ}) \rangle = \underline{\langle 3, 3\sqrt{3} \rangle}$$

$$\vec{u} = 8\langle -1, 0 \rangle = \langle -8, 0 \rangle$$

$$lpha = 0; eta = 0$$
  $lpha = 2eta$   $lpha \in \mathbb{R}; eta = 2lpha$ 

a) 
$$f'(x) = 4x^{3}; f'(1) = 4$$
$$y = 4x + 1 \Rightarrow y = 4 + 1 = 5$$
$$\underline{Q(2, 5)}$$

b) 
$$\overrightarrow{PQ} = \langle 2-1, 5-1 \rangle = \langle 1, 4 \rangle; ||u|| = \sqrt{1^2 + 4^2} = \sqrt{17}$$
 
$$\overrightarrow{u} = \left\langle \frac{1}{\sqrt{17}}, \frac{4}{\sqrt{17}} \right\rangle$$

$$\theta=30^{\circ}$$
 
$$\vec{v}=100\langle\cos(30^{\circ}),\sin(30^{\circ})\rangle=\left\langle50,50\sqrt{3}\right\rangle$$
 Horizontal: 50.00 mph, Vertical: 86.60 mph

2.2

a) 
$$(0,3,0), (2,0,0), (0,0,5), (2,0,5), (0,3,5), (2,3,0)$$

b) 
$$\sqrt{2^2 + 3^2 + 5^2} = \sqrt{4 + 9 + 25} = \sqrt{38}$$

$$z=2; z=5$$
 The set is 2 planes at  $z=2$  and  $z=5$