



Excited State Optical Transitions in Quantum Cascade Lasers for Lower Thresholds and Multi-Wavelength Emission

**Kale J. Franz, Stefan Menzel*, Anthony J. Hoffman,
Daniel Wasserman, Kuen-Ting Shiu**, Stephen R. Forrest**,
and Claire Gmachl**

Department of Electrical Engineering
Princeton University, Princeton, NJ 08544


* University of Sheffield, Sheffield S3 7RH, United Kingdom

** Current address: Department of Electrical Engineering and Computer Science
University of Michigan, Ann Arbor, MI 48109

designing a better QC laser



- QC emitters:
a “designer” material
- Limited by
design space





$$J_{th} = \frac{\alpha_m + \alpha_w}{g\Gamma}$$

$$g = \tau_u \left(1 - \frac{\tau_\ell}{\tau_{ul}} \right) \frac{4\pi q}{\epsilon_0 \lambda_0 n_{eff} L_p} \frac{z_{ul}^2}{2\gamma_{ul}}$$

Why is long wavelength so hard?



- Optical absorption
- Upper laser level lifetime
- Coupling efficiency





improving laser gain

optical dipole matrix element




$$g \propto z_{ul}^2 = \left| \langle \phi_u(z) | z | \phi_l(z) \rangle \right|^2$$

$$g \propto \frac{1}{2\gamma_{ul}}$$



especially useful for longer-wavelength optical transitions

15 μm excited state QC laser



- $E_{ul} = 83.6 \text{ meV}$
- $z_{ul} = 50.3 \text{ \AA}$
- $\tau_5 = 0.586 \text{ ps}$
- $\tau_4 = 0.215 \text{ ps}$
- $\tau_{54} = 1.9 \text{ ps}$
- $\text{FoM} = 1313 \text{ ps \AA}^2$
- $z^2 \tau_5 = 1483 \text{ ps \AA}^2$
- $z^2 \tau_5 E = 124 \text{ ps \AA}^2 \text{ eV}$


FoM comparison



$$\tau_3 z^2 E \text{ (ps } \text{\AA}^2 \text{ eV)}$$


69.8

C. Gmachl et al. (1998)
13 μm Diagonal Transition




56.0

A. Tredicucci et al. (1999)
17 μm Superlattice




70.2

M. Rochat et al. (2001)
15.6 μm B-to-C




123.9



15 μm Excited State (2007)





excited state QC laser the first attempt




lasing and EL spectra





field-dependent emission







high current EL





cascaded QC laser



Light – Current – Voltage



threshold behavior



summary



Appl. Phys. Lett. **90** 091104 (2007)

- Excited state transitions:
a strategy to lower
threshold currents
- Cascaded emission in
semiconductor lasers
- Acknowledgements
 - + MIRTHE (NSF-ERC)
 - + DARPA L-PAS
 - + POISE
 - + PCCM (NSF-MRSEC)
 - + NSF Graduate Research
Fellowship Program

