



Probiotics and prebiotics associated with aquaculture: A review



Najeeb Akhter^a, Bin Wu^{a,*}, Aamir Mahmood Memon^b, Muhammad Mohsin^b

^a Ocean College, Zhejiang University, Hangzhou 310058, China

^b College of Fisheries, Ocean University of China, Qingdao 266003, China

ARTICLE INFO

Article history:

Received 13 April 2015

Received in revised form

16 May 2015

Accepted 28 May 2015

Available online 1 June 2015

Keywords:

Aquaculture

Probiotics

Prebiotics

Immunosaccharides

Immunostimulants

Innate immunity

ABSTRACT

There is a rapidly growing literature, indicating success of probiotics and prebiotics in immunomodulation, namely the stimulation of innate, cellular and humoral immune response. Probiotics are considered to be living microorganisms administered orally and lead to health benefits. These Probiotics are microorganisms in sufficient amount to alter the microflora (by implantation or colonization) in specific host's compartment exerting beneficial health effects at this host. Nevertheless, Prebiotics are indigestible fiber which enhances beneficial commensally gut bacteria resulting in improved health of the host. The beneficial effects of prebiotics are due to by-products derived from the fermentation of intestinal commensal bacteria. Among the many health benefits attributed to probiotics and prebiotics, the modulation of the immune system is one of the most anticipated benefits and their ability to stimulate systemic and local immunity, deserves attention. They directly enhance the innate immune response, including the activation of phagocytosis, activation of neutrophils, activation of the alternative complement system, an increase in lysozyme activity, and so on. Prebiotics acting as immunosaccharides directly impact on the innate immune system of fish and shellfish. Therefore, both probiotics and prebiotics influence the immunomodulatory activity boosting up the health benefits in aquatic animals.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

A variety of useful feed additives, including probiotics and prebiotics having beneficial effects to the host was used in aquaculture to combat diseases such as supplements, to improve growth include increasing the size and weight gain, and in some cases, act as an alternative antimicrobial compounds [55], as well as to stimulate immunity response of the host. Probiotics live in microbial feed additives that modulate gastrointestinal microbial communities whereas Prebiotics, non-digestible forage additives stimulate the activity or abundance of beneficial gastrointestinal bacteria. Both of these have received widespread attention, showing the improved production, health and disease resistance of aquatic animals [30]. In addition increased research for the development of new strategies of food supplementation which were assessed in various health and growth promoting compounds such as; probiotics, prebiotics, synbiotics, phytobiotics and other functional food supplements were also evaluated [28].

Probiotic microorganisms are served for many purposes such as increasing economic growth, promoting digestion, absorption and suppression of infectious diseases [11,76]. Apart from immunomodulation, probiotics are also believed to have diverse modes of action in living organisms. For example in terrestrial animals, they competitively exclude potential pathogens by placing repressive molecules or through direct competition for space, nutrients or oxygen in the digestive tract of an organism [39]. However, in aquatic animals, they stick with the mucosal epithelium of gastrointestinal tract, such as *Brevibacillus brevis*, lactic acid bacteria, *Vagococcus fluvialis* and *Vibrio harveyi* [23,65,67,73,108,111] and help to resist pathogens [70]. However, it was also reported that probiotics do not dwell in the digestive tract [90]. Another mode of action of probiotics is that, they enhance the digestibility of food [117] in the form of enzymes such as alginate lyases, amylases and proteases [131]. They also boost the production of nutrients for example, fatty acids, biotin and vitamin B12 [112,113,120], which has positive effect on the health of an animal.

These are also used as a supplementary food for improving intestinal microbial balance that may offer beneficial effects to the host [40,105]. Probiotic bacteria are known to have immune stimulant effect [43,119]. Classically, certain prebiotics are non-digestible fibers that increase intestinal beneficial bacteria in a

* Corresponding author. Tel./fax: +86 571 8820 8540.

E-mail addresses: na_memon@yahoo.com (N. Akhter), wubin@zju.edu.cn (B. Wu), aamirmd_24@hotmail.com (A.M. Memon), mohsingrw@yahoo.com (M. Mohsin).

host [64,91]. Later proposed another type of functional saccharides; so-called immunosaccharides, which stimulate the immune system directly, rather than through the probiotic products. Prebiotics, such as fructooligosaccharide, mannanooligosaccharide, inulin or β -glucan, are known immunosaccharides. Probiotics were found to stimulate feed conversion efficiency, increase live weight gain of fish and shrimp culture [4] and by competitive exclusion to confer protection to a fixed area, to prevent pathogens [23,121], the production of organic acids, hydrogen peroxide, antibiotics, bacteriocins, siderophores and lysozyme [33,126]. They also regulate the immune response of fish [13,14].

Existing literature lack evidence that pro and prebiotics are safe for their host. Consequently, viability of an organism during its administration and retention of beneficial effects during prolonged storage or repeated sub-culturing is still questionable. A number of probiotics and prebiotics are now commercially available and in use. Although there are added costs associated with using these products, improved production efficiency and reduced disease incidence may offset such costs. Another possible way of simultaneous administration of pro and prebiotics, termed as synbiotic, is intended to improve the survival and implantation of the live microbial supplement in the GIT. However, meager information is available on the use of synbiotics in aquaculture. By understanding all the advantages that pro and prebiotics confer, it would be possible for us to apply them symbiotically only after knowing their benefits in aquaculture. So, over the last decade, the application of both probiotics and prebiotics that take advantage of their potential for pathogen control has increased in aquaculture. This document focuses on immunomodulatory action of probiotics and prebiotics, which directly stimulate the innate immune system of fish and shellfish. Furthermore, it is necessary to standardize the procedures used for the study of probiotics and prebiotics, particularly with regard to the precise nature of the congenital and or cellular immune response.

2. Probiotics

The Probiotics, has been initially defined as organisms or substances that contribute to intestinal microbial balance [84]. The term probiotic was emerged from the Greek words “Pro” and “bios” meaning “for life” [47] and is often referred as a support for life, that help naturally to improve the overall health of the host organism. According to the Food and Agriculture Organization (FAO) and the World Health Organization (WHO), probiotics are live microorganisms which, when administered in an appropriate amount provide health benefits to the host [36].

Nevertheless, aquaculture has been an expansion undertaken by FAO/WHO to include a wide range of Gram-positive and Gram-

negative bacteria, bacteriophages, microalgae and yeast (Table 1) with the application through the aqueous/water route, as well as by supplying the feed. The essence of the definition of FAO/WHO is that probiotics are living organisms, which are administered orally having some tangible health benefits, and have been widely used for the control of diseases in aquaculture, especially in developing countries [11,37,56,60,76].

In practice, there is only limited evidence that the viability of probiotic cultures checked as soon as they are added to feed, and the health benefits may be inaccurately described. Of course, it may be an impact on the viability of the processing result of diet, namely pelleting/granulation and extrusion. For example, the administration of the probiotic extruded diet led to an improvement in nonspecific immunity of *Nile tilapia* compared with granulated/pelleted diet [103].

Stimulation of nonspecific host defense mechanisms, using specific biological compounds called immune stimulant, increases resistance to disease and the growth of the host [72]. The innate immune system containing physical barriers and cellular and humoral components is a self-defense weapon in fish [38]. Live bacteria as probiotics act alternatively for antibiotics and chemicals, and functions as signaling molecules to activate the immune system.

There are many beneficial effects of probiotics, one among which is the modulation of the immune system. The role of probiotics in the modulation of the immune system was extensively studied and reviewed in humans and animals [6,41]. Previous studies dealt with growth-promoting and disease protection ability of probiotics in fish. But in recent years much attention has so far been in the direction of the immunomodulatory effects of probiotics in aquaculture. Many immunological researches have been conducted in various fish, using different probiotics and their validity to stimulate fish immunity, both within invivo and invitro conditions is remarkable [13,53]. Probiotics have also been identified for use as immunomodulatory agents in shellfish like shrimp [8,51,88].

3. Strengthening immune response

The innate immune system, also known as non-specific immune system is the first line of defense which comprises the cells and mechanisms that defend the host from infection by other organisms. Probiotics interact with the immune cells such as mononuclear phagocytes (monocytes, macrophages) and polymorphonuclear leukocytes (neutrophils), natural killer (NK) cells, to enhance innate immune responses. Some probiotics can increase the number of erythrocytes, granulocytes, macrophages and lymphocytes in various fishes [55,57,62,63,66,77]. In the same

Table 1
Probiotics used in aquaculture acting on specific pathogens (updated from Ref. [79]. The pathogens controlled by the probiotics are given as Latin small letters.

Category of organism	Genus/groupings
Gram-positive bacteria	<i>Arthrobacter</i> (q, s), <i>Bacillus</i> (b, d, q, x), <i>Brevibacillus</i> (m), <i>Brochothrix</i> (a), <i>Clostridium</i> (b, n), <i>Carnobacterium</i> (n, r, x), <i>Enterococcus</i> (e, q, s), <i>Kocuria</i> (n, r), <i>Lactobacillus</i> (c, g, j, l), <i>Lactococcus</i> (n), <i>Leuconostoc</i> (g), <i>Microbacterium</i> (n), <i>Micrococcus</i> (b, c), <i>Pediococcus</i> (h, n), <i>Rhodococcus</i> (n), <i>Streptococcus</i> (q), <i>Streptomyces</i> (q, u), <i>Vagococcus</i> (n) and <i>Weissella</i>
Gram-negative bacteria	<i>Aeromonas</i> (a, c, g, l, w), <i>Agarivorans</i> , <i>Alteromonas</i> (p, t, v), <i>Bdellovibrio</i> (b), <i>Burkholderia</i> , <i>Citrobacter</i> (b), <i>Enterobacter</i> (f), <i>Neptunomonas</i> (p, t, v), <i>Phaeobacter</i> (n, s), <i>Pseudoalteromonas</i> (n, t, v), <i>Pseudomonas</i> (c, h, n, q), <i>Rhodobacter</i> (n), <i>Rhodopseudomonas</i> , <i>Roseobacter</i> (n), <i>Shewanella</i> (i), <i>Synechococcus</i> (q), <i>Thalassobacter</i> , <i>Vibrio</i> (c, n, q, x) and <i>Zooshikella</i> (l)
Non-bacterial candidates	
Bacteriophage	<i>Myoviridae</i> (k) and <i>Podoviridae</i> (k)
Microalgae (m, o, u)	<i>Dunaliellasalina</i> , <i>D. tertiolecta</i> , <i>Isochrysisgalbana</i> , <i>Navicula</i> , <i>Phaedactylumtricornutum</i> and <i>Tetraselmissuecia</i>
Yeast	<i>Debaryomyceshansenii</i> , <i>Phaffiarhodozyma</i> , <i>Saccharomyces cerevisiae</i> (b, l), <i>S. exiguus</i> and <i>Yarrowialipolytica</i>

Pathogens: a → *A. bestiarum*, b → *A. hydrophila*, c → *A. salmonicida*, d → *Edw. ictaluri*, e → *Edw. tarda*, f → *F. psychrophilum*, g → *Lc. Garvieae*, h → *Photobacteriumdamselae* subsp. *damselae*, i → *Ph. damsela* subsp. *piscicida*, j → *Ps. fluorescens*, k → *Ps. plecoglossicida*, l → *Streptococcus* sp./*St. iniae*, m → *Vibrio* spp., n → *V. anguillarum*, o → *V. campbellii*, p → *V. coralliilyticus*, q → *V. harveyi*, r → *V. ordalii*, s → *V. parahaemolyticus*, t → *V. pectenicida*, u → *V. proteolyticus*, v → *V. splendidus*, w → *V. tubiashii*, x → *Y. ruckeri*.

way, probiotics, both in conditions of invitro and invivo; actively induce the proliferation of B lymphocytes in fish.

Increasing the level of immunoglobulin by probiotic supplementation has been reported in a number of animals including fish [63]. It has been demonstrated that oral administration of the bacteria *Clostridium butyricum* in rainbow trout has reinforced the strength of the fish to vibriosis, increasing phagocytic activity of leukocytes [94]. The use of *Bacillus* sp. (Strain S11) provided protection of the disease by activating both cellular and humoral immune defense in tiger shrimp (*Penaeus monodon*) [89].

A management of mixture of bacterial strain (*Bacillus* and *Vibrio* sp.) positively influenced the growth and survival of young white shrimp and presents a protective effect against the pathogen *V. harveyi* and white spot syndrome virus [8]. This protection is due to the stimulation of the immune system, by increasing the phagocytosis and antibacterial activity. In addition, the administration of a lactic acid bacterium *Lactobacillus rhamnosus* (ATCC 53103) in an amount of 10^5 cfu g^{-1} feed; stimulates the respiratory burst in rainbow trout (*Oncorhynchus mykiss*) [81]. However [10] found only increase in immunoglobulin levels in brown trout (*Salmo trutta*) but not significant level by lactic acid bacteria (LAB) fed groups of probiotics namely *Lactococcus Lactis* sp. *Lactis*, *Lactobacillus sakei* and *Leuconostoc mesenteroides* supplemented at 10 (6) CFU/g diet for a period of 2 weeks.

Therefore, probiotics are beneficial bacteria capable of inhibiting not only pathogens, but also regulate the host immune system [Fig. 1]. Immunomodulation by probiotics are considered as a community effort of the introduced microorganism, host and commensals. The host can detect whether the organism is pathogenic or not through pathogen pattern recognition receptors (PRRs). To identify these recognition receptors, the microbial associated molecular patterns (MAMPs) which are present in both pathogenic and non-pathogenic microorganism. Some MAMPs are lipopolysaccharides (LPS), peptidoglycan, flagellin, and microbial nucleic acids. The binding of MAMPs to PRRs trigger intracellular signaling cascade, urging the release of specific cytokines and

transmit signals to adjacent cells, or to exert anti-viral, pro- or anti-inflammatory exercise effects. The same mechanism of recognition regulates the homeostasis of the commensal of microbiota in the mucosa. Moreover, probiotics can also manipulate the richness and diversity of commensal microbiota [76].

4. Mechanism of immune response

4.1. Phagocytic activity

Phagocytes such as neutrophils and macrophages play an important role in the anti-bacterial defense. These cells ingest and kill bacteria through the production of reactive oxygen species (ROS), including superoxide anion, hydrogen peroxide and hydroxyl radicals in the respiratory burst [34]. Therefore, probiotics can effectively cause phagocytes in host and improve phagocytosis activity by lactic acid bacteria (LAB) group of probiotics; such as *L. rhamnosus*, *Lactobacillus acidophilus* and *Lactobacillus Lactis* which has been observed in several animals. In tilapia (*Oreochromis niloticus*) a two weeks feeding of *L. rhamnosus* significantly stimulates phagocytosis [86]. Similarly, the oral administrations of *C. butyricum* bacteria in *O. mykiss* have also been reported to increase their phagocytosis activity [94]. In rainbow trout, increased activity was identified relative to the control after four weeks, and furthermore, the highest activity against *Vibrio anguillarum* was observed in rainbow trout two weeks after the start of probiotic feeding mode [102].

4.2. Respiratory burst activity

Respiratory burst activity is important innate defense mechanisms of fish. In fish, probiotics have been reported to improve the respiratory burst of phagocytic cells, which play a central role in the protection of non-specific cell [12,69,82]. The two weeks feeding lactic acid bacteria (LAB 4012) significantly improved the respiratory burst of peripheral blood lymphocytes (PBL) from the fed's fish




Fig. 1. Probiotics showing the activity of host immunomodulation. Abbreviations: MAMPs → Microbe associated molecular patterns, PRRs → Pathogen pattern recognition receptors, IEC → Intestinal epithelial cell.

and sparked an effective protection of Cobia (*Rachycentron canadum*) against photobacteriosis challenge (RPS = 74.3) [124]. Several invitro and invivo studies showed a significant increase in respiratory burst activity by various probiotics in many aquatic animals and fish. The activity of head kidney leucocytes of gilthead sea bream (*Sparus aurata*) has additionally been found to improve with the feeding 5×10^7 CFU/ml of heat-inactivated *Lactobacillus delbrueckii subsp. Lactis* and *B. subtilis* under invitro conditions [96]. HK, (Head kidney) respiratory burst activity of leukocytes, is important innate immune parameter, which are widely used as an indicator of immune activity, especially when caused by stimulants [101].

4.3. Lysozyme

Lysozyme is a component of the innate immune system and plays an important role in the defense mechanism because of its anti-cancer, anti-viral and opsonization properties [59]. It is one of the most important bactericidal enzymes of intrinsic immunity and is an indispensable tool for the fish to fight infectious agents [68]. Probiotics are found either singly or in combination, trigger the lysozyme level in bony fish. It is noted that supplementation of *L. rhamnosus* (10^9 CFU g^{-1}) as feed additives for 30 days improved lysozyme activity in rainbow trout [82]. Stimulation of lysozyme activity in rainbow trout was observed after two weeks feeding with *Kocuria sp.* SM1 [102] and *Carnobacterium divergens* B33 [94]. Lysozyme activity significantly increased in the third and fourth week of the twelve diet group. The lysozyme activity of *Catla sp.*, serum was significantly increased ($p < 0.05$) after four weeks of feeding with different probiotic supplemented diets compared to that of the control [27]. In addition to serum lysozyme content probiotic can also increase the lysozyme level in the skin mucosa fish [107,116]. Taoka et al [116] reported significantly high lysozyme level in skin mucosa by supplementing commercial probiotics by water compared to oral supplementation in *O. niloticus*. These results suggest that high lysozyme activity can afford to ward off bacterial diseases in probiotic-supplemented fish.

4.4. Peroxidase and anti-protease activity

The peroxidase is an important enzyme which uses oxidizing radicals to produce hypochlorous acid to kill pathogens. During the oxidative respiratory burst, peroxidase is usually released from the azurophilic granules of neutrophils. Nutritional Supplements of probiotics such as *Bacillus subtilis*, alone or in combination with *N. delbrueckii spp. Lactis* for three weeks resulted in high serum protease activity, but failed to improve the peroxidase activity of the head kidney leukocytes *S. aurata* [95]. Likewise, probiotics such as *Enterococcus faecium* also increase the serum level of peroxidase in *O. niloticus*, if by the water in every four days for forty days supplemented at 1×10^7 CFU/ml [123]. Anti-protease activity in serum, which contains α 1-antiprotease, the α 2-antiprotease and α 2 macroglobulin, inhibits the activity of the proteases used by certain bacteria to invade the host. This activity is generally high and is not affected by immunization or infection [35,72]. Sharifuzzaman and Austin [102] reported significantly higher anti-protease activity in *O. mykiss* within two weeks of the completion of probiotic species *Kocuria*.

4.5. Complement activity

The complement system is the part of the immune system that helps or supplements the ability of phagocytes to combat pathogens by antibody opsonization. It is part of the innate immune system which is not adjustable and does not change over the

lifetime of an individual [58]. However, it can be adjusted by the adaptive immune system and brought into action. Complement, is the main component of the innate humoral immune response, plays an essential role in the modification of the host immune system of the presence of potential pathogens and their clearance. Complement is initiated by one or a combination of the three ways, namely, the classical, alternative and lectin. All three paths merge into a common amplification with C3 and navigate through the terminal path that leads to the formation of the membrane attack complex, which can directly lyses pathogenic cells [17]. According to [115]; serum C3 levels of probiotic treatments were significantly higher than that of the control after 30 days of feeding on *Epinephelus coioides* grouper. Furthermore, C4, a key component of the classical and lectin pathways, compared with the control after thirty days of feeding increased slightly in the probiotic treatments. The probiotic *Pediococcus acidilactici* as a potent food additive improves the level of the alternative complement activity in the serum of the green terror during a 56-day period [78].

4.6. Cytokines

Cytokines are a broad and loose matter of small proteins that are important in cellular signaling. They are released by cells and affect the behavior of other cells, and sometimes producing cell itself. Cytokines include chemokines, interferons, interleukins, lymphokines, tumor necrosis factor [25,26]. A number of probiotics can effectively modulate the production of pro-inflammatory cytokines, such as interleukin-1 (IL-1), IL-6, IL-12, tumor necrosis factor α (TNF- α) and interferon-gamma (IFN- γ) and anti-inflammatory cytokines, such as IL-10 and transforming growth factor β (TGF β) in many organisms [24,80,122]. Probiotics such as *L. rhamnosus*, *E. faecium* and *B. subtilis* are found to up regulate the pro-inflammatory cytokines such as IL-1 β and TGF β in the spleen and head kidney of the *O. mykiss* [83]. The cytokine gene-reaction was performed in an invitro study with head kidney cells of the Japanese pufferfish (*Takifugu rubripes*) that were exposed significantly to improve heat-killed *Lactobacillus* derived from Mongolian dairy products. Thus, the *Lactobacilli* triggered reactions of the antiviral (I-IFN-1 and IFN- γ), the cell-mediated immune regulators (IL-12p35, IL-12p40 and IL-18), pro-inflammatory (IL-1 β , IL-6, IL-17A/F-3, TNF- α and TNF-N) and regulatory (IL-2, IL-7, IL-15, IL-21, IL-10 and TGF- β 1) cytokines [16].

Interestingly, the data pointed *Lb. subsp. paracasei* having greater potential for immunomodulation by *Lb. plantarum* [16]. With real-time polymerase chain reaction (PCR), it was determined that IL-1b, IL-10 and TNF gene expression was significantly up regulated in the head kidney after feeding with *Lb. plantarum* [85]. In a separate study, pro-inflammatory cytokine TNF- α in the Nile tilapia was up regulated after feeding with *Pediococcus acidilactici* [109]. In addition, IL-12 and IFN- γ were induced in olive flounder (*Paralichthys olivaceus*) after feeding with *Lactococcus lactis* [61]. Thus, a large number of microorganisms have been considered for use as probiotics in aquaculture with a mechanism of action involving more immunomodulation, wherein some of sub cellular components act as immunostimulating agents in their own right.

5. Probiotics

The use of immunostimulants may enhance activities in a non-specific defense mechanism, the increase resistance to infectious disease by enhancing innate humoral and cellular defense mechanisms and indirectly cause increased growth of aquatic animals [7]. Probiotics are indigestible food ingredients that selectively and beneficially affect the host by stimulating the growth and/or activity of one or a limited number of bacteria in the colon.

Recruitment prebiotics can significantly alter the colonic microflora increasing the number of special bacteria and thus changing the composition of microflora [46]. Prebiotics are carbohydrates which can be classified according to molecular size or degree of polymerization (number of monosaccharide units) such as monosaccharides, polysaccharides or oligosaccharides. According to the International Union of Pure and Applied Chemistry nomenclature, oligosaccharides are defined as saccharides containing between three and ten meioties sugar [75]. Examples of prebiotics included fructo-oligosaccharide, mannanoligosaccharide, inulin or β -glucan. Despite potential benefits for health and performance as noted in various terrestrial animals, the use of prebiotics in the cultivation of fish and shellfish has been less investigated.

The pattern recognition receptors (PRRs), such as beta (β) glucan receptors dentin-1 receptors expressed in macrophages have direct interactions mediated with the immunomodulatory activity of prebiotics [19]. This type of receptor interaction activates signal transduction molecules, such as NF- κ B, which stimulate the immune cells [125], saccharides may also interact with PRRs format of microbial associated molecular patterns (MAMPs) as teichoic acid, peptidoglycan, glycosylated protein or the capsular polysaccharide of the bacteria, causing an immune response [18].

6. Immunomodulatory effects

Prebiotics act as a growth factor to specific commensal bacteria, that inhibit the adhesion and invasion of pathogenic microorganisms in the colon epithelium by competing for the same glycoconjugates found on the surface of epithelial cells, altering the pH of the colon, preference the barrier function, improvement the production of mucus, production of short chain fatty acids and induce cytokine production. The effect of prebiotics directly enhances the innate immunity, such as;

Phagocytic effect; is the process which has the endocytic ability of intracellular phagocytic leukocytes and an active host defense mechanism in the spleen, kidney, head kidney (HK) or other lymphoid organs. It can be observed by measuring the level of endocytosed zymosan in phagocytes using microscopy or colorimetric detection. Phagocytosis occurs in a series of steps. Microbes are detected by PRRs such as toll like receptors (TLRs); the microbes are devoured by phagosomes; Phagosomes fuse with lysosomes, which contain a variety of proteases; and the microbes are killed by proteolysis [1]. Furthermore antigen processing steps followed by the presentation of antigen to T cells, leading to T cell activation, and consequently the whole immune system. The Japanese flounder (*P. olivaceus*) was fed 5.0 g FOS kg^{-1} for 56 days were analyzed for lysozyme activity, the percentage of phagocytic cells that assumed a marker (phagocytic percent) and phagocytic index [127]. Fructooligosaccharides (FOS) administration significantly improved lysozyme activity, but not the phagocytic percentage or phagocytic index, compared with the control diet group. However, when fed a mixture of FOS and mannanooligosaccharides (MOS), 5.0 g kg^{-1} , the flounder showed a marginal increase in phagocytic activity.

Macrophages; play an important role in linking the innate and adaptive connection of immune system to produce a maximum immune response and kill pathogenic microbes. Macrophages of IFN- γ and their direct interactions between MAMPs are activated on bacteria and PRRs on the host cells. Activated macrophages secrete variety of inflammatory cytokines, such as tumor necrosis factor (TNF), IL-1, IL-12, etc. [1]. The major macrophage activation steps are the cytokines and can be observed by either polymerase chain reaction (PCR) or ELISA. Phosphomonoesterase or acid phosphatase is an enzyme which removes the phosphate group from the phosphorylated molecules. In activated macrophages or dendritic cells, acid phosphatase triggers a drop in the internal pH

of phagolysosomes macrophages, and increases its internal acidity. The higher internal acidity leads to activation of protease, whereby resulting microcidal activity [1].

A respiratory burst or oxidative burst is an indication of oxidative potential of reactive oxygen species such as hydrogen peroxide, superoxide anions and hydroxyl radicals. Reactive oxygen species are produced by activated phagocytes and they are responsible for killing or degrading devours materials, including microbes. Reactive oxygen species are widely used to protect the ability of the host against pathogens [1]. Respiratory bursts of innate immune cells, including blood neutrophils, are measured by using NBT (nitroblue tetrazolium) or MPO (myeloperoxidase) tests. Nile tilapia was given a dietary supplement of inulin 5 g kg^{-1} increased hematocrit NBT activity and lysozyme activity, suggesting that inulin has a stimulating effect on the innate immunity. In the group fed inulin for two months there was a statistically significant change in the NBT activity and lysozyme activity. Furthermore, the survival rate after challenge of *A. hydrophilla* increased in inulin fed group [54].

The complement system in serum is one of the most potent non-effector cell responses of the immune system and may be activated by antigen-specific antibodies, a microbial cell surface, or a lectin. Phenoloxidase can be used for measuring the state of the innate immune system of marine invertebrates, such as, Cray fish, sea cucumber, shrimp or lobster [104] [15,52]. The prophenoloxidase system is modified complement system especially for marine invertebrates comprising tyrosinases, catecholases and laccases. Phenoloxidase is critical for enhancing the antimicrobial activity through a respiratory burst and phagocytosis by opsonization. Red Swamp Cray fish proved significantly increased the activity of Phenoloxidase, superoxide dismutase (SOD) and the expression of immune genes (crustin 1, lysozyme, SOD and prop) when fed dietary FOS 8 and 10 g kg^{-1} [31]. Thus, the dietary supplement of fructooligosaccharides for thirty days is given, increasing the survival period against *Aeromonas hydrophilla*.

Lysozyme is an enzyme degrading peptidoglycan in bacterial walls by hydrolyzing β - [11,56] glycosidic bonds to N-acetylmuramic acid and N-acetylglucosamine. Lysozyme is found in many areas, such as mucus, serum, intestines and eggs of marine animals [5]. Activated macrophages are the primary producers of lysozyme [48]. The effects of dietary fructo-oligosaccharides, including 10, 20, or 30 g kg^{-1} for Caspian roach fry investigated [106]. Immunoglobulin levels of lysozyme ACH50 were fed increasing the activity significantly in the groups 20 and 30 g FOS kg^{-1} . Only lysozyme activity was improved in the group fed 10 g FOS kg^{-1} . Nutritional supplements FOS increases the resistance of the roach on salt stress challenges to complete, regardless of the level, but only the group fed 3% FOS had a significantly higher survival rate.

Antibodies are produced by B lymphocytes and recognize specific microbial antigens. The antibodies neutralize pathogens by binding to their surface antigens and preventing their attachment to their target cells. Antibodies may also facilitate the phagocytosis of antibody-bound pathogens through opsonization and activate the complement system and antibody-dependent cellular cytotoxicity (ADCC). The hematocrit or total hemocyte count reflects the total number of cells in the blood, including red blood cells, white blood cells and platelets. The hematocrit count can be used as a macro-analysis of the immunological status of fish, because the number of immune cells in the immune activation increases blood. For the immune cells in the blood are neutrophils, eosinophils, basophils, monocytes and lymphocytes. Carp fry (11.12 \pm 0.55 g) were given a diet supplemented with immunogen 0, 0.5, 1, 1.5 and 2.5 g kg^{-1} . Immunogen is a commercial product, which contains mannano-oligosaccharides (MOS) and β -glucan [32]. All the complement level of leukocyte counts increased significantly.

Prebiotics have a beneficial effect on the gut-associated lymphoid tissue (GALT). Inulin, Fructooligosaccharides, mannanooligosaccharides, galactooligosaccharides and arabinogalactans are nutritional therapeutic preparations. They are used for optimal bowel function for preferring the proliferation of normal bacterial flora and prevent the ontogenesis of pathogens. In

different species they have been investigated as prebiotics (Table 2), and are used to impede the growth of pathogenic organisms. In fact these Immunosaccharides primarily relieve the function of the phagocytic cells and increased bactericidal activities. Moreover, they also stimulate natural killer cells, complement, lysozyme and antibody response in the host.

Table 2

The following references to the table, showing related research Immunosaccharides, such as; inulin, fructooligosaccharides (FOS), mannanoligosaccharides (MOS), galactooligosaccharides (GOS) and arabinoxylane-oligosaccharides (AXOS), using as immunostimulants in aquaculture.

Prebiotic	Species	Dose, route and duration of administration	Weight	Response	Reference
Inulin	<i>Beluga</i> [Great sturgeon]	Dietary 1.0, 2.0, 3.0% inulin 8 weeks	(16.14 ± 0.38 g)	RBC count →, Total WBC count ↑, MCH → Dose dependent alkaline phosphatase ↓.	[2]
	<i>Gilthead seabream</i>	Dietary 1.0% (10 g kg ⁻¹) inulin 2 and 4 weeks	(50 g)	Serum complement activity ↑, IgM level ↑, Leucocyte phagocytic activity ↑, Leucocyte respiratory burst activity ↑, Protection from <i>P. damsela</i> challenge ↑.	[20,21]
	<i>Hybrid surubim</i>	Dietary 0.5% inulin 15 days	(73.6 ± 19.5 g)	Lactic acid bacteria ↑, <i>Vibrio</i> spp. ↓, <i>Pseudomonas</i> spp. ↓, Total Ig ↑ (restricted to symbiotic treatment), Serum protein or lysozyme →.	[74]
Fructooligosaccharides	<i>Nile tilapia</i>	Dietary 0.5% (5 g kg ⁻¹) inulin 1 and 2 months	(11.00 ± 0.2 g)	Hematocrit ↑, NBT (Superoxide activity) ↑, lysozyme activity ↑, Protection from <i>A. hydrophila</i> challenge ↑.	[54]
	<i>Atlantic salmon</i>	Dietary 1.0% (10 g kg ⁻¹) 4 months	(200.2 ± 0.6 g)	Whole Blood Neutrophil Oxidative Radical Production →, Serum Lysozyme Activity →.	[49]
	<i>Black amur bream</i>	Dietary 0.3 and 0.6%, 8 weeks	(30.5 ± 0.5 g)	ACP ↑, PO ↑, ACH50 ↑, Ig M ↑.	[128]
	<i>Caspian roach</i>	Dietary 1, 2, 3%, 7 weeks	(0.67 ± 0.03 g)	Serum Ig ↑, Lysozyme activity ↑, ACH50 ↑ Resistance to salinity challenge ↑.	[105,107]
	<i>Japanese flounder</i>	Dietary 0.005% (5.0 g kg ⁻¹) 56 days	(21 g)	Lysozyme activity ↑, Phagocytic percentage and index →.	[127]
	<i>Red swamp cray fish</i>	Dietary 0.008, 0.01% (8, 10 g kg ⁻¹) 30 days	(15–17 g)	Immune related genes (crustin1, lysozyme, SOD, and proPO) ↑, Phagocytic activity ↑, SOD ↑, Survival against <i>A. hydrophila</i> ↑	[31]
	<i>Sea cucumber</i>	Dietary 0.25, 0.5%, 8 weeks	(5.06 ± 0.10 g)	0.5% FOS TCC ↑, Phagocytosis ↑, PO ↑.	[129]
	<i>Stellate sturgeon</i>	Dietary 0.4, 0.8, 1.6%, 50 days	(3.72 ± 0.16 g)	No significant immunological improvement observed.	[114]
Nanoligosaccharides	<i>Yellow croaker</i>	Dietary 1, 2%, 11 weeks	(30.16 ± 0.14 g)	1% FOS; serum lysozyme activity ↑.	[3]
	<i>Atlantic salmon</i>	Dietary 0.2, 0.4%, 10 weeks	(7.82 ± 0.68 g)	No statistically significant innate immunity improvement observed.	[87]
	<i>Atlantic salmon</i>	Dietary 1.0% (10 g kg ⁻¹) 4 months	(200.2 ± 0.6 g)	Whole Blood Neutrophil Oxidative Radical Production ↓, Serum Lysozyme Activity ↓	[49]
	<i>Cray fish; common yabby</i>	Dietary 4%, 56 days	(35.14 ± 0.48 g)	THC ↑, GC ↑, Semi-GC ↑, Hyaline cells ↓, Bacteremia ↓	[97,99]
	<i>European sea bass</i>	Dietary 2, 4% 8 weeks	(116 g)	4% MOS head kidney (HK) macrophage phagocytic activity ↑, 2% MOS HK macrophage phagocytic activity ↓, Eosinophilic granulocytes ↑, levels of <i>V. alginolyticus</i> after exposure ↓	[118]
	<i>Japanese flounder</i>	Dietary 0.005% (5.0 g kg ⁻¹) 56 days	(21 g)	Lysozyme activity ↑, Phagocytic percentage and index →	[127]
	<i>Marron</i>	Dietary 0.2, 0.4%/0.05, 0.1, 0.2, 0.4, 0.8%, 30–112 days	(4.44 ± 0.20 g, 10.44 ± 0.20 g, 94 ± 2.17 g)	Survival in <i>V. mimicus</i> challenge ↑, Live transport; GCs ↑, Bacteremia ↓	[97,99, 100]
	<i>Rainbow trout</i>	Dietary 0.2% (2 g kg ⁻¹) 42 days	(30 g)	Antibody titer ↑, Lysozyme activity ↑	[110]
	<i>Rainbow trout</i>	Dietary 0.4% 12 weeks	(13.2 g)	Hemolytic activity ↑, Phagocytic activity ↑, Survival in <i>V. anguillarum</i> challenge ↑	[92]
	<i>Rainbow trout</i>	Dietary 0.25, 0.5% 12 weeks	(36.27 ± 0.42 g)	0.5% MOS Hematocrit ↑, Phagocytic activity ↑, Survival in <i>A. salmonicida</i> challenge ↑	[93]
	<i>Sea cucumber</i>	Dietary 0.1, 0.2% 4 weeks	(3.8 ± 0.2 g)	TCC ↑, Phagocytosis of coelomocytes ↑, Superoxide anion production ↑, SOD activity ↑, Survival in <i>V. splendidus</i> challenge ↑	[50]
	<i>Sea cucumber</i>	Dietary 0.002% (2 g kg ⁻¹) 4 weeks	(6.80 ± 0.30 g)	TCC ↑, Superoxide anion production ↑ at 15 days	[9]
	<i>Tropical spiny lobster</i>	Dietary 0.4%, 8 weeks	(1.28 ± 0.01 g)	THC ↑, GC count ↑, Bacteremia ↓	[98]
	<i>Common carp</i>	Dietary (Immunogen®) 0.0005, 0.001, 0.0015, 0.0025%, (0.5, 1, 1.5, 2.5 g kg ⁻¹) 8 weeks	(11.12 ± 0.55 g)	Leucocyte counts ↑, Dose dependent survival rate in <i>A. hydrophila</i> challenge ↑	[32]
Galactooligosaccharides	<i>Atlantic salmon</i>	Dietary 1.0% (10 g/kg) 4 months	(200.2 ± 0.6 g)	Whole Blood Neutrophil Oxidative Radical Production →, Serum Lysozyme Activity →	[49]
	<i>Red drum</i>	Dietary 1% 8 weeks	(7 g)	Lysozyme level ↑, NBT reduction level →, Microvilli heights in the pyloric caeca, proximal and mid intestine ↑	[130]
Arabinoxylan	<i>Siberian sturgeon</i>	Dietary 2% 12 weeks	(25.9 ± 0.9 g)	Phagocytic activity ↑, ACH50 ↑ (AXO-32-0.30 only), Serum peroxidase content ↑ (AXO-32-0.30 only)	[45]
	<i>Siberian sturgeon</i>	Dietary 2% 28 days	(48.4 ± 1.4 g)	Phagocytic activity ↑, Respiratory burst activity ↓	[44]

Abbreviations: ACP-plasma alkaline phosphatase, GC-granular cells, PO-phenoloxidase, SOD-superoxide dismutase, TCC-total coelomocyte count, THC-total hemocyte count, ACH50-alternative complement activity, NBT-nitroblue tetrazolium. ↑-increase, →-no change, ↓-decrease.

7. Conclusion

Fish has become one of the major aquaculture industries. In natural conditions, fish consume a larger variety of food compared to cultured fish. An indicator in this regard may be farmed fish to reduce the intestinal microbial community diversity [29]. Many dietary supplements such as probiotics, prebiotics, vitamins or plant crude extracts can compensate for this problem, resulting in prosperous result in aquaculture [42]. However, most of probiotics and prebiotics can exert immunomodulatory effects in aquatic animals. Therefore, in order to improve the probiotics and prebiotics for aquaculture, further work needs to be carried out in the exact nature of humoral innate or cellular immune response and to determine the protective probiotic life of bacteria. Finally, this paper proposes a long-term perspective of natural dietary supplement, such as probiotics and prebiotics that can be used as strategic control of diseases in aquaculture.

Acknowledgment

This work was supported by NSFC (No. 81273386).

References

- [1] A.K. Abbas, A.H. Lichtman, S. Pillai, Cellular and Molecular Immunology, seventh ed., Saunders Elsevier Philadelphia, PA, USA, 2012.
- [2] Ehsan Ahmdifar, Raza Akrami, Afshin Ghelichi, et al., Effects of different dietary prebiotic inulin levels on blood serum enzymes, hematologic, and biochemical parameters of great sturgeon (*Huso huso*) juveniles, Comp. Clin. Pathol. 20 (5) (2011) 447–451.
- [3] Reza Akrami, Yousef Iri, Hosseinali Khoshbavar Rostami, et al., Effect of dietary supplementation of fructooligosaccharide (FOS) on growth performance, survival, *Lactobacillus* bacterial population and hemato-immunological parameters of stellate sturgeon (*Acipenser stellatus*) juvenile, Fish Shellfish Immunol. 35 (4) (2013) 1235–1239.
- [4] M.A. Al-Dohail, H. Roshada, M. Aliyu-Paiko, Effects of the probiotic, *Lactobacillus acidophilus*, on the growth performance, haematology parameters and immunoglobulin concentration in African Catfish (*Clarias gariepinus*, Burchell 1822) fingerling, Aqua. Res. 40 (2009) 1642–1652.
- [5] J.B. Alexander, G.A. Ingram, Noncellular nonspecific defense mechanisms of fish, Annu. Rev. Fish Dis. 2 (1992) 249–279. Published: 1992.
- [6] S.M. Aly, Y.A.G. Ahmed, A.A.A. Ghareeb, M.F. Mohamed, Studies on *Bacillus subtilis* and *Lactobacillus acidophilus*, as potential probiotics, on the immune response and resistance of Tilapia nilotica (*Oreochromis niloticus*) to challenge infections, Fish Shellfish Immunol. 25 (1–2) (2008) 128–136.
- [7] A.S. Amlashi, B. Falahatkar, M. Sattari, M.H.T. Gilani, Effect of dietary vitamin E on growth, muscle composition, hematological and immunological parameters of sub-yearling beluga *Huso huso* L, Fish Shellfish Immunol. 30 (3) (2011) 807–814.
- [8] S.P. Antony, I.S.B. Singh, R.M. Jose, P.R.A. Kumar, R. Philip, Antimicrobial peptide gene expression in tiger shrimp, *Penaeus monodon* in response to gram-positive bacterial probiotics and white spot virus challenge, Aquaculture 316 (1–4) (2011) 6–12.
- [9] N. Bai, W. Zhang, K. Mai, M. Gu, W. Xu, Effects of continuous and alternative administration of α -glucan and mannan-oligosaccharide on the growth, immunity and resistance against *Vibrio splendidus* of sea cucumber *Apostichopus japonicus* (Selenka), Aquacult. Res. 44 (2013) 1613–1624.
- [10] J.L. Balcázar, Evaluation of Probiotic Bacterial Strains in *Litopenaeus vannamei*. Final Report, National Center for Marine and Aquaculture Research, Guayaquil, Ecuador, 2003.
- [11] J.L. Balcázar, I. de-Blas, I. Ruiz-Zazuela, D. Cunningham, D. Vandrell, J.L. Muzquiz, The role of probiotics in aquaculture, Veterinary Microbiol. 114 (3–4) (2006a) 173–186.
- [12] J.L. Balcázar, I. de-Blas, I. Ruiz-Zazuela, D. Vandrell, O. Girones, J.L. Muzquiz, Enhancement of the immune response and protection induced by probiotic lactic acid bacteria against furunculosis in rainbow trout (*Oncorhynchus mykiss*), FEMS Immunol. Med. Microbiol. 51 (2007) 85–93.
- [13] J.L. Balcázar, D. Vendrell, I. de-Blas, I. Ruiz-Zazuela, O. Girones, J.L. Muzquiz, Immune modulation by probiotic strains: quantification of phagocytosis of *Aeromonas salmonicida* by leukocytes isolated from gut of rainbow trout (*Oncorhynchus mykiss*) using a radiolabelling assay, Comp. Immunol. Microbiol. Infect. Dis. 29 (2006b) 5–43.
- [14] S. Batista, S.T. Tapia-Paniagua, M.A. Morinigo, J.A. Nunez-Diaz, J.F.M. Goncalves, R. Barros, Expression of immune response genes in sole (*Solea senegalensis*, Kaup 1858) induced by dietary probiotic supplementation following exposure to *Photobacterium damsela* subsp. piscicida, Fish Shellfish Immunol. 34 (2013) 1638–1639.
- [15] G. Beck, G.S. Habicht, Immunity and the invertebrates, Sci. Am. 275 (1996 November) 60–66.
- [16] G. Biswas, H. Korenaga, R. Nagamine, H. Takayama, S. Kawahara, S. Takeda, Y. Kikuchi, B. Dashnyam, T. Kono, M. Sakai, Cytokine responses in the Japanese pufferfish (*Takifugu rubripes*) head kidney cells induced with heat-killed probiotics isolated from the Mongolian dairy products, Fish Shellfish Immunol. 34 (5) (2013) 1170–1177.
- [17] H. Boshra, J. Li, J.O. Sunyer, Recent advances on the complement system of teleost fish, Fish Shellfish Immunol. 20 (2) (2006) 239–262.
- [18] P.A. Bron, B.P. Van, M. Kleerebezem, Emerging molecular insights into the interaction between probiotics and the host intestinal mucosa, Nat. Rev. Microbiol. 10 (1) (2012), 66–U90.
- [19] G.D. Brown, P.R. Taylor, D.M. Reid, J.A. Willment, D.L. Williams, L. Martinez-Pomares, Dectin-1 is a major β -glucan receptor on macrophages, J. Exp. Med. 196 (2002) 407–412.
- [20] R. Cerezuola, A. Cuesta, J. Meseguer, M.A. Esteban, Effects of inulin on gilt-head seabream (*Sparus aurata* L.) innate immune parameters, Fish Shellfish Immunol. 24 (5) (2008) 663–668.
- [21] R. Cerezuola, F.A. Guardiola, J. Meseguer, M.A. Esteban, Increases in immune parameters by inulin and *Bacillus subtilis* dietary administration to gilt-head seabream (*Sparus aurata* L.) did not correlate with disease resistance to *Photobacterium damsela*, Fish. Shellfish Immunol. 32 (2012) 1032–1040.
- [22] M. Chabrillon, R.M. Rico, S. Arijo, P. Diaz-Rosales, M.C. Balebona, M.A. Morinigo, Interactions of microorganisms isolated from gilt-head sea bream, *Sparus aurata* L., on *Vibrio harveyi*, a pathogen of farmed *Senegalese sole*, *Solea senegalensis* (Kaup), J. Fish. Dis. 28 (2005) 531–537.
- [23] H.R. Christensen, H. Frokiaer, J.J. Pestka, Lactobacilli differentially modulate expression of cytokines and maturation surface markers in murine dendritic cells, J. Immunol. 168 (2002) 171–178.
- [24] Cytokine in John Lackie, A Dictionary of Biomedicine, Oxford University Press, 2010. ISBN 9780199549351.
- [25] Cytokine in Stedman's Medical Dictionary, Wolters Kluwer Health, 28th ed., Lippincott, Williams & Wilkins, 2006.
- [26] Anushree Das, N.A. Khriezhato, C.B. Supratim, K.A. Dibyendu, Effects of potential probiotic *Bacillus amyloliquefaciens* FPTB16 on systemic and cutaneous mucosal immune responses and disease resistance of catla (*Catla catla*), Fish Shellfish Immunol. 35 (5) (2013) 1547–1553.
- [27] S.A. Denev, Ecological Alternatives of Antibiotic Growth Promoters in the Animal Husbandry and Aquaculture, DSc. Thesis, Department of Biochemistry Microbiology, Trakia University, Stara Zagora, Bulgaria, 2008.
- [28] A.K.S. Dhanasiri, L. Brunvold, M.F. Brinchmann, K. Korenes, Ø. Berigh, V. Kiron, Changes in the intestinal microbiota of wild Atlantic cod *Gadus morhua* L. upon captive rearing, Microb. Ecol. 61 (2011) 20–30.
- [29] A. Dimitroglou, D.L. Merrifield, O. Carnevali, S. Picchiatti, M.A. Avella, C.L. Daniels, D. Güroy, S. Davies, Microbial manipulations to improve fish health and production — a Mediterranean perspective, Fish Shellfish Immunol. 30 (2011) 1–16.
- [30] C. Dong, J. Wang, Immunostimulatory effects of dietary fructooligosaccharides on red swamp crayfish, *Procambarus clarkii* (Girard), Aquacult. Res. 44 (2013) 1416–1424.
- [31] G.H. Ebrahimi, H. Ouraji, M.K. Khalesi, M. Sudagar, A. Barari, M.Z. Dangesaraki, Effects of a prebiotic, Immunogen®, on feed utilization, body composition, immunity and resistance to *Aeromonas hydrophila* infection in the common carp *Cyprinus carpio* (Linnaeus) fingerlings, J. Anim. Physiol. Anim. Nutr. 96 (2012) 591–599.
- [32] A.Y. El-Dakar, S.M. Shalaby, I.P. Saoud, Assessing the use of a dietary probiotic/prebiotic as an enhancer of spine foot rabbit fish *Siganus rivulatus* survival and growth, Aqua. Nutr. 13 (2007) 407–412.
- [33] A.E. Ellis, Immunity to bacteria in fish, Fish Shellfish Immunol. 9 (1999) 291–308.
- [34] A.E. Ellis, Innate host defense mechanisms of fish against viruses and bacteria, Dev. Comp. Immunol. 25 (2001) 827–839.
- [35] FAO, Health and Nutritional Properties of Probiotics in Food Including Powder Milk with Live Lactic Acid Bacteria, 2001.
- [36] FAO/WHO, Expert Consultation Report on Evaluation of Health and Nutritional Properties of Probiotics in Food Including Powder Milk with Live Lactic Acid Bacteria, 2001.
- [37] L.J. Fooks, R. Fuller, G.R. Gibson, Probiotics, probiotics and human gut microbiology, Int. Dairy J. 9 (1999) 53–61.
- [38] R. Fuller, A review, probiotics in man and animals, J. Appl. Bacteriol. 66 (1987) 365–378.
- [39] R. Fuller, A review: probiotics in man and animals, J. Appl. Bacteriol. 66 (1989) 365–378.
- [40] C.M. Galdeano, G. Perdigon, The probiotic bacterium *Lactobacillus casei* induces activation of the gut mucosal immune system through innate immunity, Clin. Vacc. Immunol. 13 (2006) 9–26.
- [41] C. Galli, P.C. Calder, Effects of fat and fatty acid intake on inflammatory and immune responses. A critical review, Annu. Rev. Metab. 55 (2009) 123–139.
- [42] F.J. Gatesoupe, The use of probiotics in aquaculture, Aquaculture 180 (1999) 147–165.
- [43] Z. GERALYLOU, C. Souffreau, E. Rurangwa, L. De Meester, C.M. Courtin, J.A. Delcort, Effects of dietary arabinosyl-oligosaccharides (AXOS) and endogenous probiotics on the growth performance, non-specific immunity and gut microbiota of juvenile Siberian sturgeon (*Acipenser baerii*), Fish. Shellfish Immunol. 35 (2013) 766–775.

- [45] Z. GERALYLOU, C. Souffreau, E. Rurangwa, S. D'Hondt, L. Callewaert, C.M. Courtin, Effects of arabinoxylan-oligosaccharides (AXOS) on juvenile Siberian sturgeon (*Acipenser baerii*) performance, immune responses and gastrointestinal microbial community, *Fish. Shellfish Immunol.* 33 (2012) 718–724.
- [46] G.R. Gibson, M.B. Roberfroid, Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics, *J. Nutr.* 125 (1995) 1401–1412.
- [47] M.R. Gismondo, L. Drago, A. Lombardi, Review of probiotics available to modify gastrointestinal flora, *Int. J. Antimicrob. Agents* 12 (1999) 287–292.
- [48] R. Goethe, L. Phi-van, Posttranscriptional lipopolysaccharide regulation of the lysozyme gene at processing of the primary transcript in myelomonocytic HD11 cells, *J. Immunol.* 160 (1998) 4970–4978.
- [49] B. Grisdale-Helland, S.J. Helland, D.M. Gatlin, The effects of dietary supplementation with mannanoligosaccharide, fructooligosaccharide or galactooligosaccharide on the growth and feed utilization of Atlantic salmon (*Salmo salar* L.), *Aquaculture* 283 (2008) 163–167.
- [50] M. Gu, H. Ma, K. Mai, W. Zhang, N. Bai, X. Wang, Effects of dietary β -glucan, mannan oligosaccharide and their combinations on growth performance, immunity and resistance against *Vibrio splendidus* of sea cucumber, *Apostichopus japonicus*, *Fish. Shellfish Immunol.* 31 (2011) 303–309.
- [51] M. Gullian, F. Thompson, J. Rodriguez, Selection of probiotic bacteria and study of their immunostimulatory effect in *Penaeus vannamei*, *Aquaculture* 233 (2004) 1–14.
- [52] C. Hellio, A. Bado-Nilles, B. Gagnaire, T. Renault, H. Thomas-Guyon, Demonstration of a true phenoloxidase activity and activation of a ProPO cascade in pacific oyster, *Crassostrea gigas* (Thunberg) in vitro, *Fish. Shellfish Immunol.* 22 (2007) 433–440.
- [53] J.-B. Huang, Y.-C. Wu, S.-C. Chi, Dietary supplementation of *Pediococcus pentosaceus* enhances innate immunity, physiological health and resistance to *Vibrio anguillarum* in orange-spotted grouper (*Epinephelus coioides*), *Fish. Shellfish Immunol.* 39 (2014) 196–205.
- [54] M.D. Ibrahim, M. Fathi, S. Meslhy, A.M.A. El-Aty, Effect of dietary supplementation of inulin and vitamin C on the growth hematology, innate immunity and resistance of Nile tilapia (*Oreochromis niloticus*), *Fish. Shellfish Immunol.* 29 (2010) 241–246.
- [55] A. Irianto, B. Austin, Probiotics in aquaculture, *J. Fish. Dis.* 25 (2002a) 633–642.
- [56] A. Irianto, B. Austin, Use of probiotics to control furunculosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum), *J. Fish. Dis.* 25 (2002b) 333–342.
- [57] A. Irianto, B. Austin, Use of dead probiotic cells to control furunculosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum), *J. Fish. Dis.* 26 (2003) 59–62.
- [58] C.A. Janeway Jr., P. Travers, M. Walport, The Complement System and Innate Immunity, Immunobiology: The Immune System in Health and Disease, Garland Science, New York, 2001. Retrieved 25 February 2013.
- [59] P. Jøllès, J. Jøllès, What is new in lysozyme research? Always a model system, today as yesterday, *Mol. Cell Biochem.* 63 (1984) 156–189.
- [60] B. Kazun, K. Kazun, Probiotics in aquaculture, *Med. Weter.* 70 (2014) 25–29.
- [61] D. Kim, B.R. Beck, S.-B. Heo, J. Kim, H.B. Kim, S.-M. Lee, Y. Kim, S.Y. Oh, K. Lee, H. Do, *Lactococcus lactis* BFE920 activates the innate immune system of olive flounder (*Paralichthys olivaceus*), resulting in protection against Streptococcus iniae infection and enhancing feed efficiency and weight gain in large scale field-studies, *Fish. Shellfish Immunol.* 35 (2013) 1585–1590.
- [62] D.H. Kim, B. Austin, Cytokine expression in leucocytes and gut cells of rainbow trout, *Oncorhynchus mykiss* Walbaum, induced by probiotics, *Vet. Immunol. Immunopathol.* 114 (2006a) 297–304.
- [63] D.H. Kim, B. Austin, Innate immune responses in rainbow trout (*Oncorhynchus mykiss*, Walbaum) induced by probiotics, *Fish. Shellfish Immunol.* 21 (2006b) 513–524.
- [64] A. Kocher, The potential for immunosaccharides to maximize growth performance: a review of six published meta-analyses on bio-mos, in: L.A. Tucker, J.A. Taylor-Pickard (Eds.), *Interfacing Immunity, Gut Health and Performance*, Nottingham University Press, 2004, pp. 107–116.
- [65] T.L. Korkea-Aho, A. Papadopolou, J. Heikkinen, A. Von wright, A. Adams, B. Austin, K.D. Thompson, *Pseudomonas* M162 confers protection against rainbow trout fry syndrome, *J. Appl. Microbiol.* 113 (2012) 24–35.
- [66] R. Kumar, S.C. Mukherjee, R. Ranjan, S.K. Nayak, Enhanced innate immune parameters in *Labeo rohita* (Ham.) following oral administration of *Bacillus subtilis*, *Fish. Shellfish Immunol.* 24 (2008) 168–172.
- [67] C.C. Lazado, C.M.A. Caipang, M.F. Brinchmann, V. Kiron, In vitro adherence of two candidate probiotics from Atlantic cod and their interference with the adhesion of two pathogenic bacteria, *Vet. Microbiol.* 148 (2011) 252–259.
- [68] G.J.H. Lindsay, The significance of chitinolytic enzymes and lysozyme in rainbow trout (*Salmo gairdneri*) defence, *Aquaculture* 51 (1986) 169–173.
- [69] C.H. Liu, C.H. Chiu, S.W. Wang, W. Chen, Dietary administration of the probiotic, *Bacillus subtilis* E20, enhances the growth, innate immunity response, and disease resistance of the grouper, *Epinephelus coioides*, *Fish. Shellfish Immunol.* 33 (2012) 699–706.
- [70] I.E. Luis-Villasenor, M.E. Macias-Rodriguez, B. Gomez-Gil, F. Ascencio-Valle, A.I. Campa-Cordova, Beneficial effects of four *Bacillus* strains on the larval cultivation of *Litopenaeus vannamei*, *Aquaculture* 321 (2011) 136–144.
- [72] B. Magnadoóttir, B.K. Gudmundsdóttir, S. Lange, A. Steinarsson, M. Oddgeirsson, T. Bowden, Immunostimulation of larvae and juveniles of cod, *Gadus morhua* L., *J. Fish. Dis.* 29 (2006) 147–155.
- [73] A. Mahdhi, F. Kamoun, C. Messina, A. Santulli, A. Bakhrouf, Probiotic properties of *Brevibacillus brevis* and its influence on sea bass (*Dicentrarchus labrax*) larval rearing, *Afr. J. Microbiol. Res.* 6 (2012) 6487–6495.
- [74] J.L.P. Mourino, V.F. Do Nascimento, A.B. Jatobá, B.C. Da Silva, G.F.A. Jesus, W.Q. Seiffert, Effect of dietary supplementation of inulin and W. cibaria on haemato-immunological parameters of hybrid surubim (*Pseudoplatystoma* sp.), *Aquacult. Nutr.* 18 (2012) 73–80.
- [75] S.I. Mussatto, I.M. Mancilha, Non-digestible oligosaccharides: a review, *Carbohydr. Polym.* 68 (2007) 587–597.
- [76] S.K. Nayak, Probiotics and immunity: a fish perspectives, *Fish. Shellfish Immunol.* 29 (2010) 2–14.
- [77] S.K. Nayak, P. Swain, S.C. Mukherjee, Effect of dietary supplementation of probiotic and vitamin C on the immune response of Indian major carp, *Labeo rohita* (Ham), *Fish. Shellfish Immunol.* 23 (2007) 892–896.
- [78] A. Neissi, G. Rafiee, M. Nematollahi, O. Safari, The effect of *Pediococcus acidilactici* bacteria used as probiotic supplement on the growth and non-specific immune responses of green terror, *Aequidens rivulatus*, *Fish. Shellfish Immunol.* 35 (2013) 1976–1980.
- [79] A. Newaj-Fyzul, A.H. Al-Harbi, B. Austin, Review: developments in the use of probiotics for disease control in aquaculture, *Aquaculture* 431 (2014) 1–11.
- [80] L.E.M. Niers, H.M. Timmerman, G.T. Rijkers, G.M. van Bleek, N.O.P. van Uden, E.F. Knol, Identification of strong interleukin-10 inducing lactic acid bacteria which down regulate T helper type 2 cytokines, *Clin. Exp. Allergy* 35 (2005) 1481–1489.
- [81] S. Nikoskelainen, A.C. Ouwehand, G. Bylund, S. Salminen, E. Lilius, Immune enhancement in rainbow trout (*Oncorhynchus mykiss*) by potential probiotic bacteria (*Lactobacillus rhamnosus*), *Fish. Shellfish Immunol.* 15 (2003) 43–52.
- [82] A. Panigrahi, V. Kiron, T. Kobayashi, J. Puangkaew, S. Satoh, H. Sugita, Immune responses in rainbow trout *Oncorhynchus mykiss* induced by a potential probiotic bacteria *Lactobacillus rhamnosus* JCM 1136, *Vet. Immunol. Immunopathol.* 102 (2004) 379–388.
- [83] A. Panigrahi, V. Kiron, S. Satoh, I. Hirono, T. Kobayashi, H. Sugita, Immune modulation and expression of cytokine genes in rainbow trout *Oncorhynchus mykiss* upon probiotic feeding, *Dev. Comp. Immunol.* 31 (2007) 372–382.
- [84] R.B. Parker, Probiotics, the other half of the antibiotic story, *Anim. Nutr. Health* 29 (1974) 4–8.
- [85] T. Pérez-Sánchez, J.L. Balcázar, D.L. Merrifield, O. Carnevali, G. Gioacchini, I. de Blas, I. Ruiz-Zarzuola, Expression of immune-related genes in rainbow trout (*Oncorhynchus mykiss*) induced by probiotic bacteria during *Lactococcus garvieae* infection, *Fish. Shellfish Immunol.* 31 (2011) 196–201.
- [86] N. Pirarat, T. Kobayashi, T. Katagiri, M. Maita, M. Endo, Protective effects and mechanisms of a probiotic bacterium *Lactobacillus rhamnosus* against experimental *Edwardsiella tarda* infection in tilapia (*Oreochromis niloticus*), *Vet. Immunol. Immunopathol.* 113 (2006) 339–347.
- [87] Qinghui Ai, Hougao Xu, Kangsen Mai, Wei Xu, Jun Wang, Wenbing Zhang, Effects of dietary supplementation of *Bacillus subtilis* and fructooligosaccharide on growth performance, survival, non-specific immune response and disease resistance of juvenile large yellow croaker, *Larimichthys crocea*, *Aquaculture* 317 (1–4) (2011) 155–161.
- [88] K.M.M. Rahiman, Y. Jesmi, A.P. Thomas, A.A.M. Hatha, Probiotic effect of *Bacillus* NL110 and *Vibrio* NE17 on the survival, growth performance and immune response of *Macrobrachium rosenbergii* (de Man), *Aquac. Res.* 41 (2010) 120–134.
- [89] S. Rengpipat, S. Rukpratanporn, S. Piyatiratitivorakul, P. Menasaveta, Immunity enhancement in black tiger shrimp (*Penaeus monodon*) by a probiotic bacterium (*Bacillus* S11), *Aquaculture* 191 (2000) 271–288.
- [90] M.T. Ridha, I.S. Azad, Preliminary evaluation of growth performance and immune response of Nile tilapia *Oreochromis niloticus* supplemented with two putative probiotic bacteria, *Aquac. Res.* 43 (2012) 843–852.
- [91] M.B. Roberfroid, Introducing inulin-type fructans, *Brit. J. Nutr.* 93 (2005) 13–26.
- [92] U. Rodrigues-Estrada, S. Satoh, Y. Haga, H. Fushimi, J. Sweetman, Effects of single and combined supplementation of *Enterococcus faecalis*, mannan oligosaccharide and polyhydroxybutyric acid on growth performance and immune response of rainbow trout *Oncorhynchus mykiss*, *Aquacult. Sci.* 54 (2009) 609–617.
- [93] U. Rodrigues-Estrada, S. Satoh, Y. Haga, H. Fushimi, J. Sweetman, Effects of inactivated *Enterococcus faecalis* and mannan oligosaccharide and their combination on growth, immunity, and disease protection in rainbow trout, *N. Am. J. Aquacult.* 75 (2013) 416–428.
- [94] M. Sakai, T. Yoshida, S. Astuta, M. Kobayashi, Enhancement of resistance to vibriosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum) by oral administration of *Clostridium butyricum* bacteria, *J. Fish. Dis.* 18 (1995) 187–190.
- [95] I. Salinas, L. Abelli, F. Bertoni, S. Picchietti, A. Roque, D. Furones, Monospecies and multispecies probiotic formulations produce different systemic and local immunostimulatory effects in the gilthead seabream (*Sparus aurata* L.), *Fish. Shellfish Immunol.* 25 (2008) 114–123.
- [96] I. Salinas, P. Diaz-Rosales, A. Cuesta, J. Meseguer, M. Chabrillon, M.A. Morinigo, Effect of heat-inactivated fish and non-fish derived probiotics on the innate immune parameters of a teleost fish (*Sparus aurata* L.), *Vet. Immunol. Immunopathol.* 111 (2006) 279–286.
- [97] H.M. Sang, R. Fotedakar, K. Filer, Effects of dietary mannan oligosaccharide on the survival, growth, immunity and digestive enzyme activity of freshwater crayfish, *Cherax destructor* Clark (1936), *Aquacult. Nutr.* 17 (2011b) 629–635.

- [98] H.M. Sang, R. Fotadar, Effects of mannan oligosaccharide dietary supplementation on performances of the tropical spiny lobsters juvenile (*Panulirus ornatus*, Fabricius 1798), Fish. Shellfish Immunol. 28 (2010) 483–489.
- [99] H.M. Sang, R. Fotadar, K. Filer, Effects of dietary mannan oligosaccharide on survival, growth, physiological condition, and immunological responses of marron, *Cherax tenuimanus* (Smith 1912), J. World Aquacult Soc. 42 (2011a) 230–241.
- [100] H.M. Sang, L.T. Ky, R. Fotadar, Dietary supplementation of mannan oligosaccharide improves the immune responses and survival of marron, *Cherax tenuimanus* (Smith, 1912) when challenged with different stressors, Fish. Shellfish Immunol. 27 (2009) 341–348.
- [101] C.J. Secombes, Isolation of salmonid macrophages and analysis of their killing activity, in: J.S. Stolen, T.C. Fletcher, D.P. Anderson, B.S. Robertson, W.B. van Muiswinkel (Eds.), Techniques in Fish Immunology, vol. I, SOS Publications, Fair Haven, 1990, pp. 137–154.
- [102] S.M. Sharifuzzaman, B. Austin, Influence of probiotic feeding duration on disease resistance and immune parameters in rainbow trout, Fish. Shellfish Immunol. 27 (2009) 440–445.
- [103] J. Skjermo, T.R. Størseth, K. Hansen, Å. Handa, G. Øie, Evaluation of b-(1/3, 1/6)-glucans and high-M alginate used as immunostimulatory dietary supplement during first feeding and weaning of Atlantic cod (*Gadus morhua* L.), Aquaculture 261 (2006) 1088–1101.
- [104] V.J. Smith, K.A. Soderhall, Comparison of phenoloxidase activity in the blood of marine invertebrates, Dev. Comp. Immunol. 15 (1991) 251–261.
- [105] N. Soleimani, S.H. Hoseinifar, D.L. Merrifield, M. Barati, Z.H. Abadi, Dietary supplementation of fructooligosaccharide (FOS) improves the innate immune response, stress resistance, digestive enzyme activities and growth performance of Caspian roach (*Rutilus rutilus*) fry, Fish. Shellfish Immunol. 32 (2012a) 316–321.
- [106] N. Soleimani, S.H. Hoseinifar, D.L. Merrifield, M. Barati, Z.H. Abadi, Dietary supplementation of fructooligosaccharide (FOS) improves the innate immune response, stress resistance, digestive enzyme activities and growth performance of Caspian roach (*Rutilus rutilus*) fry, Fish. Shellfish Immunol. 32 (2012b) 316–321.
- [107] Z. Song, T. Wu, L. Cai, L. Zhang, X. Zheng, Effects of dietary supplementation with *Clostridium butyricum* on the growth performance and humoral immune response in *Miichthys miiuy*, J. Zhejiang Univ. Sci. B 7 (2006) 596–602.
- [108] L. Sorroza, D. Padilla, F. Acosta, L. Roman, V. Grasso, J. Vega, F. Real, Characterization of the probiotic strain *Vagococcus fluvialis* in the protection of European sea bass (*Dicentrarchus labrax*) against vibriosis by *Vibrio anguillarum*, Vet. Microbiol. 155 (2012) 369–373.
- [109] B.T. Standen, M.D. Rawling, S.J. Davies, M. Castex, A. Foey, G. Gioacchini, O. Carnevali, D.L. Merrifield, Probiotic *Pediococcus acidilactici* moderates both localized intestinal- and peripheral-immunity in tilapia (*Oreochromis niloticus*), Fish Shellfish Immunol. 35 (2013) 1097–1104.
- [110] Y. Staykov, P. Spring, S. Denev, J. Sweetman, Effect of a mannan oligosaccharide on the growth performance and immune status of rainbow trout (*Oncorhynchus mykiss*), Aquacult Int. 15 (2007) 153–161.
- [111] Y. Sugimura, T. Hagi, T. Hoshino, Correlation between in vitro mucin adhesion and the in vivo colonization ability of lactic acid bacteria: screening of new candidate carp probiotics, Biosci. Biotechnol. Biochem. 75 (2011) 511–515.
- [112] H. Sugita, C. Miyajima, H. Deguchi, The vitamin B12-producing ability of the intestinal microflora of freshwater fish, Aquaculture 92 (1991) 267–276.
- [113] H. Sugita, J. Takahashi, H. Deguchi, Production and consumption of biotin by the intestinal microflora of cultured freshwater fishes, Biosci. Biotechnol. Biochem. 56 (1992) 1678–1679.
- [114] Y. Sun, Z. Wen, X. Li, N. Meng, R. Mi, Y. Li, Dietary supplement of fructooligosaccharides and *Bacillus subtilis* enhances the growth rate and disease resistance of the sea cucumber *Apostichopus japonicus* (Selenka), Aquacult Res. 43 (2012) 1328–1334.
- [115] Y.Z. Sun, H.L. Yang, R.L. Ma, W.Y. Lin, Probiotic applications of two dominant gut *Bacillus* strains with antagonistic activity improved the growth performance and immune responses of grouper *Epinephelus coioides*, Fish. Shellfish Immunol. 29 (2010) 803–809.
- [116] Y. Taoka, H. Maeda, J.Y. Jo, S.M. Kim, S. Park, T. Yoshikawa, Use of live and dead probiotic cells in tilapia *Oreochromis niloticus*, Fish. Sci. 72 (2006) 55–66.
- [117] K.I. ten Doeschate, V.E. Coyne, Improved growth rate in farmed Halibutids through probiotic treatment, Aquaculture 284 (2008) 174–179.
- [118] S. Torrecillas, A. Makol, T. Benítez-Santana, M.J. Caballero, D. Montero, J. Sweetman, Reduced gut bacterial translocation in European sea bass (*Dicentrarchus labrax*) fed mannan oligosaccharides (MOS), Fish. Shellfish Immunol. 30 (2011) 674–681.
- [119] L. Verschuere, G. Rombaut, P. Sorgeloos, W. Verstraete, Probiotic bacteria as biological control agents in aquaculture, Microbiol. Mol. Biol. Rev. 64 (2000) 651–671.
- [120] N.G. Vine, W.D. Leukes, H. Kaiser, Probiotics in marine larviculture, FEMS Microbiol. Rev. 30 (2006) 404–427.
- [121] N.G. Vine, W.D. Leukes, H. Kaiser, S. Daya, J. Baxter, T. Hecht, Competition for attachment of aquaculture candidate probiotic and pathogenic bacteria on fish intestinal mucus, J. Fish. Dis. 27 (2004) 319–326.
- [122] T. von der Weid, C. Bulliard, E.J. Schiffrin, Induction by a lactic acid bacterium of a population of CD4⁺ T cells with low proliferative capacity that produces transforming growth factor b and interleukin-10, Clin. Diagn. Lab. Immunol. 8 (2001) 695–701.
- [123] Y.B. Wang, Z.Q. Tian, J.T. Yao, W.F. Li, Effect of probiotics, *Enterococcus faecium*, on tilapia (*Oreochromis niloticus*) growth performance and immune response, Aquaculture 277 (2008) 203–207.
- [124] C.F. Xing, H.H. Hu, J.B. Huang, H.C. Fang, Y.H. Kai, Y.C. Wu, S.C. Chi, Diet supplementation of *Pediococcus pentosaceus* in cobia (*Rachycentron canadum*) enhances growth rate, respiratory burst and resistance against photobacteriosis, Fish. Shellfish Immunol. 35 (2013) 1122–1128.
- [125] M. Yadav, J.S. Schorey, The b-glucan receptor dectin-1 functions together with TLR2 to mediate macrophage activation by mycobacteria, Blood 108 (2002) 3168–3175.
- [126] L. Yan, K.G. Boyd, J.G. Burgess, Surface attachment induced production of antimicrobial compounds by marine epiphytic bacteria using modified roller bottle cultivation, Mar. Biotechnol. 4 (2002) 356–366.
- [127] J.D. Ye, K. Wang, F.D. Li, Y.Z. Sun, Single or combined effects of fructo- and mannan oligosaccharide supplements and *Bacillus clausii* on the growth, feed utilization, body composition, digestive enzyme activity, innate immune response and lipid metabolism of Japanese flounder *Paralichthys olivaceus*, Aquacult Nutr. 17 (2011) 902–911.
- [128] C.N. Zhang, X.F. Li, W.N. Xu, G.Z. Jiang, K.L. Lu, L.N. Wang, Combined effects of dietary fructooligosaccharide and *Bacillus licheniformis* on innate immunity, antioxidant capability and disease resistance of triangular bream (*Megalobrama terminalis*), Fish. Shellfish Immunol. 35 (2013) 1380–1386.
- [129] Q. Zhang, H. Ma, K. Mai, W. Zhang, Z. Liufu, W. Xu, Interaction of dietary *Bacillus subtilis* and fructooligosaccharide on the growth performance, non-specific immunity of sea cucumber, *Apostichopus japonicus*, Fish. Shellfish Immunol. 29 (2010) 204–211.
- [130] Q.C. Zhou, J.A. Buentello, D.M. Gatlin III, Effects of dietary prebiotics on growth performance, immune response and intestinal morphology of red drum (*Sciaenops ocellatus*), Aquaculture 309 (2010) 253–257.
- [131] H. Zokaeifar, J.L. Balcazar, C.R. Saad, M.S. Kamarudin, K. Sijam, A. Arshad, N. Nejat, Effects of *Bacillus subtilis* on the growth performance, digestive enzymes, immune gene expression and disease resistance of white shrimp, *Litopenaeus vannamei*, Fish. Shellfish Immunol. 33 (2012) 683–689.