

# ISyE6669 Homework 5 Solution

Fall 2021

1. Compute the gradient  $\nabla f(x)$  and Hessian  $\nabla^2 f(x)$  of the Rosenbrock function

$$f(x_1, x_2) = 100(x_2 - x_1^2)^2 + (1 - x_1)^2.$$

**Solution:**

$$\begin{aligned}\nabla f(x) &= \begin{bmatrix} 200(x_2 - x_1^2)(-2x_1) + 2(1 - x_1)(-1) \\ 200(x_2 - x_1^2) \end{bmatrix} \\ &= \begin{bmatrix} 400(x_1^3 - x_1x_2) + 2(x_1 - 1) \\ 200(x_2 - x_1^2) \end{bmatrix} \end{aligned} \tag{1}$$

$$\nabla^2 f(x) = \begin{bmatrix} 1200x_1^2 - 400x_2 + 2, & -400x_1 \\ -400x_1, & 200 \end{bmatrix} \tag{2}$$

2. Implement the Newton's Method with line search given in Algorithm 1. Use the Newton's

---

**Algorithm 1** Newton's Method with Line Search

---

Start with  $x^0$ . Set  $k = 0$ ,  $\epsilon = 10^{-4}$ .  
Set  $d^0 \leftarrow -(\nabla^2 f(x^0))^{-1} \nabla f(x^0)$   
**while**  $\|\nabla f(x^k)\| > \epsilon$  **do**  
    Choose  $\bar{\alpha} > 0, \rho \in (0, 1), c \in (0, 1)$ ; Set  $\alpha^k \leftarrow \bar{\alpha}$   
    **while**  $f(x^k + \alpha^k d^k) > f(x^k) + c\alpha^k \nabla f(x^k)^\top d^k$  **do**  
         $\alpha^k \leftarrow \rho \alpha^k$   
    **end while**  
     $x^{k+1} \leftarrow x^k + \alpha^k d^k$   
     $d^{k+1} \leftarrow -(\nabla^2 f(x^{k+1}))^{-1} \nabla f(x^{k+1})$   
     $k \leftarrow k + 1$   
**end while**

---

Method to minimize the Rosenbrock function in Problem 1. Set the initial stepsize  $\bar{\alpha} = 1$ . Select your own choice of  $\rho \in (0, 1), c \in (0, 1)$ . First run the algorithm from the initial point  $x^0 = (1.2, 1.2)^\top$ , and then try the more difficult starting point  $x^0 = (-1.2, 1)^\top$ . For each starting point, print out the step length  $\alpha^k$  used by the algorithm as well as the point  $x^k$  for every step  $k$ . You should observe that Newton's Method converges very fast.

Note that answers may vary depending on what  $\rho$  and  $c$  values were used. We use  $\rho = 0.9$  and  $c = 10^{-5}$ . We do not break the inner backtracking while loop in our code, but you can if  $\alpha$  falls below 0.2 or 0.3. Such answers will be accepted. Using initial point  $x^0 = (1.2, 1.2)^\top$ , we get the following output from our code:

```
iter =  
1  
  
alpha =  
1  
  
x =  
1.195918367346939  
1.430204081632654  
  
iter =  
2  
  
alpha =  
0.6561000000000000  
  
x =  
1.067803199227371  
1.123784446698436  
  
iter =  
3  
  
alpha =
```

0.9000000000000000

x =

1.053558314632527  
1.108140283036032

iter =

4

alpha =

0.9000000000000000

x =

1.018347492953793  
1.035607330512453

iter =

5

alpha =

0.9000000000000000

x =

1.005495680368007  
1.010713965562699

iter =

6

```
alpha =  
  
0.9000000000000000
```

```
x =  
  
1.000836216112118  
1.001620661106868
```

```
iter =  
  
7
```

```
alpha =  
  
0.9000000000000000
```

```
x =  
  
1.000091437375132  
1.000177081378216
```

```
iter =  
  
8
```

```
alpha =  
  
0.9000000000000000
```

```
x =  
  
1.000009239115984  
1.000017891387492
```

```
iter =
```

9

alpha =

0.9000000000000000

x =

1.000000924887601

1.000001791013948

Note that Newton's method converges very fast. Our code converges in 9 iterations to optimal solution  $x_* = (1, 1)^\top$ .

Using initial point  $x^0 = (-1.2, 1)^\top$ , we get the following output from our code:

iter =

1

alpha =

1

x =

-1.175280898876405

1.380674157303371

iter =

2

alpha =

0.166771816996666

```
x =  
  
-0.8520111114246042  
0.620910454427278
```

```
iter =  
  
3
```

```
alpha =  
  
0.9000000000000000
```

```
x =  
  
-0.776255621024348  
0.586332645981484
```

```
iter =  
  
4
```

```
alpha =  
  
0.7290000000000000
```

```
x =  
  
-0.471434147807713  
0.124932946380622
```

```
iter =  
  
5
```

```
alpha =
```

0.9000000000000000

x =

-0.406719192152260  
0.151500754845643

iter =

6

alpha =

0.7290000000000000

x =

-0.135706497703422  
-0.058803878313132

iter =

7

alpha =

0.9000000000000000

x =

-0.073548007735245  
-0.006176381616507

iter =

8

```
alpha =  
  
    0.8100000000000000
```

```
x =  
  
    0.188597803134905  
   -0.035352576110539
```

```
iter =  
  
    9
```

```
alpha =  
  
    0.9000000000000000
```

```
x =  
  
    0.236690898089599  
    0.046617464710355
```

```
iter =  
  
    10
```

```
alpha =  
  
    0.8100000000000000
```

```
x =  
  
    0.451295330945479  
    0.155825440991456
```

```
iter =
```



11

alpha =

0.9000000000000000

x =

0.498022733151194

0.241058989144317

iter =

12

alpha =

0.9000000000000000

x =

0.686772990971616

0.435333715941717

iter =

13

alpha =

0.9000000000000000

x =

0.720882495066941

0.514875770905670

```

iter =

    14

alpha =

    0.9000000000000000

x =

    0.849103637916556
    0.704056746370642

iter =

    15

alpha =

    0.9000000000000000

x =

    0.880081108112290
    0.771891129041179

iter =

    16

alpha =

    0.9000000000000000

x =

```

```
0.950606625890149
0.898413945746921
```

```
iter =
```

```
17
```

```
alpha =
```

```
0.9000000000000000
```

```
x =
```

```
0.972314794184649
0.944400913275701
```

```
iter =
```

```
18
```

```
alpha =
```

```
0.9000000000000000
```

```
x =
```

```
0.993095511134515
0.985707341467110
```

```
iter =
```

```
19
```

```
alpha =
```

```
0.9000000000000000
```

```
x =  
  
    0.998712618051308  
    0.997342206287941
```

```
iter =  
  
    20
```

```
alpha =  
  
    0.9000000000000000
```

```
x =  
  
    0.999851964205533  
    0.999694183499308
```

```
iter =  
  
    21
```

```
alpha =  
  
    0.9000000000000000
```

```
x =  
  
    0.999984936676745  
    0.999968879216080
```

```
iter =  
  
    22
```

```
alpha =

    0.9000000000000000
```

```
x =

    0.999998490972099
    0.999996882326324
```

Using this more difficult initial point, we converge in 22 iterations to optimal solution  $x_* = (1, 1)^\top$ .

Here is the Matlab code (you can use Python alternatively) that generates the above output.

```
1 clc
2 clear all
3 close all
4 x0=[-1.2;1];
5 epsilon=10^-4;
6 alphabar=1;
7 c=1*10^-5;
8 rho=0.9;
9
10 f=@(x1,x2) 100*(x2-x1^2)^2+(1-x1)^2;
11 gradf=@(x1,x2) [400*(x1^3-x1*x2)+2*(x1-1);200*(x2-x1^2)];
12 hessf=@(x1,x2) [1200*x1^2-400*x2+2 -400*x1; -400*x1 200];
13
14 d=-inv(hessf(x0(1),x0(2)))*gradf(x0(1),x0(2));
15
16 x=x0;
17 z=x+alphabar*d;
18 iter=0;
19 while norm (gradf(x(1),x(2)))>epsilon
20     alpha=alphabar;
21     while f(z(1),z(2))>f(x(1),x(2))+(c*alpha)*transpose(gradf(x(1),
22         x(2)))*d
23         alpha=rho*alpha;
24         z=x+alpha*d;
25     end
26     iter=iter+1
27     alpha
28     x=x+alpha*d
29
30     d=-inv(hessf(x(1),x(2)))*gradf(x(1),x(2));
```

31 **end**

Observe that Newton's method converges in less iterations for initial point  $x^0 = (1.2, 1.2)^\top$  than initial point  $x^0 = (-1.2, 1)^\top$  because  $x^0 = (1.2, 1.2)^\top$  is closer in distance to the optimal solution  $x_* = (1, 1)^\top$ . Note we also take  $c$  to be very small (i.e.  $10^{-5}$ ) and  $\rho$  to be large (i.e. 0.9) so the condition in the backtracking line search is satisfied faster and so we do not reduce the stepsize as much in this loop. This way, since the stepsize is not too small, we make significant progress towards the optimal solution at each iteration and thus converge in a very small number of iterations.

- Figure 1 below illustrates the water network of Newvillage. The lines are water pipelines numbered from 1 through 13. The arrows on the lines are possible direction(s) of flow of water in these pipelines. The circles are water sources numbered A, B, C. The rectangles are houses D, E, F, G, H. The maximum possible capacity of the water sources are (the sources can operate at less than the maximum capacity): A: 100 Units, B: 100 Units, C: 80 Units Demands of water in the houses are: D: 50 Units, E: 60 Units, F: 40 Units, G: 30 Units, H: 70 Units Since the houses are at different elevation and the pipes are of different diameter, the cost of transporting water is different in the different pipes. These costs per unit of water are: Pipe 1: \$2, Pipe 2: \$3, Pipe 3: \$4, Pipe 4: \$2, Pipe 5: \$3, Pipe 6: \$2, Pipe 7: \$4, Pipe 8: \$1, Pipe 9: \$2, Pipe 10: \$4, Pipe 11: \$5, Pipe 12: \$1, Pipe 13: \$2. Formulate an LP to minimize the total cost of transporting water so as to meet the water demands of each house.




Figure 1: Water Network for Question 2

### Solution:

We introduce the following decision variables.  $x_A, x_B, x_C$ , respectively, is the amount of water pumped through water source A, B, C, and  $f_{ij}$  is the flow of water from location  $i$  to location  $j$ , for arcs  $(i, j)$  in the set of arc  $\mathcal{A} := \{(A, D), (A, E), (B, D), (B, F), (B, G), (C, F), (C, G), (C, H), (D, F), (E, F), (E, H), (F, D), (F, G), (H, E), (H, F)\}$ .

$$\begin{aligned}
\min \quad & 2f_{AD} + 4f_{AE} + 3f_{BD} + 3f_{BF} + 2f_{BG} + 5f_{CF} + f_{CG} + 2f_{CH} + 2f_{DF} + 4f_{EF} + 2f_{EH} + 2f_{FD} \\
& + f_{FG} + 2f_{HE} + 4f_{HF} \tag{3} \\
\text{s.t.} \quad & x_A - f_{AD} - f_{AE} = 0 \tag{4} \\
& x_B - f_{BD} - f_{BF} - f_{BG} = 0 \tag{5} \\
& x_C - f_{CF} - f_{CG} - f_{CH} = 0 \tag{6} \\
& f_{AD} + f_{BD} - f_{DF} + f_{FD} = 50 \tag{7} \\
& f_{AE} + f_{HE} - f_{EH} - f_{EF} = 60 \tag{8} \\
& f_{BF} + f_{CF} + f_{DF} + f_{EF} + f_{HF} - f_{FD} - f_{FG} = 40 \tag{9} \\
& f_{BG} + f_{FG} + f_{CG} = 30 \tag{10} \\
& f_{CH} + f_{EH} - f_{HE} - f_{HF} = 70 \tag{11} \\
& x_A \leq 100, \tag{12} \\
& x_B \leq 100, \tag{13} \\
& x_C \leq 80, \tag{14} \\
& x_i \geq 0, \quad \forall i \in \{A, B, C\}, \tag{15} \\
& f_{ij} \geq 0, \quad \forall (i, j) \in \mathcal{A}. \tag{16}
\end{aligned}$$

**Note:** Edges  $DF$  and  $EH$  are bidirectional, meaning that there is flow in both directions. Thus we have two variables for each edge:  $f_{DF}$  and  $f_{FD}$  for  $DF$ , and  $f_{EH}$  and  $f_{HE}$  for  $EH$ . Some of the students assumed that arrows pointing to both directions, that may be flow in one direction or the other, but not both at the same time. We will accept that formulation too. Following is the solution in that case.

We introduce the following decision variables.  $x_A, x_B, x_C$ , respectively, is the amount of water pumped through water source A, B, C, and  $f_{ij}$  is the flow of water from location  $i$  to location  $j$ , for arcs  $(i, j)$  in the set of arc  $\mathcal{A} := \{(A, D), (A, E), (B, D), (B, F), (B, G), (C, F), (C, G), (C, H), (D, F), (E, F), (E, H), (F, G), (H, F)\}$ . And let  $\mathcal{D} = \{(D, F), (E, H)\} \subset \mathcal{A}$  denotes set of edges for which flow can go in either direction.

$$\begin{aligned}
\min \quad & 2f_{AD} + 4f_{AE} + 3f_{BD} + 3f_{BF} + 2f_{BG} + 5f_{CF} + f_{CG} + 2f_{CH} + 2|f_{DF}| + 4f_{EF} + 2|f_{EH}| \\
& + f_{FG} + 4f_{HF} \\
\text{s.t.} \quad & x_A - f_{AD} - f_{AE} = 0 \\
& x_B - f_{BD} - f_{BF} - f_{BG} = 0 \\
& x_C - f_{CF} - f_{CG} - f_{CH} = 0 \\
& f_{AD} + f_{BD} - f_{DF} = 50 \\
& f_{AE} + f_{HE} - f_{EH} - f_{EF} = 60 \\
& f_{BF} + f_{CF} + f_{DF} + f_{EF} + f_{HF} - f_{FG} = 40 \\
& f_{BG} + f_{FG} + f_{CG} = 30 \\
& f_{CH} + f_{EH} - f_{HF} = 70 \\
& x_A \leq 100, \\
& x_B \leq 100, \\
& x_C \leq 80, \\
& x_i \geq 0, \quad \forall i \in \{A, B, C\}, \\
& f_{ij} \geq 0, \quad \forall (i, j) \in \mathcal{A} \setminus \mathcal{D}.
\end{aligned}$$

This is not a linear problem due to two absolute values in the objective function. Hence it will need to be converted into:

$$\begin{aligned}
\min \quad & 2f_{AD} + 4f_{AE} + 3f_{BD} + 3f_{BF} + 2f_{BG} + 5f_{CF} + f_{CG} + 2f_{CH} + 2z + 4f_{EF} + 2w \\
& + f_{FG} + 4f_{HF} \\
\text{s.t.} \quad & x_A - f_{AD} - f_{AE} = 0 \\
& x_B - f_{BD} - f_{BF} - f_{BG} = 0 \\
& x_C - f_{CF} - f_{CG} - f_{CH} = 0 \\
& f_{AD} + f_{BD} - f_{DF} = 50 \\
& f_{AE} + f_{HE} - f_{EH} - f_{EF} = 60 \\
& f_{BF} + f_{CF} + f_{DF} + f_{EF} + f_{HF} - f_{FG} = 40 \\
& f_{BG} + f_{FG} + f_{CG} = 30 \\
& f_{CH} + f_{EH} - f_{HF} = 70 \\
& x_A \leq 100, \\
& x_B \leq 100, \\
& x_C \leq 80, \\
& x_i \geq 0, \quad \forall i \in \{A, B, C\}, \\
& f_{ij} \geq 0, \quad \forall (i, j) \in \mathcal{A} \setminus \mathcal{D} \\
& -z \leq f_{DF} \leq z \\
& -w \leq f_{EH} \leq w.
\end{aligned}$$



4. Consider the following electric power network shown in Figure 2. This network is taken from a real-world electric power system. Electricity generators are located at nodes 1, 3, and 5 and producing  $p_1, p_2, p_3$  amounts of electricity, respectively. Electricity loads are located at nodes 2, 4, and 6 and are consuming  $d_1, d_2, d_3$  amounts of electricity, respectively.

The demand is fixed and given as  $d_1 = 110$ ,  $d_2 = 65$ ,  $d_3 = 95$ .

Each generator  $i$ 's production must be within an upper and a lower bound as  $p_i^{\min} \leq p_i \leq p_i^{\max}$ . The bounds are given as  $p_1^{\min} = 20$ ,  $p_1^{\max} = 200$ ,  $p_2^{\min} = 20$ ,  $p_2^{\max} = 150$ ,  $p_3^{\min} = 10$ ,  $p_3^{\max} = 150$ .

The flow limits over lines are given as  $f_{12}^{\max} = 100$ ,  $f_{23}^{\max} = 110$ ,  $f_{34}^{\max} = 50$ ,  $f_{45}^{\max} = 80$ ,  $f_{56}^{\max} = 60$ ,  $f_{61}^{\max} = 40$ .

The line parameters are given as  $B_{12} = 11.6$ ,  $B_{23} = 5.9$ ,  $B_{34} = 13.7$ ,  $B_{45} = 9.8$ ,  $B_{56} = 5.6$ ,  $B_{61} = 10.5$ . The unit generation costs are given as  $c_1 = 16$ ,  $c_2 = 20$ ,  $c_3 = 8$ .

- Formulate the power system scheduling problem using the model discussed in Lecture 2.
- Implement and solve the model using CVXPY. Write down the optimal solution.
- Find the electricity prices for demand at nodes 2, 4, 6. To do this, use the command `constraints[0].dual_value` to find the dual variable of `constraints[0]`. Hint: Recall the electricity price at node  $i$  is the dual variable for the flow conservation constraint at node  $i$ .




Figure 2: Electric network for Question 3.

(a) The LP can be formulated as follows:

$$\min \sum_{i=1..3} c_i p_i$$

s.t.

*Flow Conservation:*

$$f_{12} - f_{61} = p_1,$$

$$f_{34} - f_{23} = p_2,$$

$$f_{56} - f_{45} = p_3,$$

$$f_{23} - f_{12} = -d_1,$$

$$f_{45} - f_{34} = -d_2,$$

$$f_{61} - f_{56} = -d_3,$$

*Constraints linking branch flow and nodal potential:*

$$f_{12} = B_{12}(\theta_1 - \theta_2),$$

$$f_{23} = B_{23}(\theta_2 - \theta_3),$$

$$f_{34} = B_{34}(\theta_3 - \theta_4),$$

$$f_{45} = B_{45}(\theta_4 - \theta_5),$$

$$f_{56} = B_{56}(\theta_5 - \theta_6),$$

$$f_{61} = B_{61}(\theta_6 - \theta_1),$$

*Flow limit constraints:*

$$-f_{12}^{\max} \leq f_{12} \leq f_{12}^{\max},$$

$$-f_{23}^{\max} \leq f_{23} \leq f_{23}^{\max},$$

$$-f_{34}^{\max} \leq f_{34} \leq f_{34}^{\max},$$

$$-f_{45}^{\max} \leq f_{45} \leq f_{45}^{\max},$$

$$-f_{56}^{\max} \leq f_{56} \leq f_{56}^{\max},$$

$$-f_{61}^{\max} \leq f_{61} \leq f_{61}^{\max},$$

*Generation constraints:*

$$p_1^{\min} \leq p_1 \leq p_1^{\max},$$

$$p_2^{\min} \leq p_2 \leq p_2^{\max},$$

$$p_3^{\min} \leq p_3 \leq p_3^{\max}.$$

(b) The optimal objective value is . The quantum of electricity produced are  $p_1 = 113.12, p_2 = 20.0, p_3 = 136.88$ .

Following is the output from python model

-The optimal value is 3304.96

-The quantum of electricity produced at node 1:  $p_1 = 113.12$  , at node 3:  $p_2 = 20.0$  , at node 5:  $p_3 = 136.88$

-The flows in different lines are  $f_{12} = 78.12$  ,  $f_{23} = -31.88$  ,  $f_{34} = -11.88$  ,  $f_{45} = -76.88$  ,  $f_{56} = 60.0$  and  $f_{61} = -35.0$

(c) The electricity price at node 2 is 14.4 , at node 4 is 9.9 and at node 6 is 17.77 .