

## Introduction of a new variable

- Consider the problem (**P**),  
Min  $\mathbf{c}^T \mathbf{x}$   
subject to  
 $A\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .  
Let  $\mathbf{x}_0$  be an **optimal BFS** of this problem.
- A new variable  $x_{n+1}$  is added to the LPP given above.

## Introduction of a new variable

- Consider the problem (P),

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}.$$

Let  $\mathbf{x}_0$  be an optimal BFS of this problem.

- A new variable  $x_{n+1}$  is added to the LPP given above.
- A column is added to the matrix  $A$  call it  $\tilde{\mathbf{a}}_{n+1}$  and a component  $c_{n+1}$  is added to the cost vector  $\mathbf{c}$ , which is the cost associated with the new variable  $x_{n+1}$ .

## Introduction of a new variable

- Consider the problem (**P**),

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}.$$

Let  $\mathbf{x}_0$  be an **optimal BFS** of this problem.

- A new variable  $x_{n+1}$  is added to the LPP given above.
- A column is added to the matrix  $A$  call it  $\tilde{\mathbf{a}}_{n+1}$  and a component  $c_{n+1}$  is added to the cost vector  $\mathbf{c}$ , which is the cost associated with the new variable  $x_{n+1}$ .
- Since  $\mathbf{x}_0$  is optimal for **P**  
 $\mathbf{x}'_0 = [\mathbf{x}_0^T, 0]^T$  is feasible for the new problem (P'), given by:

## Introduction of a new variable

- Consider the problem (**P**),

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}.$$

Let  $\mathbf{x}_0$  be an **optimal BFS** of this problem.

- A new variable  $x_{n+1}$  is added to the LPP given above.
- A column is added to the matrix  $A$  call it  $\tilde{\mathbf{a}}_{n+1}$  and a component  $c_{n+1}$  is added to the cost vector  $\mathbf{c}$ , which is the cost associated with the new variable  $x_{n+1}$ .
- Since  $\mathbf{x}_0$  is optimal for **P**  
 $\mathbf{x}'_0 = [\mathbf{x}_0^T, 0]^T$  is feasible for the new problem (P'), given by:
- Min  $[\mathbf{c}, c_{n+1}]^T \mathbf{x}_{(n+1) \times 1}$   
subject to  
 $[\mathbf{A} : \tilde{\mathbf{a}}_{n+1}] \mathbf{x}_{(n+1) \times 1} = \mathbf{b}, \quad \mathbf{x}_{(n+1) \times 1} \geq \mathbf{0}.$

- To check whether  $\mathbf{x}'_0$  is optimal for  $(P')$ , add a new column to the optimal table and calculate  $C_{n+1} - Z_{n+1}$ .

- To check whether  $\mathbf{x}'_0$  is optimal for (P'), add a new column to the optimal table and calculate  $c_{n+1} - z_{n+1}$ .
- If  $c_{n+1} - z_{n+1} \geq 0$ , then  $\mathbf{x}'_0$  is optimal for the new problem.

- To check whether  $\mathbf{x}'_0$  is optimal for (P'), add a new column to the optimal table and calculate  $c_{n+1} - z_{n+1}$ .
- If  $c_{n+1} - z_{n+1} \geq 0$ , then  $\mathbf{x}'_0$  is optimal for the new problem.
- If **not**, then enter  $\tilde{\mathbf{a}}_{n+1}$  in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.

- To check whether  $\mathbf{x}'_0$  is optimal for (P'), add a new column to the optimal table and calculate  $c_{n+1} - z_{n+1}$ .
- If  $c_{n+1} - z_{n+1} \geq 0$ , then  $\mathbf{x}'_0$  is optimal for the new problem.
- If **not**, then enter  $\tilde{\mathbf{a}}_{n+1}$  in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.
- Adding a **variable** to **P** results in adding a **constraint** to the dual,



- To check whether  $\mathbf{x}'_0$  is optimal for (P'), add a new column to the optimal table and calculate  $c_{n+1} - z_{n+1}$ .
- If  $c_{n+1} - z_{n+1} \geq 0$ , then  $\mathbf{x}'_0$  is optimal for the new problem.
- If **not**, then enter  $\tilde{\mathbf{a}}_{n+1}$  in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.
- Adding a **variable** to **P** results in adding a **constraint** to the dual, If the feasible region of the new dual becomes empty,

- To check whether  $\mathbf{x}'_0$  is optimal for (P'), add a new column to the optimal table and calculate  $c_{n+1} - z_{n+1}$ .
- If  $c_{n+1} - z_{n+1} \geq 0$ , then  $\mathbf{x}'_0$  is optimal for the new problem.
- If **not**, then enter  $\tilde{a}_{n+1}$  in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.
- Adding a **variable** to **P** results in adding a **constraint** to the dual, If the feasible region of the new dual becomes empty, then **P'** will **not** have an optimal solution.

## Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.

## Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ , where  $\mathbf{a}_{m+1}^T$  is a row vector.

## Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ , where  $\mathbf{a}_{m+1}^T$  is a row vector.
- **Case 1:**  $\mathbf{x}_0$  satisfies  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ .

## Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ , where  $\mathbf{a}_{m+1}^T$  is a row vector.
- **Case 1:**  $\mathbf{x}_0$  satisfies  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ .  
Then  $\mathbf{x}_0$  is optimal for the changed problem also.

## Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ , where  $\mathbf{a}_{m+1}^T$  is a row vector.
- **Case 1:**  $\mathbf{x}_0$  satisfies  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ .  
Then  $\mathbf{x}_0$  is optimal for the changed problem also.
- **Case 2:**  $\mathbf{x}_0$  does not satisfy the new constraint, that is  $\mathbf{a}_{m+1}^T \mathbf{x}_0 > b_{m+1}$ ,

## Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ , where  $\mathbf{a}_{m+1}^T$  is a row vector.
- **Case 1:**  $\mathbf{x}_0$  satisfies  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ .  
Then  $\mathbf{x}_0$  is optimal for the changed problem also.
- **Case 2:**  $\mathbf{x}_0$  does not satisfy the new constraint, that is  $\mathbf{a}_{m+1}^T \mathbf{x}_0 > b_{m+1}$ ,  
or  $\mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} > b_{m+1}$ ,



## Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ , where  $\mathbf{a}_{m+1}^T$  is a row vector.
- **Case 1:**  $\mathbf{x}_0$  satisfies  $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$ .  
Then  $\mathbf{x}_0$  is optimal for the changed problem also.
- **Case 2:**  $\mathbf{x}_0$  does not satisfy the new constraint, that is  $\mathbf{a}_{m+1}^T \mathbf{x}_0 > b_{m+1}$ ,  
or  $\mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} > b_{m+1}$ ,  
where  $\mathbf{a}_{B,m+1}^T$  and  $\mathbf{a}_{N,m+1}^T$  are the components of  $\mathbf{a}_{m+1}^T$  corresponding to the basic variables and nonbasic variables of  $\mathbf{x}_0$ , respectively.

- The new constraint can be written as  $\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}$ , where  $s_{m+1} \geq 0$ .

- The new constraint can be written as  $\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}$ , where  $s_{m+1} \geq 0$ .
- If  $B$  denotes the basis matrix corresponding to  $\mathbf{x}_0$ , then

- The new constraint can be written as  $\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}$ , where  $s_{m+1} \geq 0$ .
- If  $B$  denotes the basis matrix corresponding to  $\mathbf{x}_0$ , then 
$$\begin{bmatrix} B & \mathbf{0} \\ \mathbf{a}_{B,m+1}^T & 1 \end{bmatrix}$$
 is the new basis matrix after the new constraint is added, the newly added **basic variable** being  $s_{m+1}$ .

- The new constraint can be written as  $\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}$ , where  $s_{m+1} \geq 0$ .
- If  $B$  denotes the basis matrix corresponding to  $\mathbf{x}_0$ , then 
$$\begin{bmatrix} B & \mathbf{0} \\ \mathbf{a}_{B,m+1}^T & 1 \end{bmatrix}$$
 is the new basis matrix after the new constraint is added, the newly added **basic variable** being  $s_{m+1}$ .
- The **inverse** of this new basis matrix is given by 
$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix}.$$

- The new row added to the (main) simplex table will be of the form

$$\begin{aligned} & [-\mathbf{a}_{B,m+1}^T B^{-1}, 1] \begin{bmatrix} \mathbf{a}_{B,m+1}^T B & \mathbf{a}_{N,m+1}^T N & \mathbf{0} \\ \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & 1 \end{bmatrix} = \\ & [\mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1] \end{aligned}$$

- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ \mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ \mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- If  $\mathbf{x}_0$  does not satisfy the newly added constraint then the  $(m+1)$  th entry of the RHS will be **strictly** less than zero.



- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ \mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- If  $\mathbf{x}_0$  does not satisfy the newly added constraint then the  $(m+1)$  th entry of the RHS will be **strictly** less than zero.
- Since the cost associated with  $s_{m+1}$  is equal to zero, all the  $c_j - z_j$  entries remain unchanged and are non negative.

- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ \mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- If  $\mathbf{x}_0$  does not satisfy the newly added constraint then the  $(m+1)$  th entry of the RHS will be **strictly** less than zero.
- Since the cost associated with  $s_{m+1}$  is equal to zero, all the  $c_j - z_j$  entries remain unchanged and are non negative.
- Then **Dual Simplex** can be used either to obtain the new optimal solution or to conclude that new (P) is **infeasible**.

## Changing entries in the coefficient matrix $A$ :

- **Case 1:** The column corresponding to a **nonbasic variable** say  $\tilde{a}_j$  of the optimal solution is changed to  $\tilde{a}'_j$ .

## Changing entries in the coefficient matrix $A$ :

- **Case 1:** The column corresponding to a **nonbasic variable** say  $\tilde{\mathbf{a}}_j$  of the optimal solution is changed to  $\tilde{\mathbf{a}}'_j$ .
- The column corresponding to that variable in the optimal simplex table  $B^{-1}\tilde{\mathbf{a}}_j$  is changed to  $B^{-1}\tilde{\mathbf{a}}'_j$ .

## Changing entries in the coefficient matrix $A$ :

- **Case 1:** The column corresponding to a **nonbasic variable** say  $\tilde{\mathbf{a}}_j$  of the optimal solution is changed to  $\tilde{\mathbf{a}}'_j$ .
- The column corresponding to that variable in the optimal simplex table  $B^{-1}\tilde{\mathbf{a}}_j$  is changed to  $B^{-1}\tilde{\mathbf{a}}'_j$ .
- If the new  $c_j - z'_j$  value satisfies the **optimality condition**, then the previous optimal solution will be **optimal** for the new problem.

## Changing entries in the coefficient matrix $A$ :

- **Case 1:** The column corresponding to a **nonbasic variable** say  $\tilde{\mathbf{a}}_j$  of the optimal solution is changed to  $\tilde{\mathbf{a}}'_j$ .
- The column corresponding to that variable in the optimal simplex table  $B^{-1}\tilde{\mathbf{a}}_j$  is changed to  $B^{-1}\tilde{\mathbf{a}}'_j$ .
- If the new  $c_j - z'_j$  value satisfies the **optimality condition**, then the previous optimal solution will be **optimal** for the new problem.
- If **not**, then use **Simplex** either to obtain the new **optimal solution** or to conclude that new (P) does **not** have an optimal solution.

## Changing entries in the coefficient matrix $A$ :

- **Case 2:** If the column corresponding to the  $j$  th **basic variable** say  $\tilde{\mathbf{a}}_j$  is changed to  $\tilde{\mathbf{a}}'_j$ .

## Changing entries in the coefficient matrix $A$ :

- **Case 2:** If the column corresponding to the  $j$  th **basic variable** say  $\tilde{\mathbf{a}}_j$  is changed to  $\tilde{\mathbf{a}}'_j$ .
- Then treat this case as the case when a variable (say a variable  $x_{n+1}$ ) is added to the problem with column  $\tilde{\mathbf{a}}'_j$  and cost  $c_j$ .



## Changing entries in the coefficient matrix $A$ :

- **Case 2:** If the column corresponding to the  $j$  th **basic variable** say  $\tilde{\mathbf{a}}_j$  is changed to  $\tilde{\mathbf{a}}'_j$ .
- Then treat this case as the case when a variable (say a variable  $x_{n+1}$ ) is added to the problem with column  $\tilde{\mathbf{a}}'_j$  and cost  $c_j$ .
- Add a column  $B^{-1}\tilde{\mathbf{a}}'_j$  and the corresponding  $c_{n+1} - z_{n+1}$  value to the previous simplex table.

## Changing entries in the coefficient matrix $A$ :

- **Case 2:** If the column corresponding to the  $j$  th **basic variable** say  $\tilde{\mathbf{a}}_j$  is changed to  $\tilde{\mathbf{a}}'_j$ .
- Then treat this case as the case when a variable (say a variable  $x_{n+1}$ ) is added to the problem with column  $\tilde{\mathbf{a}}'_j$  and cost  $c_j$ .
- Add a column  $B^{-1}\tilde{\mathbf{a}}'_j$  and the corresponding  $c_{n+1} - z_{n+1}$  value to the previous simplex table.
- If  $u_{j,(n+1)} = 0$ , then it implies that  $\{\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_{j-1}, \tilde{\mathbf{a}}'_j, \tilde{\mathbf{a}}_{j+1}, \dots, \tilde{\mathbf{a}}_m\}$  is **LD** and we have to find an **initial BFS** for the changed problem.

## Changing entries in the coefficient matrix $A$ :

- **Case 2:** If the column corresponding to the  $j$  th **basic variable** say  $\tilde{\mathbf{a}}_j$  is changed to  $\tilde{\mathbf{a}}'_j$ .
- Then treat this case as the case when a variable (say a variable  $x_{n+1}$ ) is added to the problem with column  $\tilde{\mathbf{a}}'_j$  and cost  $c_j$ .
- Add a column  $B^{-1}\tilde{\mathbf{a}}'_j$  and the corresponding  $c_{n+1} - z_{n+1}$  value to the previous simplex table.
- If  $u_{j,(n+1)} = 0$ , then it implies that  $\{\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_{j-1}, \tilde{\mathbf{a}}'_j, \tilde{\mathbf{a}}_{j+1}, \dots, \tilde{\mathbf{a}}_m\}$  is **LD** and we have to find an **initial BFS** for the changed problem.
- If  $u_{j,(n+1)} \neq 0$ , then pivot on this element and make  $x_{n+1}$  enter the basis and  $x_j$  leave the basis.

- Perform the necessary elementary row operations to make the  $(n+1)$  th column as the  $j$  **th column** of  $I$  and  $C_{n+1} - Z_{n+1}$  equal to 0.

- Perform the necessary elementary row operations to make the  $(n+1)$  th column as the  $j$  **th column** of  $I$  and  $c_{n+1} - z_{n+1}$  equal to 0.
- Delete the column corresponding to  $\tilde{a}_j$  from all subsequent calculations, and the variable  $x_{n+1}$  is now indexed as  $x_j$ .

- Perform the necessary elementary row operations to make the  $(n+1)$  th column as the  $j$  **th column** of  $I$  and  $C_{n+1} - Z_{n+1}$  equal to 0.
- Delete the column corresponding to  $\tilde{a}_j$  from all subsequent calculations, and the variable  $x_{n+1}$  is now indexed as  $x_j$ .
- The necessary calculations mentioned above might disturb the optimality as well as the feasibility of the current **BFS**.

- Perform the necessary elementary row operations to make the  $(n+1)$  th column as the  $j$  **th column** of  $I$  and  $c_{n+1} - z_{n+1}$  equal to 0.
- Delete the column corresponding to  $\tilde{a}_j$  from all subsequent calculations, and the variable  $x_{n+1}$  is now indexed as  $x_j$ .
- The necessary calculations mentioned above might disturb the optimality as well as the feasibility of the current **BFS**.
- If however the new  $c_j - z_j$  values are all **nonnegative** and all the RHS entries remain **non negative**, then the table is **optimal** and the corresponding BFS is **optimal** for the new (P).

- If the new table has all RHS entries **nonnegative**, but atleast one of the  $c_j - z_j$  values is **negative** then use **Simplex** either to obtain the new optimal solution, or to conclude that the new **(P)** **does not** have an optimal solution.



- If the new table has all RHS entries **nonnegative**, but atleast one of the  $c_j - z_j$  values is **negative** then use **Simplex** either to obtain the new optimal solution, or to conclude that the new **(P)** **does not** have an optimal solution.
- If all the  $c_j - z_j$  values are **nonnegative** but atleast one of the RHS entries in the simplex table is **negative** then use **Dual Simplex** to obtain the new optimal solution or to conclude that the new **(P)** has **no** feasible solution.

## Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:  
Min  $\mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .

## Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:  
Min  $\mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .
- Assume  $\mathbf{b} \geq \mathbf{0}$ .

## Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:  
Min  $\mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .
- Assume  $\mathbf{b} \geq \mathbf{0}$ .
- Add artificial variables  $w_1, \dots, w_m$  with cost  $M$  ( $M$  large).

## Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:  
Min  $\mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .
- Assume  $\mathbf{b} \geq \mathbf{0}$ .
- Add artificial variables  $w_1, \dots, w_m$  with cost  $M$  ( $M$  large).
- We get an initial BFS to the following problem **LP(M)** given by,  
Min  $\mathbf{c}^T \mathbf{x} + [M, \dots, M] \mathbf{w}$   
subject to  
 $\mathbf{Ax} + \mathbf{w} = [A : I] \begin{bmatrix} \mathbf{x} \\ \mathbf{w} \end{bmatrix} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \mathbf{w} \geq \mathbf{0},$   
where  $\mathbf{w} = [w_1, \dots, w_m]^T$ .

## Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:  
Min  $\mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .
- Assume  $\mathbf{b} \geq \mathbf{0}$ .
- Add artificial variables  $w_1, \dots, w_m$  with cost  $M$  ( $M$  large).
- We get an initial BFS to the following problem **LP(M)** given by,  
Min  $\mathbf{c}^T \mathbf{x} + [M, \dots, M] \mathbf{w}$   
subject to  
$$\mathbf{Ax} + \mathbf{w} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{w} \end{bmatrix} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \mathbf{w} \geq \mathbf{0},$$
  
where  $\mathbf{w} = [w_1, \dots, w_m]^T$ .
- An initial BFS is  $[\mathbf{0}_{1 \times n}, w_1, \dots, w_m]^T$ .

## Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:  
Min  $\mathbf{c}^T \mathbf{x}$   
subject to  $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ .
- Assume  $\mathbf{b} \geq \mathbf{0}$ .
- Add artificial variables  $w_1, \dots, w_m$  with cost  $M$  ( $M$  large).
- We get an initial BFS to the following problem **LP(M)** given by,  
Min  $\mathbf{c}^T \mathbf{x} + [M, \dots, M] \mathbf{w}$   
subject to
$$\mathbf{Ax} + \mathbf{w} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{w} \end{bmatrix} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \mathbf{w} \geq \mathbf{0},$$
where  $\mathbf{w} = [w_1, \dots, w_m]^T$ .
- An initial BFS is  $[\mathbf{0}_{1 \times n}, w_1, \dots, w_m]^T$ .
- $\mathbf{x} \in \text{Fea}(\mathbf{P}) \Leftrightarrow [\mathbf{x}^T, \mathbf{0}_{1 \times m}]^T \in \text{Fea}(\text{LP}(\mathbf{M}))$ .

- **Case 1:**  $LP(M)$  has an optimal solution.



- **Case 1:**  $\text{LP}(\mathbf{M})$  has an **optimal** solution.
- **Case 1a:**  $[\mathbf{x}_*^T, \mathbf{0}_{1 \times m}]^T$  is an **optimal** BFS of  $\text{LP}(\mathbf{M})$  with all the artificial variables as nonbasic variables.

- **Case 1:**  $\text{LP}(\mathbf{M})$  has an **optimal** solution.
- **Case 1a:**  $[\mathbf{x}_*^T, \mathbf{0}_{1 \times m}]^T$  is an **optimal** BFS of  $\text{LP}(\mathbf{M})$  with all the artificial variables as nonbasic variables.  
Then  $\mathbf{x}_*$  is an **optimal** BFS for  $\mathbf{P}$ .

- **Case 1:**  $\text{LP}(\mathbf{M})$  has an **optimal** solution.
- **Case 1a:**  $[\mathbf{x}_*^T, \mathbf{O}_{1 \times m}]^T$  is an **optimal** BFS of  $\text{LP}(\mathbf{M})$  with all the artificial variables as nonbasic variables.  
Then  $\mathbf{x}_*$  is an **optimal** BFS for  $\mathbf{P}$ .
- **Case 1b:**  $[\mathbf{x}_*^T, \mathbf{w}_*^T]^T$  is optimal for  $\text{LP}(\mathbf{M})$  and  $\mathbf{w}_* \neq \mathbf{O}$ .

- **Case 1:**  $\text{LP}(\mathbf{M})$  has an **optimal** solution.
- **Case 1a:**  $[\mathbf{x}_*^T, \mathbf{0}_{1 \times m}]^T$  is an **optimal** BFS of  $\text{LP}(\mathbf{M})$  with all the artificial variables as nonbasic variables.  
Then  $\mathbf{x}_*$  is an **optimal** BFS for  $\mathbf{P}$ .
- **Case 1b:**  $[\mathbf{x}_*^T, \mathbf{w}_*^T]^T$  is optimal for  $\text{LP}(\mathbf{M})$  and  $\mathbf{w}_* \neq \mathbf{0}$ .  
Then  $\mathbf{P}$  **does not** have a **feasible solution**.

- **Case 1:**  $\text{LP}(\mathbf{M})$  has an **optimal** solution.
- **Case 1a:**  $[\mathbf{x}_*^T, \mathbf{0}_{1 \times m}]^T$  is an **optimal** BFS of  $\text{LP}(\mathbf{M})$  with all the artificial variables as nonbasic variables.  
Then  $\mathbf{x}_*$  is an **optimal** BFS for  $\mathbf{P}$ .
- **Case 1b:**  $[\mathbf{x}_*^T, \mathbf{w}_*^T]^T$  is optimal for  $\text{LP}(\mathbf{M})$  and  $\mathbf{w}_* \neq \mathbf{0}$ .  
Then  $\mathbf{P}$  **does not** have a **feasible solution**.
- **Case 2:**  $\text{LP}(\mathbf{M})$  **does not** have an **optimal** solution.

- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\text{LP}(\mathbf{M})$ .

- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .
- Then there exists a **direction**  $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$  of  $\text{Fea}(\mathbf{LP}(\mathbf{M}))$  such that

- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .
- Then there exists a **direction**  $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$  of  $\text{Fea}(\mathbf{LP}(\mathbf{M}))$  such that  $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$ .



- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .
- Then there exists a **direction**  $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$  of  $\text{Fea}(\mathbf{LP}(\mathbf{M}))$  such that  $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$ .
- Then  $\mathbf{d}_1$  is a **direction** of  $\text{Fea}(\mathbf{P})$  and  $\mathbf{c}^T \mathbf{d}_1 < 0$ .

- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .
- Then there exists a **direction**  $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$  of  $\text{Fea}(\mathbf{LP}(\mathbf{M}))$  such that  $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$ .
- Then  $\mathbf{d}_1$  is a **direction** of  $\text{Fea}(\mathbf{P})$  and  $\mathbf{c}^T \mathbf{d}_1 < 0$ .  
Hence  $\mathbf{P}$  does **not** have an **optimal** solution but  $\mathbf{P}$  has a **feasible** solution.

- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .
- Then there exists a **direction**  $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$  of  $\text{Fea}(\mathbf{LP}(\mathbf{M}))$  such that  $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$ .
- Then  $\mathbf{d}_1$  is a **direction** of  $\text{Fea}(\mathbf{P})$  and  $\mathbf{c}^T \mathbf{d}_1 < 0$ .  
Hence  $\mathbf{P}$  does **not** have an **optimal** solution but  $\mathbf{P}$  has a **feasible** solution.
- **Case 2b:** In some iteration (or simplex table),  $c_k - z_k = \min\{c_j - z_j : c_j - z_j < 0\}$ ,  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} \neq \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .

- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .
- Then there exists a **direction**  $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$  of  $\text{Fea}(\mathbf{LP}(\mathbf{M}))$  such that  $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$ .
- Then  $\mathbf{d}_1$  is a **direction** of  $\text{Fea}(\mathbf{P})$  and  $\mathbf{c}^T \mathbf{d}_1 < 0$ .  
Hence  $\mathbf{P}$  does **not** have an **optimal** solution but  $\mathbf{P}$  has a **feasible** solution.
- **Case 2b:** In some iteration (or simplex table),  $c_k - z_k = \min\{c_j - z_j : c_j - z_j < 0\}$ ,  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} \neq \mathbf{0}$  in the corresponding BFS for  $\mathbf{LP}(\mathbf{M})$ .  
Then  $\mathbf{P}$  does **not** have a **feasible** solution.

- **Case 2a:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} = \mathbf{0}$  in the corresponding BFS for  $\text{LP}(\mathbf{M})$ .
- Then there exists a **direction**  $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$  of  $\text{Fea}(\text{LP}(\mathbf{M}))$  such that  $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$ .
- Then  $\mathbf{d}_1$  is a **direction** of  $\text{Fea}(\mathbf{P})$  and  $\mathbf{c}^T \mathbf{d}_1 < 0$ .  
Hence  $\mathbf{P}$  does **not** have an **optimal** solution but  $\mathbf{P}$  has a **feasible** solution.
- **Case 2b:** In some iteration (or simplex table),  $c_k - z_k = \min\{c_j - z_j : c_j - z_j < 0\}$ ,  $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$  and  $\mathbf{w} \neq \mathbf{0}$  in the corresponding BFS for  $\text{LP}(\mathbf{M})$ .  
Then  $\mathbf{P}$  does **not** have a **feasible** solution.
- The above conclusion of **Case 2b** that  $\mathbf{P}$  is **infeasible** may not be true if  $c_k - z_k$  is **not** the **most negative** among the  $c_j - z_j$  values.

- **Case 2c:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{a}_k \leq 0$  for **LP(M)**.

- **Case 2c:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{a}_k \leq \mathbf{0}$  for  $\mathbf{LP}(\mathbf{M})$ .  
Then  $\mathbf{P}$  does **not** have an **optimal** solution (may be **infeasible**, may not be **infeasible**).

- **Case 2c:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{a}_k \leq \mathbf{0}$  for  $\mathbf{LP}(\mathbf{M})$ .

Then  $\mathbf{P}$  does **not** have an **optimal** solution (may be **infeasible**, may not be **infeasible**).

- **Remark 1:** If at some stage of solving the artificial variables **leave** the basis, then the corresponding **BFS** is a **BFS** of  $\mathbf{P}$ .



- **Case 2c:** In some iteration (or simplex table), there exists a  $k$  such that  $c_k - z_k < 0$ , the corresponding column  $B^{-1}\tilde{a}_k \leq \mathbf{0}$  for **LP(M)**.

Then **P** does **not** have an **optimal** solution (may be **infeasible**, may not be **infeasible**).

- **Remark 1:** If at some stage of solving the artificial variables **leave** the basis, then the corresponding **BFS** is a **BFS** of **P**.

Then **delete** all columns corresponding to the **artificial variables** and continue.

- **Example 1** : Consider the problem **P**,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- **Example 1** : Consider the problem **P**,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- **Example 1** : Consider the problem **P**,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- If we consider the initial basic solution with basic variables  $s_1$  and  $s_2$ , then the  $c_j - z_j$  values are not  $\geq 0$  for all  $j$ .

- **Example 1** : Consider the problem **P**,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- If we consider the initial basic solution with basic variables  $s_1$  and  $s_2$ , then the  $c_j - z_j$  values are not  $\geq 0$  for all  $j$ .  
We do **not** have a **feasible solution** of the **dual** of **P** at hand.

- **Example 1** : Consider the problem **P**,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize  $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- If we consider the initial basic solution with basic variables  $s_1$  and  $s_2$ , then the  $c_j - z_j$  values are not  $\geq 0$  for all  $j$ .  
We do **not** have a **feasible solution** of the **dual** of **P** at hand.
- The (**Big-M**) is used which provides an **initial BFS** of **P** for Simplex algorithm.

- Consider the modified problem

Minimize  $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

- Consider the modified problem

Minimize  $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

Here  $w$  is called the artificial variable and cost  $M$  associated with it is **very large**.



- Consider the modified problem

Minimize  $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

Here  $w$  is called the artificial variable and cost  $M$  associated with it is **very large**.

- The initial table corresponding to the basic variables  $w$  and  $s_2$  is given below.

$c_j - z_j$	$1 - 2M$	$-M - 1$	$M$	$0$	$0$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$w$	2	1	-1	0	1	4
$s_2$	1	-1	0	1	0	1

- Consider the modified problem

Minimize  $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

Here  $w$  is called the artificial variable and cost  $M$  associated with it is **very large**.

- The initial table corresponding to the basic variables  $w$  and  $s_2$  is given below.

$c_j - z_j$	$1 - 2M$	$-M - 1$	$M$	$0$	$0$	
	$B^{-1}\tilde{a}_1$	$B^{-1}\tilde{a}_2$	$B^{-1}s_1$	$B^{-1}s_2$	$B^{-1}w$	$B^{-1}b$
$w$	2	1	-1	0	1	4
$s_2$	1	-1	0	1	0	1

- $x_1$  will be the **entering variable** and  $s_2$  will be the **leaving variable** for the next table.



$c_j - z_j$	0	$-3M$	$M$	$2M - 1$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$w$	0	3	-1	-2	1	2
$x_1$	1	-1	0	1	0	1



$c_j - z_j$	0	$-3M$	$M$	$2M - 1$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$w$	0	3	-1	-2	1	2
$x_1$	1	-1	0	1	0	1

- $x_2$  will be the **entering variable** and the artificial variable  $w$  will be the **leaving variable**.



$c_j - z_j$	0	$-3M$	$M$	$2M - 1$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$w$	0	3	-1	-2	1	2
$x_1$	1	-1	0	1	0	1

- $x_2$  will be the **entering variable** and the artificial variable  $w$  will be the **leaving variable**.



$c_j - z_j$	0	0	0	-1	$M$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$x_2$	0	1				$\frac{2}{3}$
$x_1$	1	0				$1 + \frac{2}{3}$

- Continue with the BFS corresponding to the basic vectors  $x_1$  and  $x_2$  and drop the column corresponding to the artificial variable  $w$  from all future calculations.

- Continue with the BFS corresponding to the basic vectors  $x_1$  and  $x_2$  and drop the column corresponding to the artificial variable  $w$  from all future calculations.  
Now proceed with **simplex method**.

- Continue with the BFS corresponding to the basic vectors  $x_1$  and  $x_2$  and drop the column corresponding to the artificial variable  $w$  from all future calculations.  
Now proceed with **simplex method**.

- Example 2:** Consider the problem,

Minimize  $x_1 + x_2$

subject to

$$x_1 + 2x_2 \leq 2$$

$$3x_1 + 5x_2 \geq 15$$

$$x_1 \geq 0, x_2 \geq 0.$$



- Continue with the BFS corresponding to the basic vectors  $x_1$  and  $x_2$  and drop the column corresponding to the artificial variable  $w$  from all future calculations.  
Now proceed with **simplex method**.

- Example 2:** Consider the problem,  
Minimize  $x_1 + x_2$   
subject to

$$x_1 + 2x_2 \leq 2$$

$$3x_1 + 5x_2 \geq 15$$

$$x_1 \geq 0, x_2 \geq 0.$$

We consider the corresponding modified problem

- Minimize  $x_1 + x_2 + Mw$   
subject to
- $$x_1 + 2x_2 + s_1 = 2$$
- $$3x_1 + 5x_2 - s_2 + w = 15$$
- $$x_1, x_2, s_1, s_2, w \geq 0.$$

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- The initial table is given by

$c_j - z_j$	$1 - 3M$	$1 - 5M$	$0$	$M$	$0$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$s_1$	1	2	1	0	0	2
$w$	3	5	0	-1	1	15

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- The initial table is given by

$c_j - z_j$	$1 - 3M$	$1 - 5M$	$0$	$M$	$0$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$s_1$	1	2	1	0	0	2
$w$	3	5	0	-1	1	15

- Here  $x_2$  is the **entering variable** and  $s_1$  the **leaving variable**, hence the next table is given by,

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- The initial table is given by

$c_j - z_j$	$1 - 3M$	$1 - 5M$	$0$	$M$	$0$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$s_1$	1	2	1	0	0	2
$w$	3	5	0	-1	1	15

- Here  $x_2$  is the **entering variable** and  $s_1$  the **leaving variable**, hence the next table is given by,



$c_j - z_j$	$\frac{1}{2}(1 - M)$	$0$	$\frac{1}{2}(5M - 1)$	$M$	$0$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$x_2$	$\frac{1}{2}$	1	$\frac{1}{2}$	0	0	1
$w$	$\frac{1}{2}$	0	$-\frac{5}{2}$	-1	1	10

- Here  $x_1$  is the entering variable and  $x_2$  the leaving variable and the next table is given by

- Here  $x_1$  is the entering variable and  $x_2$  the leaving variable and the next table is given by

$c_j - z_j$	0	$M - 1$	$3M - 1$	$M$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$x_1$	1	2	1	0	0	2
$w$	0	-1	-3	-1	1	9

- Here  $x_1$  is the entering variable and  $x_2$  the leaving variable and the next table is given by

$c_j - z_j$	0	$M - 1$	$3M - 1$	$M$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
$x_1$	1	2	1	0	0	2
$w$	0	-1	-3	-1	1	9

- Since the above optimal table has an artificial variable taking positive value hence the original problem ( without the artificial variable ) has no feasible solution.



- If we solve the above problem by the **Dual simplex method** then the problem is written as:

- If we solve the above problem by the **Dual simplex method** then the problem is written as:

Minimize  $x_1 + x_2$

subject to

$$x_1 + 2x_2 + s_1 = 2$$

$$-3x_1 - 5x_2 + s_2 = -15$$

$$x_1, x_2, s_1, s_2 \geq 0.$$

- If we solve the above problem by the **Dual simplex method** then the problem is written as:

Minimize  $x_1 + x_2$

subject to

$$x_1 + 2x_2 + s_1 = 2$$

$$-3x_1 - 5x_2 + s_2 = -15$$

$$x_1, x_2, s_1, s_2 \geq 0.$$

The initial table is given by:

$c_j - z_j$	1	1	0	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
$s_1$	1	2	1	0	2
$s_2$	-3	-5	0	1	-15

- If we solve the above problem by the **Dual simplex method** then the problem is written as:

Minimize  $x_1 + x_2$

subject to

$$x_1 + 2x_2 + s_1 = 2$$

$$-3x_1 - 5x_2 + s_2 = -15$$

$$x_1, x_2, s_1, s_2 \geq 0.$$

The initial table is given by:

$c_j - z_j$	1	1	0	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
$s_1$	1	2	1	0	2
$s_2$	-3	-5	0	1	-15

Here  $s_2$  is the **leaving variable** and  $x_2$  the **entering variable**.

- The next table is given by,

- The next table is given by,

$c_j - z_j$	$\frac{2}{5}$	0	0	$\frac{1}{5}$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
$s_1$	$-\frac{1}{5}$	0	1	$\frac{2}{5}$	-4
$x_2$	$\frac{3}{5}$	1	0	$-\frac{1}{5}$	3

- The next table is given by,

$c_j - z_j$	$\frac{2}{5}$	0	0	$\frac{1}{5}$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
$s_1$	$-\frac{1}{5}$	0	1	$\frac{2}{5}$	-4
$x_2$	$\frac{3}{5}$	1	0	$-\frac{1}{5}$	3

- Here  $s_1$  is the leaving variable and  $x_1$  the entering variable. The next table is given by:

$c_j - z_j$	0	0			
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
$x_1$	1	0	-5	-2	20
$x_2$	0	1	3	1	-9

- The next table is given by,

$c_j - z_j$	$\frac{2}{5}$	0	0	$\frac{1}{5}$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
$s_1$	$-\frac{1}{5}$	0	1	$\frac{2}{5}$	-4
$x_2$	$\frac{3}{5}$	1	0	$-\frac{1}{5}$	3

- Here  $s_1$  is the leaving variable and  $x_1$  the entering variable. The next table is given by:

$c_j - z_j$	0	0			
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
$x_1$	1	0	-5	-2	20
$x_2$	0	1	3	1	-9

Hence clearly the problem is infeasible.