

MA 372 : Stochastic Calculus for Finance

July - November 2023

Department of Mathematics, Indian Institute of Technology Guwahati

Exercises 4

October 12, 2023

1. We shall call $f(t), t \in [0, T]$ a simple process if there is a finite sequence of numbers $0 = t_0 < t_1 < \dots < t_n = T$ and square integrable random variables $\eta_0, \eta_1, \dots, \eta_{n-1}$ such that $f(t, w) = \sum_{j=0}^{n-1} \eta_j(w) \mathbb{I}_{[t_j, t_{j+1})}(t)$, where η_j is \mathcal{F}_{t_j} -measurable. The set of simple processes will be denoted by $M_{step}^2([0, T] \times \Omega)$
 - a) Show that $M_{step}^2([0, T] \times \Omega)$ is a vector space, that is, $af + bg \in M_{step}^2([0, T] \times \Omega)$ for any $f, g \in M_{step}^2([0, T] \times \Omega)$ and $a, b \in \mathbb{R}$.
 - b) Show that $I : M_{step}^2([0, T] \times \Omega) \rightarrow L^2$ is a linear map, i.e., for any $f, g \in M_{step}^2([0, T] \times \Omega)$ and $a, b \in \mathbb{R}$

$$I(af + bg) = aI(f) + bI(g).$$

- c) For any $f, g \in M_{step}^2([0, T] \times \Omega)$

$$E[I(f)I(g)] = E\left[\int_0^T f(t)g(t)dt\right]$$

2. Check whether the following processes $X(t)$ are martingale with respect to Brownian filtration
 - a) $X(t) = W(t) + 4t$ b) $X(t) = W^2(t)$
 - c) $X(t) = t^2W(t) - 2 \int_0^t sW(s)ds$
3. Use Ito's formula to prove that the following stochastic process are martingale with respect to Brownian filtration
 - a) $X(t) = e^{\frac{t}{2}} \cos W(t)$ b) $X(t) = e^{\frac{t}{2}} \sin W(t)$
 - c) $X(t) = e^{W(t) - \frac{t}{2}}$ d) $X(t) = (W(t) + t)e^{-W(t) - \frac{t}{2}}$
4. Define $\beta_k(t) = \mathbb{E}[W^k(t)]$; $k = 0, 1, 2, \dots$; $t \geq 0$
Use Ito's formula to prove that

$$\beta_k(t) = \frac{1}{2}k(k-1) \int_0^t \beta_{k-2}(s)ds; \quad k \geq 2$$

- a) Deduce that $\mathbb{E}[W^4(t)] = 3t^2$ and find $\mathbb{E}[W^6(t)]$.
 - b) Show that $\mathbb{E}[W^{2k+1}(t)] = 0$ and $\mathbb{E}[W^{2k}(t)] = \frac{(2k)!t^k}{2^k k!}$
5. For c, α constants, define

$$X(t) = e^{ct + \alpha W(t)}.$$

Prove that

$$dX(t) = (c + \frac{1}{2}\alpha^2)X(t)dt + \alpha X(t)dW(t)$$

6. Let $\Pi = \{t_0, t_1, \dots, t_n\}$ be a partition of $[0, T]$ with $0 = t_0 < t_1 < \dots < t_n = T$. For $\alpha \in [0, 1]$, consider the sum

$$S_\alpha(\Pi) = \sum_{j=0}^{n-1} \left[(1 - \alpha)W(t_j) + \alpha W(t_{j+1}) \right] (W(t_{j+1}) - W(t_j)).$$

Evaluate the limit $\lim_{\|\Pi\| \rightarrow 0} S_\alpha(\Pi)$ (in L^2), where $\|\pi\| = \max_{j=1,2,\dots,n} (t_j - t_{j-1})$.

7. If $f(t, x) = e^{t/2}(\sin x + \cos x)$, then check whether the process $f(t, W(t))$ is a martingale with respect to Brownian filtration.
8. Suppose f, g are square integrable, deterministic (non random) functions and that there exist constants C, D such that

$$C + \int_0^T f(s) dW(s) = D + \int_0^T g(s) dW(s), \text{ a.e. } \omega \in \Omega.$$

- (i) What is the relationship between C and D ?
- (ii) What is the relationship between $f(t)$ and $g(t)$?

9. If $f(t, x) = x^5 - 10tx^3 + 15t^2x$, then check whether the process $f(t, W(t))$ is a martingale with respect to Brownian filtration.
10. Suppose that $\{W(t); t \geq 0\}$ is a standard Brownian motion with $W(0) = 0$. Determine an expression for

$$\int_0^t \sin(W(s)) dW(s)$$

that does not involve Ito integrals.

11. Suppose $f(t)$ is a deterministic function. Let $X(t) = X(0) + \int_0^t f(s) dW(s)$. Determine an expression for

$$\int_0^t f(s) X(s) dW(s)$$

that does not involve Ito integrals.

12. Let

$$X(t) = \int_0^t W(s) ds$$

for $t \geq 0$. Find the mean, variance and distribution function of the random variable $X(2)$.