

FJ conditions and Karush Kuhn Tucker (KKT) conditions:

- Consider the following nonlinear programming problem (P) of the form,
- Minimize $f(\mathbf{x})$

FJ conditions and Karush Kuhn Tucker (KKT) conditions:

- Consider the following nonlinear programming problem (P) of the form,
- Minimize $f(\mathbf{x})$
- subject to $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, \quad \mathbf{x} \in \mathbb{R}^n.$

FJ conditions and Karush Kuhn Tucker (KKT) conditions:

- Consider the following nonlinear programming problem (P) of the form,
- Minimize $f(\mathbf{x})$
- subject to $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, \quad \mathbf{x} \in \mathbb{R}^n.$
- $f, g_i : \mathbb{R}^n \rightarrow \mathbb{R}.$

FJ conditions and Karush Kuhn Tucker (KKT) conditions:

- Consider the following nonlinear programming problem (P) of the form,
- Minimize $f(\mathbf{x})$
- subject to $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, \quad \mathbf{x} \in \mathbb{R}^n.$
- $f, g_i : \mathbb{R}^n \rightarrow \mathbb{R}.$
- f, g_i may not be linear functions.

FJ conditions and Karush Kuhn Tucker (KKT) conditions:

- Consider the following nonlinear programming problem (P) of the form,
- Minimize $f(\mathbf{x})$
- subject to $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, \quad \mathbf{x} \in \mathbb{R}^n.$
- $f, g_i : \mathbb{R}^n \rightarrow \mathbb{R}.$
- f, g_i may not be linear functions.
- $S = \text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, \text{ for } i = 1, \dots, m\}.$

- The set **D** of **feasible directions** at $\mathbf{x}^* \in \text{Fea}(P)$ is,
 $\mathbf{D}_{\mathbf{x}^*} = \{\mathbf{d} \in \mathbb{R}^n : \exists \ c > 0 \text{ such that } g_i(\mathbf{x}^* + t\mathbf{d}) \leq 0, \text{ for all } i = 1, \dots, m, \text{ for all } 0 \leq t \leq c\}.$

- The set **D** of **feasible directions** at $\mathbf{x}^* \in \text{Fea}(P)$ is,
 $\mathbf{D}_{\mathbf{x}^*} = \{\mathbf{d} \in \mathbb{R}^n : \exists c > 0 \text{ such that } g_i(\mathbf{x}^* + t\mathbf{d}) \leq 0, \text{ for all } i = 1, \dots, m, \text{ for all } 0 \leq t \leq c\}.$
- Let $I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) = 0\}.$

- The set **D** of **feasible directions** at $\mathbf{x}^* \in \text{Fea}(P)$ is,
 $\mathbf{D}_{\mathbf{x}^*} = \{\mathbf{d} \in \mathbb{R}^n : \exists c > 0 \text{ such that } g_i(\mathbf{x}^* + t\mathbf{d}) \leq 0, \text{ for all } i = 1, \dots, m, \text{ for all } 0 \leq t \leq c\}.$
- Let $I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) = 0\}.$
- Let $I^* = \{1, \dots, m\} \setminus I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) < 0\}.$

- The set **D** of **feasible directions** at $\mathbf{x}^* \in \text{Fea}(P)$ is,
 $\mathbf{D}_{\mathbf{x}^*} = \{\mathbf{d} \in \mathbb{R}^n : \exists c > 0 \text{ such that } g_i(\mathbf{x}^* + t\mathbf{d}) \leq 0, \text{ for all } i = 1, \dots, m, \text{ for all } 0 \leq t \leq c\}.$
- Let $I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) = 0\}.$
- Let $I^* = \{1, \dots, m\} \setminus I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) < 0\}.$
- For all $i \in I^*$, let g_i be **continuous** at \mathbf{x}^* .

- The set **D** of **feasible directions** at $\mathbf{x}^* \in \text{Fea}(P)$ is,
 $\mathbf{D}_{\mathbf{x}^*} = \{\mathbf{d} \in \mathbb{R}^n : \exists \ c > 0 \text{ such that } g_i(\mathbf{x}^* + t\mathbf{d}) \leq 0, \text{ for all } i = 1, \dots, m, \text{ for all } 0 \leq t \leq c\}.$
- Let $I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) = 0\}.$
- Let $I^* = \{1, \dots, m\} \setminus I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) < 0\}.$
- For all $i \in I^*$, let g_i be **continuous** at \mathbf{x}^* .
- For each $i \in I^*$ there exists $c_i > 0$ such that

- The set **D** of **feasible directions** at $\mathbf{x}^* \in \text{Fea}(P)$ is,
 $\mathbf{D}_{\mathbf{x}^*} = \{\mathbf{d} \in \mathbb{R}^n : \exists \ c > 0 \text{ such that } g_i(\mathbf{x}^* + t\mathbf{d}) \leq 0, \text{ for all } i = 1, \dots, m, \text{ for all } 0 \leq t \leq c\}.$
- Let $I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) = 0\}.$
- Let $I^* = \{1, \dots, m\} \setminus I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) < 0\}.$
- For all $i \in I^*$, let g_i be **continuous** at \mathbf{x}^* .
- For each $i \in I^*$ there exists $c_i > 0$ such that $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq c_i$,

- The set **D** of **feasible directions** at $\mathbf{x}^* \in \text{Fea}(P)$ is,
 $\mathbf{D}_{\mathbf{x}^*} = \{\mathbf{d} \in \mathbb{R}^n : \exists c > 0 \text{ such that } g_i(\mathbf{x}^* + t\mathbf{d}) \leq 0, \text{ for all } i = 1, \dots, m, \text{ for all } 0 \leq t \leq c\}.$
- Let $I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) = 0\}.$
- Let $I^* = \{1, \dots, m\} \setminus I = \{i \in \{1, \dots, m\} : g_i(\mathbf{x}^*) < 0\}.$
- For all $i \in I^*$, let g_i be **continuous** at \mathbf{x}^* .
- For each $i \in I^*$ there exists $c_i > 0$ such that
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0, \text{ for all } 0 \leq t \leq c_i,$
- If $c = \min_{i \in I^*} \{c_i\}$, then
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0, \text{ for all } 0 \leq t \leq c, \text{ for all } i \in I^*.$

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .
- If \mathbf{d} is such that, $\nabla g_i(\mathbf{x}^*)\mathbf{d} < 0$, for all $i \in I$,

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .
- If \mathbf{d} is such that, $\nabla g_i(\mathbf{x}^*)\mathbf{d} < 0$, for all $i \in I$,
then for each $i \in I$ there exists an $a_i > 0$ such that
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a_i$.

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .
- If \mathbf{d} is such that, $\nabla g_i(\mathbf{x}^*)\mathbf{d} < 0$, for all $i \in I$,
then for each $i \in I$ there exists an $a_i > 0$ such that
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a_i$.
- If $a = \min_{i \in I} \{a_i\} > 0$, $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a$, for
all $i \in I$.

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .
- If \mathbf{d} is such that, $\nabla g_i(\mathbf{x}^*)\mathbf{d} < 0$, for all $i \in I$,
then for each $i \in I$ there exists an $a_i > 0$ such that
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a_i$.
- If $a = \min_{i \in I} \{a_i\} > 0$, $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a$, for
all $i \in I$.
- If for all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .
- If \mathbf{d} is such that, $\nabla g_i(\mathbf{x}^*)\mathbf{d} < 0$, for all $i \in I$,
then for each $i \in I$ there exists an $a_i > 0$ such that
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a_i$.
- If $a = \min_{i \in I} \{a_i\} > 0$, $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a$, for
all $i \in I$.
- If for all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*
and for all $i \in I$, g_i is **continuously differentiable** at \mathbf{x}^*
then

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .
- If \mathbf{d} is such that, $\nabla g_i(\mathbf{x}^*)\mathbf{d} < 0$, for all $i \in I$,
then for each $i \in I$ there exists an $a_i > 0$ such that
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a_i$.
- If $a = \min_{i \in I} \{a_i\} > 0$, $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a$, for
all $i \in I$.
- If for all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*
and for all $i \in I$, g_i is **continuously differentiable** at \mathbf{x}^*
then
 $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$,

- For $i \in I$, let g_i be **continuously differentiable** at \mathbf{x}^* .
- If \mathbf{d} is such that, $\nabla g_i(\mathbf{x}^*)\mathbf{d} < 0$, for all $i \in I$,
then for each $i \in I$ there exists an $a_i > 0$ such that
 $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a_i$.
- If $a = \min_{i \in I} \{a_i\} > 0$, $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$, for all $0 \leq t \leq a$, for
all $i \in I$.
- If for all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*
and for all $i \in I$, g_i is **continuously differentiable** at \mathbf{x}^*
then
 $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$,
where $\mathbf{G}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla g_i(\mathbf{x}^*)\mathbf{d} < 0 \text{ for all } i \in I\}$.

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^*

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.
Consider the problem **P** of minimizing f subject to
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.
Consider the problem **P** of minimizing f subject to
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .
Let $S = \text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and
 $\mathbf{x}^* \in S$.

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.
Consider the problem **P** of minimizing f subject to
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .
Let $S = \text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and
 $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*

- If \mathbf{x}^* is a local **minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.
Consider the problem **P** of minimizing f subject to
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .
Let $S = \text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and
 $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*
and for all $i \in I$, g_i is **continuously differentiable** at \mathbf{x}^* .

- If \mathbf{x}^* is a **local minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.
Consider the problem **P** of minimizing f subject to
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .
Let $S = \text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and
 $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*
and for all $i \in I$, g_i is **continuously differentiable** at \mathbf{x}^* .
If \mathbf{x}^* is a **local minimum** of f over S

- If \mathbf{x}^* is a **local minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.
Consider the problem **P** of minimizing f subject to
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .
Let $S = \text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and
 $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*
and for all $i \in I$, g_i is **continuously differentiable** at \mathbf{x}^* .
If \mathbf{x}^* is a **local minimum** of f over S
then there exists **non negative** constants, $u_0, u_i, i \in I$, **not**
all zeros such that

- If \mathbf{x}^* is a **local minimum**, then $\mathbf{F}_0 \cap \mathbf{D}_{\mathbf{x}^*} = \phi$,
where $\mathbf{D}_{\mathbf{x}^*}$ is the set of feasible directions at \mathbf{x}^* and
 $\mathbf{F}_0 = \{\mathbf{d} \in \mathbb{R}^n : \nabla f(\mathbf{x}^*)\mathbf{d} < 0\}$.
- Since $\mathbf{G}_0 \subseteq \mathbf{D}_{\mathbf{x}^*}$, \mathbf{x}^* is a **local minimum** implies
 $\mathbf{F}_0 \cap \mathbf{G}_0 = \phi$.
- **Theorem 7: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be
a **continuously differentiable** function.
Consider the problem **P** of minimizing f subject to
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .
Let $S = \text{Fea}(P) = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and
 $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i is **continuous** at \mathbf{x}^*
and for all $i \in I$, g_i is **continuously differentiable** at \mathbf{x}^* .
If \mathbf{x}^* is a **local minimum** of f over S
then there exists **non negative** constants, $u_0, u_i, i \in I$, **not**
all zeros such that
$$u_0 \nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (*)$$

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$,

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$,

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.
At $\mathbf{x}^* = [0, 2]^T$ the binding constraint are g_2 and g_3

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.
At $\mathbf{x}^* = [0, 2]^T$ the binding constraints are g_2 and g_3
 $\nabla g_2(\mathbf{x}^*) = [1, 2]$,

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.
At $\mathbf{x}^* = [0, 2]^T$ the binding constraint are g_2 and g_3
 $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla g_3(\mathbf{x}^*) = [-1, 0]$,

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.
At $\mathbf{x}^* = [0, 2]^T$ the binding constraint are g_2 and g_3
 $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla g_3(\mathbf{x}^*) = [-1, 0]$, $\nabla f(\mathbf{x}^*) = [-6, 0]$.

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.
At $\mathbf{x}^* = [0, 2]^T$ the binding constraint are g_2 and g_3
 $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla g_3(\mathbf{x}^*) = [-1, 0]$, $\nabla f(\mathbf{x}^*) = [-6, 0]$.
 $\nabla g_2(\mathbf{x}^*)$ and $\nabla f(\mathbf{x}^*)$ are LI.

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$
 $-x_2 \leq 0.$
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2], \nabla g_2(\mathbf{x}^*) = [1, 2], \nabla f(\mathbf{x}^*) = [-2, -2].$
- Take $u_0 = 1, u_1 = \frac{1}{3}, u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}.$
- $[2, 1]^T$ satisfies the FJ condition.
 At $\mathbf{x}^* = [0, 2]^T$ the binding constraint are g_2 and g_3
 $\nabla g_2(\mathbf{x}^*) = [1, 2], \nabla g_3(\mathbf{x}^*) = [-1, 0], \nabla f(\mathbf{x}^*) = [-6, 0].$
 $\nabla g_2(\mathbf{x}^*)$ and $\nabla f(\mathbf{x}^*)$ are LI.
- $u_0 \nabla f(\mathbf{x}^*) + u_2 \nabla g_2(\mathbf{x}^*) + u_3 \nabla g_3(\mathbf{x}^*) = \mathbf{0}.$

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.
At $\mathbf{x}^* = [0, 2]^T$ the binding constraint are g_2 and g_3
 $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla g_3(\mathbf{x}^*) = [-1, 0]$, $\nabla f(\mathbf{x}^*) = [-6, 0]$.
 $\nabla g_2(\mathbf{x}^*)$ and $\nabla f(\mathbf{x}^*)$ are LI.
- $u_0 \nabla f(\mathbf{x}^*) + u_2 \nabla g_2(\mathbf{x}^*) + u_3 \nabla g_3(\mathbf{x}^*) = \mathbf{0}$.
has only the trivial solution.

- **Example 1:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
- Take $u_0 = 1$, $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- $[2, 1]^T$ satisfies the FJ condition.
At $\mathbf{x}^* = [0, 2]^T$ the binding constraint are g_2 and g_3
 $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla g_3(\mathbf{x}^*) = [-1, 0]$, $\nabla f(\mathbf{x}^*) = [-6, 0]$.
 $\nabla g_2(\mathbf{x}^*)$ and $\nabla f(\mathbf{x}^*)$ are LI.
- $u_0 \nabla f(\mathbf{x}^*) + u_2 \nabla g_2(\mathbf{x}^*) + u_3 \nabla g_3(\mathbf{x}^*) = \mathbf{0}$.
has only the trivial solution.
Hence $[0, 2]^T$ **does not** satisfy the FJ condition.

- If g_i is **continuously differentiable** at \mathbf{x}^* for all $i = 1, \dots, m$, then

- If g_i is **continuously differentiable** at \mathbf{x}^* for all $i = 1, \dots, m$, then
- $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$

- If g_i is **continuously differentiable** at \mathbf{x}^* for all $i = 1, \dots, m$, then
- $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$
and $u_i g_i(\mathbf{x}^*) = 0$, for all $i = 1, \dots, m$. **(**)**

- If g_i is **continuously differentiable** at \mathbf{x}^* for all $i = 1, \dots, m$, then
- $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$
and $u_i g_i(\mathbf{x}^*) = 0$, for all $i = 1, \dots, m$. (**)
- $\mathbf{x}^* \in S$ is called the **primal feasibility condition**,

- If g_i is **continuously differentiable** at \mathbf{x}^* for all $i = 1, \dots, m$, then
- $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$
and $u_i g_i(\mathbf{x}^*) = 0$, for all $i = 1, \dots, m$. (**)
- $\mathbf{x}^* \in S$ is called the **primal feasibility condition**,
- (*) together with nonnegativity of the u_i 's, is called the **dual feasibility condition**.

- If g_i is **continuously differentiable** at \mathbf{x}^* for all $i = 1, \dots, m$, then
- $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$
and $u_i g_i(\mathbf{x}^*) = 0$, for all $i = 1, \dots, m$. (**)
- $\mathbf{x}^* \in S$ is called the **primal feasibility condition**,
- (*) together with nonnegativity of the u_i 's, is called the **dual feasibility condition**.
- $u_i g_i(\mathbf{x}^*) = 0$, for all $i = 1, \dots, m$, is called the **complementary slackness condition**.

- If g_i is **continuously differentiable** at \mathbf{x}^* for all $i = 1, \dots, m$, then
- $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$
and $u_i g_i(\mathbf{x}^*) = 0$, for all $i = 1, \dots, m$. (**)
- $\mathbf{x}^* \in S$ is called the **primal feasibility condition**,
- (*) together with nonnegativity of the u_i 's, is called the **dual feasibility condition**.
- $u_i g_i(\mathbf{x}^*) = 0$, for all $i = 1, \dots, m$, is called the **complementary slackness condition**.
- The above conditions are called the **FJ (Fritz John)** conditions and the point $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is called a **Fritz John**, or an **FJ**, point.

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that
$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$
- The above conditions given by (***) are called **KKT (Karush, Kuhn, Tucker)** conditions.

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$

- The above conditions given by (***) are called **KKT (Karush, Kuhn, Tucker)** conditions.

Any $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) which satisfies the **Karush Kuhn Tucker (KKT)** conditions is called a **KKT point**.

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$

- The above conditions given by (***) are called **KKT (Karush, Kuhn, Tucker)** conditions.

Any $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) which satisfies the **Karush Kuhn Tucker (KKT)** conditions is called a **KKT point**.

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$

- The above conditions given by (***) are called **KKT (Karush, Kuhn, Tucker)** conditions.

Any $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) which satisfies the **Karush Kuhn Tucker (KKT)** conditions is called a **KKT point**.

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI**

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$

- The above conditions given by (***) are called **KKT (Karush, Kuhn, Tucker)** conditions.

Any $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) which satisfies the **Karush Kuhn Tucker (KKT)** conditions is called a **KKT** point.

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI** and \mathbf{x}^* is a local minimum of f then

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$

- The above conditions given by (***) are called **KKT (Karush, Kuhn, Tucker)** conditions.

Any $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) which satisfies the **Karush Kuhn Tucker (KKT)** conditions is called a **KKT** point.

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI** and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that

- **Gordon's Theorem:** Exactly one of the following two systems has a solution:

$$\mathbf{u} \neq \mathbf{0}, \mathbf{u} \geq \mathbf{0}, \mathbf{u}^T \mathbf{A} = \mathbf{0} \quad (1)$$

$$\mathbf{A}\mathbf{y} > \mathbf{0} \quad (2)$$

- \mathbf{x}^* is said to satisfy **KKT** condition if there exists non negative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (***)$$

- The above conditions given by (***) are called **KKT (Karush, Kuhn, Tucker)** conditions.

Any $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) which satisfies the **Karush Kuhn Tucker (KKT)** conditions is called a **KKT point**.

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI**

and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (****)$$

- **Remark:** Hence from **Theorem 8** it is clear that if $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is an **FJ** point and if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI** then $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is a **KKT** point.

- **Remark:** Hence from **Theorem 8** it is clear that if $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is an **FJ** point and if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI** then $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is a **KKT** point.
- **Remark:** If \mathbf{x}^* is a **Karush Kuhn Tucker (KKT)** point then it is also an **FJ** point.

- **Remark:** Hence from **Theorem 8** it is clear that if $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is an **FJ** point and if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI** then $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is a **KKT** point.
- **Remark:** If \mathbf{x}^* is a **Karush Kuhn Tucker (KKT)** point then it is also an **FJ** point.
- **Remark:** **KKT** conditions says that under the assumptions of the above theorem, if \mathbf{x}^* is a **local minimum**,

- **Remark:** Hence from **Theorem 8** it is clear that if $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is an **FJ** point and if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI** then $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is a **KKT** point.
- **Remark:** If \mathbf{x}^* is a **Karush Kuhn Tucker (KKT)** point then it is also an **FJ** point.
- **Remark:** **KKT** conditions says that under the assumptions of the above theorem, if \mathbf{x}^* is a **local minimum**, then $-\nabla f(\mathbf{x}^*)$ lies in the **cone generated** by the $\nabla g_i(\mathbf{x}^*)$'s, $i \in I$.

- **Remark:** Hence from **Theorem 8** it is clear that if $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is an **FJ** point and if $\nabla g_i(\mathbf{x}^*)$'s for $i \in I$ are **LI** then $(\mathbf{x}^*, \mathbf{u})$ (or \mathbf{x}^*) is a **KKT** point.
- **Remark:** If \mathbf{x}^* is a **Karush Kuhn Tucker (KKT)** point then it is also an **FJ** point.
- **Remark:** **KKT** conditions says that under the assumptions of the above theorem, if \mathbf{x}^* is a **local minimum**, then $-\nabla f(\mathbf{x}^*)$ lies in the **cone generated** by the $\nabla g_i(\mathbf{x}^*)$'s, $i \in I$.

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$

subject to

$$x_1^2 + x_2^2 \leq 5.$$

$$x_1 + 2x_2 \leq 4.$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0.$$

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$
 $-x_2 \leq 0.$
- f takes its minimum value at $[2, 1]^T$.

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
$$x_1^2 + x_2^2 \leq 5.$$
$$x_1 + 2x_2 \leq 4.$$
$$-x_1 \leq 0$$
$$-x_2 \leq 0.$$
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5,$
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4,$

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$,

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$,

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
$$x_1^2 + x_2^2 \leq 5.$$
$$x_1 + 2x_2 \leq 4.$$
$$-x_1 \leq 0$$
$$-x_2 \leq 0.$$
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
 $\nabla g_1(\mathbf{x}^*)$ and $\nabla g_2(\mathbf{x}^*)$ are LI.

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
 $\nabla g_1(\mathbf{x}^*)$ and $\nabla g_2(\mathbf{x}^*)$ are LI.
- Take $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
 $\nabla g_1(\mathbf{x}^*)$ and $\nabla g_2(\mathbf{x}^*)$ are LI.
- Take $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.

- **Example 1: revisited** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.
 $x_1 + 2x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
- f takes its minimum value at $[2, 1]^T$.
- $g_1(\mathbf{x}) = x_1^2 + x_2^2 - 5$,
 $g_2(\mathbf{x}) = x_1 + 2x_2 - 4$, $g_3(\mathbf{x}) = -x_1$ and $g_4(\mathbf{x}) = -x_2$.
- At $\mathbf{x}^* = [2, 1]^T$ the binding constraints are g_1 and g_2 .
- $\nabla g_1(\mathbf{x}^*) = [4, 2]$, $\nabla g_2(\mathbf{x}^*) = [1, 2]$, $\nabla f(\mathbf{x}^*) = [-2, -2]$.
 $\nabla g_1(\mathbf{x}^*)$ and $\nabla g_2(\mathbf{x}^*)$ are LI.
- Take $u_1 = \frac{1}{3}$, $u_2 = \frac{2}{3}$, then
- $\nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
Hence $[2, 1]^T$ satisfies the KKT condition.

- **Example 2:** $f(x_1, x_2) = -x_1$.

- **Example 2:** $f(x_1, x_2) = -x_1$.

subject to

$$x_2 - (1 - x_1)^3 \leq 0$$

- **Example 2:** $f(x_1, x_2) = -x_1$.

subject to

$$x_2 - (1 - x_1)^3 \leq 0$$

$$-x_2 \leq 0$$

- **Example 2:** $f(x_1, x_2) = -x_1$.
subject to
 $x_2 - (1 - x_1)^3 \leq 0$
 $-x_2 \leq 0$
- $\mathbf{x}^* = [1, 0]^T$ is a **local minimum**.

- **Example 2:** $f(x_1, x_2) = -x_1$.

subject to

$$x_2 - (1 - x_1)^3 \leq 0$$

$$-x_2 \leq 0$$

- $\mathbf{x}^* = [1, 0]^T$ is a **local minimum**.
- At $[1, 0]^T$ both the constraints are binding.

- **Example 2:** $f(x_1, x_2) = -x_1$.

subject to

$$x_2 - (1 - x_1)^3 \leq 0$$

$$-x_2 \leq 0$$

- $\mathbf{x}^* = [1, 0]^T$ is a **local minimum**.
- At $[1, 0]^T$ both the constraints are binding.
- $\nabla f(\mathbf{x}^*) = [-1, 0]$, $\nabla g_1(\mathbf{x}^*) = [0, 1]$
and $\nabla g_2(\mathbf{x}^*) = [0, -1]$.

- **Example 2:** $f(x_1, x_2) = -x_1$.

subject to

$$x_2 - (1 - x_1)^3 \leq 0$$

$$-x_2 \leq 0$$

- $\mathbf{x}^* = [1, 0]^T$ is a **local minimum**.
- At $[1, 0]^T$ both the constraints are binding.
- $\nabla f(\mathbf{x}^*) = [-1, 0]$, $\nabla g_1(\mathbf{x}^*) = [0, 1]$
and $\nabla g_2(\mathbf{x}^*) = [0, -1]$.
- Take $u_0 = 0$, $u_1 = 1$, $u_2 = 1$, then

- **Example 2:** $f(x_1, x_2) = -x_1$.

subject to

$$x_2 - (1 - x_1)^3 \leq 0$$

$$-x_2 \leq 0$$

- $\mathbf{x}^* = [1, 0]^T$ is a **local minimum**.
- At $[1, 0]^T$ both the constraints are binding.
- $\nabla f(\mathbf{x}^*) = [-1, 0]$, $\nabla g_1(\mathbf{x}^*) = [0, 1]$
and $\nabla g_2(\mathbf{x}^*) = [0, -1]$.
- Take $u_0 = 0$, $u_1 = 1$, $u_2 = 1$, then
 $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.

- **Example 2:** $f(x_1, x_2) = -x_1$.
subject to
 $x_2 - (1 - x_1)^3 \leq 0$
 $-x_2 \leq 0$
- $\mathbf{x}^* = [1, 0]^T$ is a **local minimum**.
- At $[1, 0]^T$ both the constraints are binding.
- $\nabla f(\mathbf{x}^*) = [-1, 0]$, $\nabla g_1(\mathbf{x}^*) = [0, 1]$
and $\nabla g_2(\mathbf{x}^*) = [0, -1]$.
- Take $u_0 = 0$, $u_1 = 1$, $u_2 = 1$, then
 $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- Since $-\nabla f(\mathbf{x}^*)$ does not lie in the cone generated by the $\nabla g_i(\mathbf{x}^*)$'s, $i = 1, 2$,
 $[1, 0]^T$ is an **FJ point** but **not a KKT point**.

- **Example 2:** $f(x_1, x_2) = -x_1$.
subject to
 $x_2 - (1 - x_1)^3 \leq 0$
 $-x_2 \leq 0$
- $\mathbf{x}^* = [1, 0]^T$ is a **local minimum**.
- At $[1, 0]^T$ both the constraints are binding.
- $\nabla f(\mathbf{x}^*) = [-1, 0]$, $\nabla g_1(\mathbf{x}^*) = [0, 1]$
and $\nabla g_2(\mathbf{x}^*) = [0, -1]$.
- Take $u_0 = 0$, $u_1 = 1$, $u_2 = 1$, then
 $u_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^2 u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- Since $-\nabla f(\mathbf{x}^*)$ does not lie in the cone generated by the $\nabla g_i(\mathbf{x}^*)$'s, $i = 1, 2$,
 $[1, 0]^T$ is an **FJ point** but **not a KKT point**.
- So **KKT** condition is **not** a necessary condition for a **local minimum**, although FJ conditions are **necessary conditions** for a **local minimum**.

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \emptyset$

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \emptyset$ and \mathbf{x}^* is a local minimum of f then

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \emptyset$ and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \emptyset$ and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that
$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (**)$$

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \emptyset$ and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that
$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (**)$$
- **Example 2:** Minimize $-x_1$.

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \emptyset$ and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that
$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (**)$$
- **Example 2:** Minimize $-x_1$.
subject to
$$-x_2 - x_1 \leq 0$$

- **Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \emptyset$ and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that
$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (**)$$

- **Example 2:** Minimize $-x_1$.

subject to

$$-x_2 - x_1 \leq 0$$

$$-x_2 \leq 0$$

$$-x_1 \leq 0$$

- Theorem 8: (KKT conditions)** If in addition to the conditions assumed for f and g_i 's, as in the previous theorem, if $G_0 \neq \phi$ and \mathbf{x}^* is a local minimum of f then there exists nonnegative constants $u_i, i \in I$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (**)$$
- Example 2:** Minimize $-x_1$.
 subject to

$$-x_2 - x_1 \leq 0$$

$$-x_2 \leq 0$$

$$-x_1 \leq 0$$
- $\nabla g_i(\mathbf{x}^*)$'s, $i \in I$ are **LD** at $\mathbf{x}^* = [0, 0]^T$ but $G_0 \neq \phi$.
- Remark:** If $\nabla g_i(\mathbf{x}^*)$'s, $i \in I$ are **LI** then $G_0 \neq \phi$ but the converse is **not** true.

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**.

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**. Consider the problem of **minimizing** f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$,

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**. Consider the problem of **minimizing** f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**.

Consider the problem of **minimizing** f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .

Let $S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and $\mathbf{x}^* \in S$.

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**.

Consider the problem of **minimizing** f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .

Let $S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and $\mathbf{x}^* \in S$. For all $i \in I^*$, assume that g_i is **continuous** at \mathbf{x}^* and for all $i \in I$, g_i 's are assumed to be **continuously differentiable** at \mathbf{x}^* .

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**.

Consider the problem of **minimizing** f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .

Let $S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and $\mathbf{x}^* \in S$. For all $i \in I^*$, assume that g_i is **continuous** at \mathbf{x}^* and for all $i \in I$, g_i 's are assumed to be **continuously differentiable** at \mathbf{x}^* .

Let all the g_i 's be **convex functions**, so that $S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, 2, \dots, m\}$ is **convex**.

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**.

Consider the problem of **minimizing** f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .

Let $S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and $\mathbf{x}^* \in S$. For all $i \in I^*$, assume that g_i is **continuous** at \mathbf{x}^* and for all $i \in I$, g_i 's are assumed to be **continuously differentiable** at \mathbf{x}^* .

Let all the g_i 's be **convex functions**, so that

$S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, 2, \dots, m\}$ is **convex**. Then \mathbf{x}^* is a **global minimum** of f over S if

- **Theorem 9:(KKT sufficient conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be **convex** and **continuously differentiable**.

Consider the problem of **minimizing** f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i .

Let $S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, \dots, m\}$ and $\mathbf{x}^* \in S$. For all $i \in I^*$, assume that g_i is **continuous** at \mathbf{x}^* and for all $i \in I$, g_i 's are assumed to be **continuously differentiable** at \mathbf{x}^* .

Let all the g_i 's be **convex functions**, so that $S = \{\mathbf{x} \in \mathbb{R}^n : g_i(\mathbf{x}) \leq 0, i = 1, 2, \dots, m\}$ is **convex**. Then \mathbf{x}^* is a **global minimum** of f over S if there exists non negative constants, $u_i, i \in I$ such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}. \quad (*)$$

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5$.

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$
 $-x_2 \leq 0.$

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$
 $-x_2 \leq 0.$
- Check that f, g_i for all i are convex functions.

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$
 $-x_2 \leq 0.$
- Check that f, g_i for all i are convex functions.
- $\mathbf{x}^* = [2, 1]^T$ is a KKT point.

- **Example 1 revisited:** $f(x_1, x_2) = (x_1 - 3)^2 + (x_2 - 2)^2$
subject to
 $x_1^2 + x_2^2 \leq 5.$
 $x_1 + 2x_2 \leq 4.$
 $-x_1 \leq 0$
 $-x_2 \leq 0.$
- Check that f, g_i for all i are convex functions.
- $\mathbf{x}^* = [2, 1]^T$ is a KKT point.
- $\mathbf{x}^* = [2, 1]^T$ is the global minimum of f .

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.
- For $x^* = 0$, $\nabla f(x^*) = 0$, hence $x^* = 0$ satisfies the **KKT** conditions but x^* not be a local minimum point.

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.
- For $x^* = 0$, $\nabla f(x^*) = 0$, hence $x^* = 0$ satisfies the **KKT** conditions but x^* not be a local minimum point.
- f is clearly **not** convex.

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.
- For $x^* = 0$, $\nabla f(x^*) = 0$, hence $x^* = 0$ satisfies the **KKT** conditions but x^* not be a local minimum point.
- f is clearly **not** convex.
- Any x^* for which $\nabla f(x^*) = 0$ is a KKT point but x^* may not be a point of local minimum.

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.
- For $x^* = 0$, $\nabla f(x^*) = 0$, hence $x^* = 0$ satisfies the **KKT** conditions but x^* not be a local minimum point.
- f is clearly **not** convex.
- Any x^* for which $\nabla f(x^*) = 0$ is a KKT point but x^* may not be a point of local minimum.
- $f(x_1, x_2) = x_1$

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.
- For $x^* = 0$, $\nabla f(x^*) = 0$, hence $x^* = 0$ satisfies the **KKT** conditions but x^* not be a local minimum point.
- f is clearly **not** convex.
- Any x^* for which $\nabla f(x^*) = 0$ is a KKT point but x^* may not be a point of local minimum.
- $f(x_1, x_2) = x_1$
 subject to
 $(x_1 - 1)^2 + (x_2 - 1)^2 \leq 1$

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.
- For $x^* = 0$, $\nabla f(x^*) = 0$, hence $x^* = 0$ satisfies the **KKT** conditions but x^* not be a local minimum point.
- f is clearly **not** convex.
- Any x^* for which $\nabla f(x^*) = 0$ is a KKT point but x^* may not be a point of local minimum.
- $f(x_1, x_2) = x_1$
subject to
 $(x_1 - 1)^2 + (x_2 - 1)^2 \leq 1$
 $(x_1 - 1)^2 + (x_2 + 1)^2 \leq 1$

- **Example 3:** $f(x) = -x^2$ for $x \leq 0$
 $= x^2$ for $x \geq 0$.
- For $x^* = 0$, $\nabla f(x^*) = 0$, hence $x^* = 0$ satisfies the **KKT** conditions but x^* not be a local minimum point.
- f is clearly **not** convex.
- Any x^* for which $\nabla f(x^*) = 0$ is a KKT point but x^* may not be a point of local minimum.
- $f(x_1, x_2) = x_1$
 subject to
 $(x_1 - 1)^2 + (x_2 - 1)^2 \leq 1$
 $(x_1 - 1)^2 + (x_2 + 1)^2 \leq 1$
 $x^* = [1, 0]^T$ is the **global minimum** of f but not a KKT point.

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ are given by:

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ are given by:

- $\mathbf{c}^T - \mathbf{y}^T A - \mathbf{v}^T = \mathbf{0}$, or $\mathbf{c}^T - \mathbf{y}^T A = \mathbf{v}^T$

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ are given by:

- $\mathbf{c}^T - \mathbf{y}^T A - \mathbf{v}^T = \mathbf{0}$, or $\mathbf{c}^T - \mathbf{y}^T A = \mathbf{v}^T$
and $v_i x_i = 0$ for all $i = 1, \dots, n$.

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$ are given by:

- $\mathbf{c}^T - \mathbf{y}^T A - \mathbf{v}^T = \mathbf{0}$, or $\mathbf{c}^T - \mathbf{y}^T A = \mathbf{v}^T$

and $v_i x_i = 0$ for all $i = 1, \dots, n$.

where $\mathbf{v} = [v_1, \dots, v_n]^T$ is a non negative vector,

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$ are given by:

- $\mathbf{c}^T - \mathbf{y}^T A - \mathbf{v}^T = \mathbf{0}$, or $\mathbf{c}^T - \mathbf{y}^T A = \mathbf{v}^T$

and $v_i x_i = 0$ for all $i = 1, \dots, n$.

where $\mathbf{v} = [v_1, \dots, v_n]^T$ is a non negative vector,

and $\mathbf{y} = [y_1, \dots, y_m]^T$ is unrestricted in sign.

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$ are given by:

- $\mathbf{c}^T - \mathbf{y}^T A - \mathbf{v}^T = \mathbf{0}$, or $\mathbf{c}^T - \mathbf{y}^T A = \mathbf{v}^T$
and $v_i x_i = 0$ for all $i = 1, \dots, n$.
where $\mathbf{v} = [v_1, \dots, v_n]^T$ is a non negative vector,
and $\mathbf{y} = [y_1, \dots, y_m]^T$ is unrestricted in sign.
- The above conditions are equivalent to the **dual feasibility** and the **complementary slackness** conditions for the above LPP.

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$ are given by:

- $\mathbf{c}^T - \mathbf{y}^T A - \mathbf{v}^T = \mathbf{0}$, or $\mathbf{c}^T - \mathbf{y}^T A = \mathbf{v}^T$
and $v_i x_i = 0$ for all $i = 1, \dots, n$.
where $\mathbf{v} = [v_1, \dots, v_n]^T$ is a non negative vector,
and $\mathbf{y} = [y_1, \dots, y_m]^T$ is unrestricted in sign.
- The above conditions are equivalent to the **dual feasibility** and the **complementary slackness** conditions for the above LPP.
- **Conclusion:** $\mathbf{x}^* \in \text{Fea}(P)$ is **optimal** for (P) if and only if \mathbf{x}^* is a **KKT point** of (P).
- **Exercise:** What are the FJ points of the above problem?

- **Exercise:** The KKT conditions for the linear programming problem (LPP):

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A_{m \times n} \mathbf{x} = \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$ are given by:

- $\mathbf{c}^T - \mathbf{y}^T A - \mathbf{v}^T = \mathbf{0}$, or $\mathbf{c}^T - \mathbf{y}^T A = \mathbf{v}^T$
and $v_i x_i = 0$ for all $i = 1, \dots, n$.
where $\mathbf{v} = [v_1, \dots, v_n]^T$ is a non negative vector,
and $\mathbf{y} = [y_1, \dots, y_m]^T$ is unrestricted in sign.
- The above conditions are equivalent to the **dual feasibility** and the **complementary slackness** conditions for the above LPP.
- **Conclusion:** $\mathbf{x}^* \in \text{Fea}(P)$ is **optimal** for (P) if and only if \mathbf{x}^* is a **KKT point** of (P).
- **Exercise:** What are the FJ points of the above problem?
- **Exercise:** What are the KKT conditions for the following linear programming problem
$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to, $A\mathbf{x} \geq \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$.

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,

for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,

for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,

and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,

for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,

and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .

Then if \mathbf{x}^* is a **local minimum** of f over S

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,

for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,

and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .

Then if \mathbf{x}^* is a **local minimum** of f over S

there exists $u_0, u_i, i \in I$, **non negative** constants, and v_j constants (**unrestricted in sign**), not all zeros, such that

- **Theorem 10: (FJ necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

Consider the problem **P** of minimizing f subject to the conditions

$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

$h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,

for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,

and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .

Then if \mathbf{x}^* is a **local minimum** of f over S

there exists $u_0, u_i, i \in I$, **non negative** constants, and v_j constants (**unrestricted in sign**), not all zeros, such that

$$u_0 \nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^l v_j \nabla h_j(\mathbf{x}^*) = \mathbf{0}. \quad (*)$$

- **Example 5:** Minimize $x_1 + x_2$

- **Example 5:** Minimize $x_1 + x_2$
subject to
 $x_2 - x_1 = 0$.

- **Example 5:** Minimize $x_1 + x_2$
subject to
 $x_2 - x_1 = 0$.
- No point of the above problem satisfies the **FJ** conditions (as in Theorem 10).

- **Example 5:** Minimize $x_1 + x_2$
subject to
 $x_2 - x_1 = 0$.
- No point of the above problem satisfies the **FJ** conditions (as in Theorem 10).
- **Example 6:** Minimize $(x_1 - 1)^2 + x_2$

- **Example 5:** Minimize $x_1 + x_2$
subject to
 $x_2 - x_1 = 0$.
- No point of the above problem satisfies the **FJ** conditions (as in Theorem 10).
- **Example 6:** Minimize $(x_1 - 1)^2 + x_2$
subject to
 $x_2 - x_1 = 1$

- **Example 5:** Minimize $x_1 + x_2$
subject to
 $x_2 - x_1 = 0$.
- No point of the above problem satisfies the **FJ** conditions (as in Theorem 10).
- **Example 6:** Minimize $(x_1 - 1)^2 + x_2$
subject to
 $x_2 - x_1 = 1$
 $x_1 + x_2 \leq 2$.

- **Example 5:** Minimize $x_1 + x_2$
subject to
 $x_2 - x_1 = 0$.
- No point of the above problem satisfies the **FJ** conditions (as in Theorem 10).
- **Example 6:** Minimize $(x_1 - 1)^2 + x_2$
subject to
 $x_2 - x_1 = 1$
 $x_1 + x_2 \leq 2$.
- $\mathbf{x}^* = [\frac{1}{2}, \frac{3}{2}]^T$ is an **FJ** point with $u_0 = 1, v_1 = -1$ (as in Theorem 10).

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function.

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,
 $h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,
 $h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .
Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,
 $h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .
Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,
 $h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .
Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,
for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,
 $h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .
Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,
for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,
and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .

- **Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions
 $g_i(\mathbf{x}) \leq 0, i = 1, \dots, m$, where $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i ,
 $h_j(\mathbf{x}) = 0, j = 1, \dots, l$, where $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ for all j .
Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.
For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,
for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,
and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .
Let $\{\nabla g_i(\mathbf{x}^*), \nabla h_j(\mathbf{x}^*) \text{ for } i \in I, j \in \{1, \dots, l\}\}$ be **LI**. Then if \mathbf{x}^* is a **local minimum** of f over S

- Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions

$$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, \text{ where } g_i : \mathbb{R}^n \rightarrow \mathbb{R} \text{ for all } i,$$

$$h_j(\mathbf{x}) = 0, j = 1, \dots, l, \text{ where } h_j : \mathbb{R}^n \rightarrow \mathbb{R} \text{ for all } j.$$
 Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.
 For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,
 for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,
 and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .
 Let $\{\nabla g_i(\mathbf{x}^*), \nabla h_j(\mathbf{x}^*) \text{ for } i \in I, j \in \{1, \dots, l\}\}$ be **LI**. Then if \mathbf{x}^* is a **local minimum** of f over S
 there exists $u_i, i \in I$, **non negative** constants, and v_j constants (**unrestricted in sign**) such that

- Theorem 11: (KKT necessary conditions)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuously differentiable** function. Consider the problem **P** of minimizing f subject to the conditions

$$g_i(\mathbf{x}) \leq 0, i = 1, \dots, m, \text{ where } g_i : \mathbb{R}^n \rightarrow \mathbb{R} \text{ for all } i,$$

$$h_j(\mathbf{x}) = 0, j = 1, \dots, l, \text{ where } h_j : \mathbb{R}^n \rightarrow \mathbb{R} \text{ for all } j.$$
 Let $S = \text{Fea}(P)$ and $\mathbf{x}^* \in S$.
 For all $i \in I^*$, g_i 's are **continuous** at \mathbf{x}^* ,
 for all $i \in I$, g_i 's are **continuously differentiable** at \mathbf{x}^* ,
 and for all j , h_j 's are **continuously differentiable** at \mathbf{x}^* .
 Let $\{\nabla g_i(\mathbf{x}^*), \nabla h_j(\mathbf{x}^*) \text{ for } i \in I, j \in \{1, \dots, l\}\}$ be **LI**. Then if \mathbf{x}^* is a **local minimum** of f over S
 there exists $u_i, i \in I$, **non negative** constants, and v_j constants (**unrestricted in sign**) such that

$$\nabla f(\mathbf{x}^*) + \sum_{i \in I} u_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^l v_j \nabla h_j(\mathbf{x}^*) = \mathbf{0}. \quad (*)$$

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,
 $x_1 - x_2 = 0$.

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,
 $x_1 - x_2 = 0$.
- Check that at $\mathbf{x}^* = [0, 0]^T$, the first constraint is binding.

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,
 $x_1 - x_2 = 0$.
- Check that at $\mathbf{x}^* = [0, 0]^T$, the first constraint is binding.
- But $\nabla g_1(\mathbf{x}^*) = 2[-1, 1]^T$, and $\nabla h_1(\mathbf{x}^*) = [1, -1]^T$.

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,
 $x_1 - x_2 = 0$.
- Check that at $\mathbf{x}^* = [0, 0]^T$, the first constraint is binding.
- But $\nabla g_1(\mathbf{x}^*) = 2[-1, 1]^T$, and $\nabla h_1(\mathbf{x}^*) = [1, -1]^T$.
- Since $\{\nabla g_1(\mathbf{x}^*), \nabla h_1(\mathbf{x}^*)\}$ is **LD**,

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,
 $x_1 - x_2 = 0$.
- Check that at $\mathbf{x}^* = [0, 0]^T$, the first constraint is binding.
- But $\nabla g_1(\mathbf{x}^*) = 2[-1, 1]^T$, and $\nabla h_1(\mathbf{x}^*) = [1, -1]^T$.
- Since $\{\nabla g_1(\mathbf{x}^*), \nabla h_1(\mathbf{x}^*)\}$ is **LD**, Theorem 11 is not applicable.

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,
 $x_1 - x_2 = 0$.
- Check that at $\mathbf{x}^* = [0, 0]^T$, the first constraint is binding.
- But $\nabla g_1(\mathbf{x}^*) = 2[-1, 1]^T$, and $\nabla h_1(\mathbf{x}^*) = [1, -1]^T$.
- Since $\{\nabla g_1(\mathbf{x}^*), \nabla h_1(\mathbf{x}^*)\}$ is **LD**, Theorem 11 is not applicable.
- Check that $[0, 0]^T$ satisfies the KKT conditions as given in **(3)**, immediately after Theorem 7.
- Since f, g_1, g_2, g_3 is convex, where $g_2(x_1, x_2) = x_1 - x_2$ and $g_3(x_1, x_2) = -x_1 + x_2$,

- **Example 7:** Minimize $f(x_1, x_2) = x_1^3 - x_1 + x_2$,
subject to
 $(x_1 - 1)^2 + 2x_2 \leq 1$,
 $x_1 - x_2 = 0$.
- Check that at $\mathbf{x}^* = [0, 0]^T$, the first constraint is binding.
- But $\nabla g_1(\mathbf{x}^*) = 2[-1, 1]^T$, and $\nabla h_1(\mathbf{x}^*) = [1, -1]^T$.
- Since $\{\nabla g_1(\mathbf{x}^*), \nabla h_1(\mathbf{x}^*)\}$ is **LD**, Theorem 11 is not applicable.
- Check that $[0, 0]^T$ satisfies the KKT conditions as given in **(3)**, immediately after Theorem 7.
- Since f, g_1, g_2, g_3 is convex, where $g_2(x_1, x_2) = x_1 - x_2$ and $g_3(x_1, x_2) = -x_1 + x_2$,
- $[0, 0]^T$ is the global minimum of f in the given feasible region.

- Consider the problem (P) of minimizing $f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$ over R^2 .

- Consider the problem (P) of minimizing $f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$ over R^2 .
 - 1 Write the first order necessary optimality condition for this problem. Is this condition also sufficient for optimality for (P)?

- Consider the problem (P) of minimizing
$$f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$$
over R^2 .
 - 1 Write the first order necessary optimality condition for this problem. Is this condition also sufficient for optimality for (P)?
 - 2 Is $[0, 0]^T$ a local minimum point for this problem? If not find $\mathbf{d} \in D_{\mathbf{x}^*}$ along which the function f will decrease.

- Consider the problem (P) of minimizing
 $f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$ over R^2 .
 - Write the first order necessary optimality condition for this problem. Is this condition also sufficient for optimality for (P)?
 - Is $[0, 0]^T$ a local minimum point for this problem? If not find $\mathbf{d} \in D_{\mathbf{x}^*}$ along which the function f will decrease.
- Minimize $(x_1 - 2)^2 + (x_2 - 3)^2$
 subject to
 $x_1^2 + x_2^2 \leq 5$.
 $2x_1 + x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.

- Consider the problem (P) of minimizing $f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$ over R^2 .
 - Write the first order necessary optimality condition for this problem. Is this condition also sufficient for optimality for (P)?
 - Is $[0, 0]^T$ a local minimum point for this problem? If not find $\mathbf{d} \in D_{\mathbf{x}^*}$ along which the function f will decrease.
- Minimize $(x_1 - 2)^2 + (x_2 - 3)^2$
 subject to
 $x_1^2 + x_2^2 \leq 5$.
 $2x_1 + x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
 - $\mathbf{x}^* = [1, 2]^T$ is a KKT point.

- Consider the problem (P) of minimizing $f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$ over R^2 .
 - Write the first order necessary optimality condition for this problem. Is this condition also sufficient for optimality for (P)?
 - Is $[0, 0]^T$ a local minimum point for this problem? If not find $\mathbf{d} \in D_{\mathbf{x}^*}$ along which the function f will decrease.
- Minimize $(x_1 - 2)^2 + (x_2 - 3)^2$
 subject to
 $x_1^2 + x_2^2 \leq 5$.
 $2x_1 + x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
 - $\mathbf{x}^* = [1, 2]^T$ is a KKT point.
 - \mathbf{x}^* a global minimum of f over this feasible region.

- Consider the problem (P) of minimizing $f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$ over R^2 .
 - Write the first order necessary optimality condition for this problem. Is this condition also sufficient for optimality for (P)?
 - Is $[0, 0]^T$ a local minimum point for this problem? If not find $\mathbf{d} \in D_{\mathbf{x}^*}$ along which the function f will decrease.
- Minimize $(x_1 - 2)^2 + (x_2 - 3)^2$
 subject to
 $x_1^2 + x_2^2 \leq 5$.
 $2x_1 + x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
 - $\mathbf{x}^* = [1, 2]^T$ is a KKT point.
 - \mathbf{x}^* a global minimum of f over this feasible region.
 - Does there exist an $\mathbf{x}^* \in \text{Fea}(P)$ such that $G_{0, \mathbf{x}^*} = D_{\mathbf{x}^*}$?

- Consider the problem (P) of minimizing $f(x_1, x_2) = -x_1 x_2 + x_1^2 + 2x_2^2 - 2x_1 + e^{x_1+x_2}$ over R^2 .
 - Write the first order necessary optimality condition for this problem. Is this condition also sufficient for optimality for (P)?
 - Is $[0, 0]^T$ a local minimum point for this problem? If not find $\mathbf{d} \in D_{\mathbf{x}^*}$ along which the function f will decrease.
- Minimize $(x_1 - 2)^2 + (x_2 - 3)^2$
 subject to
 $x_1^2 + x_2^2 \leq 5$.
 $2x_1 + x_2 \leq 4$.
 $-x_1 \leq 0$
 $-x_2 \leq 0$.
 - $\mathbf{x}^* = [1, 2]^T$ is a KKT point.
 - \mathbf{x}^* a global minimum of f over this feasible region.
 - Does there exist an $\mathbf{x}^* \in \text{Fea}(P)$ such that $G_{0, \mathbf{x}^*} = D_{\mathbf{x}^*}$?
 - Does there exist an $\mathbf{x}^* \in \text{Fea}(P)$ such that $G_{0, \mathbf{x}^*} \neq D_{\mathbf{x}^*}$?
 - Does there exist an FJ point which is not a KKT point?

- Consider the following problem:

$$\text{Minimize } 4x_1^2 - x_2^2 + 8x_1x_2$$

subject to

$$2x_1 + x_1^2 - x_2 \geq 0$$

$$x_1 \geq 0, x_2 \geq 0.$$

- Consider the following problem:

$$\text{Minimize } 4x_1^2 - x_2^2 + 8x_1x_2$$

subject to

$$2x_1 + x_1^2 - x_2 \geq 0$$

$$x_1 \geq 0, x_2 \geq 0.$$

- Does $[0, 0]^T$ satisfy the first and the second order necessary conditions for a local minimum?

- Consider the following problem:

$$\text{Minimize } 4x_1^2 - x_2^2 + 8x_1x_2$$

subject to

$$2x_1 + x_1^2 - x_2 \geq 0$$

$$x_1 \geq 0, x_2 \geq 0.$$

- Does $[0, 0]^T$ satisfy the first and the second order necessary conditions for a local minimum?
- Find all points \mathbf{x}^* in the feasible region at which $G_{0,\mathbf{x}^*} = \phi$.

- Consider the following problem:

$$\text{Minimize } 4x_1^2 - x_2^2 + 8x_1x_2$$

subject to

$$2x_1 + x_1^2 - x_2 \geq 0$$

$$x_1 \geq 0, x_2 \geq 0.$$

- Does $[0, 0]^T$ satisfy the first and the second order necessary conditions for a local minimum?
 - Find all points \mathbf{x}^* in the feasible region at which $G_{0, \mathbf{x}^*} = \phi$.
- If $\mathbf{x}^* \in S$ is such that for all $\mathbf{d} \in D_{\mathbf{x}^*} (\neq \phi)$, $\nabla f(\mathbf{x}^*)\mathbf{d} > 0$ then does it imply that \mathbf{x}^* is a local minimizer of f ?

- Consider the following problem:

$$\text{Minimize } -x_1^2 - 4x_1x_2 - x_2^2$$

subject to

$$x_2^2 + x_1^2 = 1.$$

- 1 If possible find an FJ point which is not a KKT point.

- Consider the following problem:

Minimize $-x_1^2 - 4x_1x_2 - x_2^2$

subject to

$$x_2^2 + x_1^2 = 1.$$

- 1 If possible find an FJ point which is not a KKT point.
- 2 If possible find a KKT point which is not an optimal point.

- Consider the following problem:

$$\text{Minimize } -x_1^2 - 4x_1x_2 - x_2^2$$

subject to

$$x_2^2 + x_1^2 = 1.$$

- 1 If possible find an FJ point which is not a KKT point.
- 2 If possible find a KKT point which is not an optimal point.
- 3 Are the first and the second order necessary conditions for a local minimum satisfied at the KKT point/s?

- Consider the following problem:

Minimize $-x_1^2 - 4x_1x_2 - x_2^2$

subject to

$$x_2^2 + x_1^2 = 1.$$

- 1 If possible find an FJ point which is not a KKT point.
- 2 If possible find a KKT point which is not an optimal point.
- 3 Are the first and the second order necessary conditions for a local minimum satisfied at the KKT point/s?
- 4 If the objective function is changed to $2x_1^2 - x_1x_2 + x_2^2 - x_2$ and if $S = \text{Fea}(P) = \{(x_1, x_2) : x_2^2 + x_1^2 \leq 1\}$ then find all optimum solutions to this problem.

- For a nonlinear programming problem (P) of the form,

Minimize $f(\mathbf{x})$

subject to $g_i(\mathbf{x}) \leq 0$, for $i = 1, \dots, m$, $\mathbf{x} \in \mathbb{R}^n$,

where all the g_i 's and f are continuously differentiable throughout \mathbb{R}^n , check the correctness of the following statements with proper justification.

- 1 If $\mathbf{x}^* \in \text{Fea}(P)$ is a KKT point of the above problem then $-\nabla f(\mathbf{x}^*)$ lies in the cone generated by $\nabla g_i(\mathbf{x}^*)$, $i \in I$, where I gives the indices of the binding constraints (given by g_i 's) at \mathbf{x}^* .

- For a nonlinear programming problem (P) of the form,

Minimize $f(\mathbf{x})$

subject to $g_i(\mathbf{x}) \leq 0$, for $i = 1, \dots, m$, $\mathbf{x} \in \mathbb{R}^n$,

where all the g_i 's and f are continuously differentiable throughout \mathbb{R}^n , check the correctness of the following statements with proper justification.

- 1 If $\mathbf{x}^* \in \text{Fea}(P)$ is a KKT point of the above problem then $-\nabla f(\mathbf{x}^*)$ lies in the cone generated by $\nabla g_i(\mathbf{x}^*)$, $i \in I$, where I gives the indices of the binding constraints (given by g_i 's) at \mathbf{x}^* .
- 2 If $g_2 = -g_3$ then all points $\mathbf{x} \in \text{Fea}(P)$ are FJ (Fritz John) points.

- For a nonlinear programming problem (P) of the form,

Minimize $f(\mathbf{x})$

subject to $g_i(\mathbf{x}) \leq 0$, for $i = 1, \dots, m$, $\mathbf{x} \in \mathbb{R}^n$,

where all the g_i 's and f are continuously differentiable throughout \mathbb{R}^n , check the correctness of the following statements with proper justification.

- 1 If $\mathbf{x}^* \in \text{Fea}(P)$ is a KKT point of the above problem then $-\nabla f(\mathbf{x}^*)$ lies in the cone generated by $\nabla g_i(\mathbf{x}^*)$, $i \in I$, where I gives the indices of the binding constraints (given by g_i 's) at \mathbf{x}^* .
- 2 If $g_2 = -g_3$ then all points $\mathbf{x} \in \text{Fea}(P)$ are FJ (Fritz John) points.
- 3 If $\nabla f(\mathbf{x}^*) = 0$ for some $\mathbf{x}^* \in \text{Fea}(P)$ then \mathbf{x}^* is a KKT point.

- For a nonlinear programming problem (P) of the form,

Minimize $f(\mathbf{x})$

subject to $g_i(\mathbf{x}) \leq 0$, for $i = 1, \dots, m$, $\mathbf{x} \in \mathbb{R}^n$,

where all the g_i 's and f are continuously differentiable throughout \mathbb{R}^n , check the correctness of the following statements with proper justification.

- 1 If $\mathbf{x}^* \in \text{Fea}(P)$ is a KKT point of the above problem then $-\nabla f(\mathbf{x}^*)$ lies in the cone generated by $\nabla g_i(\mathbf{x}^*)$, $i \in I$, where I gives the indices of the binding constraints (given by g_i 's) at \mathbf{x}^* .
- 2 If $g_2 = -g_3$ then all points $\mathbf{x} \in \text{Fea}(P)$ are FJ (Fritz John) points.
- 3 If $\nabla f(\mathbf{x}^*) = 0$ for some $\mathbf{x}^* \in \text{Fea}(P)$ then \mathbf{x}^* is a KKT point.
- 4 If \mathbf{x}^* is an FJ point and there is a solution to the FJ conditions at \mathbf{x}^* with $u_0 = 0$ (u_0 is the coefficient of $\nabla f(\mathbf{x}^*)$ in the FJ conditions), then \mathbf{x}^* is not a KKT point.

- If \mathbf{x}^* is a KKT point then it is also an FJ point.

- If \mathbf{x}^* is a KKT point then it is also an FJ point.
- If \mathbf{x}^* is an FJ point and $\nabla g_i(\mathbf{x}^*)$'s are LD, then $G_{0,\mathbf{x}^*} = \phi$ (or in other words there exists a solution to the FJ conditions with $u_0 = 0$).

- If \mathbf{x}^* is a KKT point then it is also an FJ point.
- If \mathbf{x}^* is an FJ point and $\nabla g_i(\mathbf{x}^*)$'s are LD, then $G_{0,\mathbf{x}^*} = \phi$ (or in other words there exists a solution to the FJ conditions with $u_0 = 0$).
- If $\nabla g_i(\mathbf{x}^*)$'s are LI, then $G_{0,\mathbf{x}^*} \neq \phi$.

- If \mathbf{x}^* is a KKT point then it is also an FJ point.
- If \mathbf{x}^* is an FJ point and $\nabla g_i(\mathbf{x}^*)$'s are LD, then $G_{0,\mathbf{x}^*} = \phi$ (or in other words there exists a solution to the FJ conditions with $u_0 = 0$).
- If $\nabla g_i(\mathbf{x}^*)$'s are LI, then $G_{0,\mathbf{x}^*} \neq \phi$.
- If $G_{0,\mathbf{x}^*} = \phi$ then $\nabla g_i(\mathbf{x}^*)$'s are LD.

- If \mathbf{x}^* is a KKT point then it is also an FJ point.
- If \mathbf{x}^* is an FJ point and $\nabla g_i(\mathbf{x}^*)$'s are LD, then $G_{0,\mathbf{x}^*} = \phi$ (or in other words there exists a solution to the FJ conditions with $u_0 = 0$).
- If $\nabla g_i(\mathbf{x}^*)$'s are LI, then $G_{0,\mathbf{x}^*} \neq \phi$.
- If $G_{0,\mathbf{x}^*} = \phi$ then $\nabla g_i(\mathbf{x}^*)$'s are LD.
- If \mathbf{x}^* is not an interior point of the feasible region S of (P) and $F_{0,\mathbf{x}^*} = \phi$ then \mathbf{x}^* is a local minimizer.

- If \mathbf{x}^* is a KKT point then it is also an FJ point.
- If \mathbf{x}^* is an FJ point and $\nabla g_i(\mathbf{x}^*)$'s are LD, then $G_{0,\mathbf{x}^*} = \phi$ (or in other words there exists a solution to the FJ conditions with $u_0 = 0$).
- If $\nabla g_i(\mathbf{x}^*)$'s are LI, then $G_{0,\mathbf{x}^*} \neq \phi$.
- If $G_{0,\mathbf{x}^*} = \phi$ then $\nabla g_i(\mathbf{x}^*)$'s are LD.
- If \mathbf{x}^* is not an interior point of the feasible region S of (P) and $F_{0,\mathbf{x}^*} = \phi$ then \mathbf{x}^* is a local minimizer.
- If all the g_i 's and f are convex functions and \mathbf{x}^* is such that $F_{0,\mathbf{x}^*} \cap G_{0,\mathbf{x}^*} = \phi$, then \mathbf{x}^* is a global minimum of f in S .

- If \mathbf{x}^* is a KKT point then it is also an FJ point.
- If \mathbf{x}^* is an FJ point and $\nabla g_i(\mathbf{x}^*)$'s are LD, then $G_{0,\mathbf{x}^*} = \phi$ (or in other words there exists a solution to the FJ conditions with $u_0 = 0$).
- If $\nabla g_i(\mathbf{x}^*)$'s are LI, then $G_{0,\mathbf{x}^*} \neq \phi$.
- If $G_{0,\mathbf{x}^*} = \phi$ then $\nabla g_i(\mathbf{x}^*)$'s are LD.
- If \mathbf{x}^* is not an interior point of the feasible region S of (P) and $F_{0,\mathbf{x}^*} = \phi$ then \mathbf{x}^* is a local minimizer.
- If all the g_i 's and f are convex functions and \mathbf{x}^* is such that $F_{0,\mathbf{x}^*} \cap G_{0,\mathbf{x}^*} = \phi$, then \mathbf{x}^* is a global minimum of f in S .
- If \mathbf{x}^* is a local minima of (P) with $F_{0,\mathbf{x}^*} \neq \phi$ but $G_{0,\mathbf{x}^*} = \phi$ then \mathbf{x}^* is not a KKT point.

- There exists a (P) such that for all $\mathbf{x}^* \in S$, $G_{0,\mathbf{x}^*} = D_{\mathbf{x}^*} \neq \phi$.

- There exists a (P) such that for all $\mathbf{x}^* \in S$, $G_{0,\mathbf{x}^*} = D_{\mathbf{x}^*} \neq \phi$.
- There exists a (P) such that $G_{0,\mathbf{x}^*} \neq D_{\mathbf{x}^*}$, for all $\mathbf{x}^* \in S$.

- There exists a (P) such that for all $\mathbf{x}^* \in S$, $G_{0,\mathbf{x}^*} = D_{\mathbf{x}^*} \neq \phi$.
- There exists a (P) such that $G_{0,\mathbf{x}^*} \neq D_{\mathbf{x}^*}$, for all $\mathbf{x}^* \in S$.
- There exists a (P) with $Fea(P) \neq \phi$ such that $G_{0,\mathbf{x}^*} = \phi$ for all $\mathbf{x}^* \in S$.

- There exists a (P) such that for all $\mathbf{x}^* \in S$, $G_{0,\mathbf{x}^*} = D_{\mathbf{x}^*} \neq \phi$.
- There exists a (P) such that $G_{0,\mathbf{x}^*} \neq D_{\mathbf{x}^*}$, for all $\mathbf{x}^* \in S$.
- There exists a (P) with $Fea(P) \neq \phi$ such that $G_{0,\mathbf{x}^*} = \phi$ for all $\mathbf{x}^* \in S$.
- There exists a (P) with $Fea(P) \neq \phi$ such that $D_{\mathbf{x}^*} = \phi$ for all $\mathbf{x}^* \in S$.

- There exists a (P) such that for all $\mathbf{x}^* \in S$, $G_{0,\mathbf{x}^*} = D_{\mathbf{x}^*} \neq \phi$.
- There exists a (P) such that $G_{0,\mathbf{x}^*} \neq D_{\mathbf{x}^*}$, for all $\mathbf{x}^* \in S$.
- There exists a (P) with $Fea(P) \neq \phi$ such that $G_{0,\mathbf{x}^*} = \phi$ for all $\mathbf{x}^* \in S$.
- There exists a (P) with $Fea(P) \neq \phi$ such that $D_{\mathbf{x}^*} = \phi$ for all $\mathbf{x}^* \in S$.
- Give examples of nonconstant functions on \mathbb{R}^n which are both convex and concave and those which are neither convex nor concave.

- For a linear programming problem (P) of the form,
Minimize $\mathbf{c}^T \mathbf{x}$ subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$,
find the KKT conditions.

- For a linear programming problem (P) of the form,
Minimize $\mathbf{c}^T \mathbf{x}$ subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$,
find the KKT conditions.
- Consider the linear programming problem.
Minimize $2x_1 - 3x_2$
subject to $x_1 + 2x_2 \leq 3$
 $2x_1 + 3x_2 \leq 5$
 $x_1 \geq 0, x_2 \geq 0$.

- For a linear programming problem (P) of the form,
Minimize $\mathbf{c}^T \mathbf{x}$ subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$,
find the KKT conditions.
- Consider the linear programming problem.
Minimize $2x_1 - 3x_2$
subject to $x_1 + 2x_2 \leq 3$
 $2x_1 + 3x_2 \leq 5$
 $x_1 \geq 0, x_2 \geq 0$. Find the KKT conditions for this problem at
a local minimum point of this problem.
Solve the KKT conditions for u_i 's. Hence find an optimal
solution of the dual.

- For a linear programming problem (P) of the form,
Minimize $\mathbf{c}^T \mathbf{x}$ subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$,
find the KKT conditions.
- Consider the linear programming problem.
Minimize $2x_1 - 3x_2$
subject to $x_1 + 2x_2 \leq 3$
 $2x_1 + 3x_2 \leq 5$
 $x_1 \geq 0, x_2 \geq 0$. Find the KKT conditions for this problem at
a local minimum point of this problem.
Solve the KKT conditions for u_i 's. Hence find an optimal
solution of the dual.
- Minimize $-x_1$
subject to
 $-(1 - x_1)^3 + x_2 = 0$
 $-(1 - x_1)^3 - x_2 = 0$.

- For a linear programming problem (P) of the form,
Minimize $\mathbf{c}^T \mathbf{x}$ subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$,
find the KKT conditions.
- Consider the linear programming problem.
Minimize $2x_1 - 3x_2$
subject to $x_1 + 2x_2 \leq 3$
 $2x_1 + 3x_2 \leq 5$
 $x_1 \geq 0, x_2 \geq 0$. Find the KKT conditions for this problem at a local minimum point of this problem.
Solve the KKT conditions for u_i 's. Hence find an optimal solution of the dual.
- Minimize $-x_1$
subject to
 $-(1 - x_1)^3 + x_2 = 0$
 $-(1 - x_1)^3 - x_2 = 0$.
- Check whether $[1, 0]^T$ is a KKT point of the above problem. How many feasible points does this problem have? What is your conclusion?