

Plan

- Directions of Fea(LPP)
- Extreme directions of Fea(LPP)
- Representation Theorem for Fea(LPP)
- Necessary and sufficient conditions for the existence of optimal solutions
- Optimal solutions in atleast one corner point

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector $\mathbf{d} \neq \mathbf{0}$ such that starting from any $\mathbf{x} \in Fea(LPP)$ and moving in the positive direction of \mathbf{d} , always gives elements of $Fea(LPP)$.

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$
- **Observation 5:** Suppose if a LPP has an unbounded feasible region, then there exists a vector $\mathbf{d} \neq \mathbf{0}$ such that starting from any $\mathbf{x} \in Fea(LPP)$ and moving in the positive direction of \mathbf{d} , always gives elements of $Fea(LPP)$.

That is for any $\mathbf{x} \in Fea(LPP)$, $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$ for all $\alpha \geq 0$,

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector $\mathbf{d} \neq \mathbf{0}$ such that starting from any $\mathbf{x} \in Fea(LPP)$ and moving in the positive direction of \mathbf{d} , always gives elements of $Fea(LPP)$.
That is for any $\mathbf{x} \in Fea(LPP)$, $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$ for all $\alpha \geq 0$,
then $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of $S = Fea(LPP)$.

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$.
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector $\mathbf{d} \neq \mathbf{0}$ such that starting from any $\mathbf{x} \in Fea(LPP)$ and moving in the positive direction of \mathbf{d} , always gives elements of $Fea(LPP)$.

That is for any $\mathbf{x} \in Fea(LPP)$, $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$ for all $\alpha \geq 0$,

then $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of $S = Fea(LPP)$.

Throughout our discussion, $\mathbf{d} = [d_1, \dots, d_n]^T$ will denote a column vector.

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$.
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector $\mathbf{d} \neq \mathbf{0}$ such that starting from any $\mathbf{x} \in Fea(LPP)$ and moving in the positive direction of \mathbf{d} , always gives elements of $Fea(LPP)$.

That is for any $\mathbf{x} \in Fea(LPP)$, $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$ for all $\alpha \geq 0$,

then $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of $S = Fea(LPP)$.

Throughout our discussion, $\mathbf{d} = [d_1, \dots, d_n]^T$ will denote a column vector.

- **Definition:** Given a non empty **convex set** S , $S \subset \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of S if for all $\mathbf{x} \in S$, $\mathbf{x} + \alpha \mathbf{d} \in S$ for all $\alpha \geq 0$.

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$.
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector $\mathbf{d} \neq \mathbf{0}$ such that starting from any $\mathbf{x} \in Fea(LPP)$ and moving in the positive direction of \mathbf{d} , always gives elements of $Fea(LPP)$.

That is for any $\mathbf{x} \in Fea(LPP)$, $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$ for all $\alpha \geq 0$,

then $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of $S = Fea(LPP)$.

Throughout our discussion, $\mathbf{d} = [d_1, \dots, d_n]^T$ will denote a column vector.

- **Definition:** Given a non empty **convex set** S , $S \subset \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of S if for all $\mathbf{x} \in S$, $\mathbf{x} + \alpha \mathbf{d} \in S$ for all $\alpha \geq 0$.
- If \mathbf{d} is a **direction** of a convex set S , then for all $\gamma > 0$, $\mathbf{x} + \alpha \mathbf{d} = \mathbf{x} + \left(\frac{\alpha}{\gamma}\right) \gamma \mathbf{d} \in S$ for all $\alpha > 0$,

- $Fea(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$.
- **Observation 5:** Suppose if a LPP has an **unbounded feasible region**, then there exists a vector $\mathbf{d} \neq \mathbf{0}$ such that starting from any $\mathbf{x} \in Fea(LPP)$ and moving in the positive direction of \mathbf{d} , always gives elements of $Fea(LPP)$.

That is for any $\mathbf{x} \in Fea(LPP)$, $\mathbf{x} + \alpha \mathbf{d} \in Fea(LPP)$ for all $\alpha \geq 0$,

then $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of $S = Fea(LPP)$.

Throughout our discussion, $\mathbf{d} = [d_1, \dots, d_n]^T$ will denote a column vector.

- **Definition:** Given a non empty **convex set** S , $S \subset \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is called a **direction** of S if for all $\mathbf{x} \in S$, $\mathbf{x} + \alpha \mathbf{d} \in S$ for all $\alpha \geq 0$.
- If \mathbf{d} is a **direction** of a convex set S , then for all $\gamma > 0$, $\mathbf{x} + \alpha \mathbf{d} = \mathbf{x} + \left(\frac{\alpha}{\gamma}\right) \gamma \mathbf{d} \in S$ for all $\alpha > 0$,
 $\Rightarrow \gamma \mathbf{d}$ is again a **direction** for all $\gamma > 0$.

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,
 $\text{Min } -x + 2y$

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,
Min $-x + 2y$
subject to
 $x + 2y \geq 1$

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,
Min $-x + 2y$
subject to
 $x + 2y \geq 1$
 $-x + y \leq 1,$

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$
 subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,
Min $-x + 2y$
subject to
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,
Min $-x + 2y$
subject to
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$
are all equal as directions of $Fea(LPP)$.
- Similarly $\mathbf{d}_2 = [1, 0]^T, \mathbf{d}'_2 = [2, 0]^T, \dots$

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,
Min $-x + 2y$
subject to
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$
are all equal as directions of $Fea(LPP)$.
- Similarly $\mathbf{d}_2 = [1, 0]^T, \mathbf{d}'_2 = [2, 0]^T, \dots$
are all equal as directions of $Fea(LPP)$.

- Directions $\mathbf{d}_1, \mathbf{d}_2$ of S are said to be **distinct** if $\mathbf{d}_1 \neq \gamma \mathbf{d}_2$ for any $\gamma > 0$
(or equivalently $\mathbf{d}_2 \neq \beta \mathbf{d}_1$ for any $\beta > 0$).
- **Example 2: (revisited)** Consider the problem,
Min $-x + 2y$
subject to
 $x + 2y \geq 1$
 $-x + y \leq 1$,
 $x \geq 0, y \geq 0$.
- $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}'_1 = [2, 2]^T, \dots$
are all equal as directions of $Fea(LPP)$.
- Similarly $\mathbf{d}_2 = [1, 0]^T, \mathbf{d}'_2 = [2, 0]^T, \dots$
are all equal as directions of $Fea(LPP)$.
- Whereas $\mathbf{d}_1 = [1, 1]^T, \mathbf{d}_2 = [1, 0]^T$ give two **distinct directions**.

- **Result:** The set of all directions of
 $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$

- **Result:** The set of all directions of $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is given by

- **Result:** The set of all directions of
 $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is given by
 $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \quad A_{m \times n} \mathbf{d} \leq \mathbf{0}, \quad \mathbf{d} \geq \mathbf{0}\}$ or by

- **Result:** The set of all directions of

$S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is given by

$D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \quad A_{m \times n} \mathbf{d} \leq \mathbf{0}, \quad \mathbf{d} \geq \mathbf{0}\}$ or by

$D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \quad \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \quad \mathbf{d} \geq \mathbf{0}\}.$

- Result:** The set of all directions of $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is given by $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} \leq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ or by $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \mathbf{d} \geq \mathbf{0}\}$.
- If $\mathbf{d}_1, \mathbf{d}_2$ are directions of S , then $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ will again be a direction of S , for any α, β non negative

- Result:** The set of all directions of $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is given by $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} \leq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ or by $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \mathbf{d} \geq \mathbf{0}\}$.
- If $\mathbf{d}_1, \mathbf{d}_2$ are directions of S , then $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ will again be a direction of S , for any α, β non negative (as long as both α, β are not equal to zero, or $\alpha + \beta \neq 0$).

- Result:** The set of all directions of $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is given by $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, A_{m \times n} \mathbf{d} \leq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$ or by $D = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \neq \mathbf{0}, \mathbf{a}_i^T \mathbf{d} \leq 0, \text{ for all } i = 1, 2, \dots, m, \mathbf{d} \geq \mathbf{0}\}$.
- If $\mathbf{d}_1, \mathbf{d}_2$ are directions of S , then $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$ will again be a direction of S , for any α, β non negative (as long as both α, β are not equal to zero, or $\alpha + \beta \neq 0$).
- The set of all directions of $S = \text{Fea}(LPP)$ is a **convex set**.

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S ,

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of S ,

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of S , that is, if \mathbf{d} an **extreme direction** of S and

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of S , that is, if \mathbf{d} an **extreme direction** of S and $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$, for $\alpha, \beta > 0$ and $\mathbf{d}_1, \mathbf{d}_2 \in D$ then $\mathbf{d}_1 = \gamma \mathbf{d}_2$ for some $\gamma > 0$.

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of S , that is, if \mathbf{d} an **extreme direction** of S and $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$, for $\alpha, \beta > 0$ and $\mathbf{d}_1, \mathbf{d}_2 \in D$ then $\mathbf{d}_1 = \gamma \mathbf{d}_2$ for some $\gamma > 0$.
- If D denotes the set of all directions of S ($D = \phi$ if S is bounded), then

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of S , that is, if \mathbf{d} an **extreme direction** of S and $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$, for $\alpha, \beta > 0$ and $\mathbf{d}_1, \mathbf{d}_2 \in D$ then $\mathbf{d}_1 = \gamma \mathbf{d}_2$ for some $\gamma > 0$.
- If D denotes the set of all directions of S ($D = \emptyset$ if S is bounded), then $D' = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, A\mathbf{d} \leq \mathbf{0}, \sum_i d_i = 1\}$ is a set of all **distinct directions** of S .

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of S , that is, if \mathbf{d} an **extreme direction** of S and $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$, for $\alpha, \beta > 0$ and $\mathbf{d}_1, \mathbf{d}_2 \in D$ then $\mathbf{d}_1 = \gamma \mathbf{d}_2$ for some $\gamma > 0$.
- If D denotes the set of all directions of S ($D = \emptyset$ if S is bounded), then $D' = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, A\mathbf{d} \leq \mathbf{0}, \sum_i d_i = 1\}$ is a set of all **distinct directions** of S .
- Also each $\mathbf{d} \in D$ is of the form $\mathbf{d} = \alpha \mathbf{d}'$ for some $\mathbf{d}' \in D'$ where $\alpha = \sum_i d_i (> 0)$.
- $D' = \left\{ \mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, \begin{bmatrix} & & A \\ 1 & 1, \dots, & 1 \\ -1 & -1, \dots, & -1 \end{bmatrix} \mathbf{d} \leq \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right\}.$

- **Definition:** A direction \mathbf{d} of S is called an **extreme direction** of S , if it **cannot** be written as a **positive linear combination** of **two distinct directions** of S , that is, if \mathbf{d} an **extreme direction** of S and $\mathbf{d} = \alpha \mathbf{d}_1 + \beta \mathbf{d}_2$, for $\alpha, \beta > 0$ and $\mathbf{d}_1, \mathbf{d}_2 \in D$ then $\mathbf{d}_1 = \gamma \mathbf{d}_2$ for some $\gamma > 0$.

- If D denotes the set of all directions of S ($D = \emptyset$ if S is bounded), then

$D' = \{\mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, A\mathbf{d} \leq \mathbf{0}, \sum_i d_i = 1\}$ is a set of all **distinct directions** of S .

- Also each $\mathbf{d} \in D$ is of the form $\mathbf{d} = \alpha \mathbf{d}'$ for some $\mathbf{d}' \in D'$ where $\alpha = \sum_i d_i (> 0)$.

- $D' = \left\{ \mathbf{d} \in \mathbb{R}^n : \mathbf{d} \geq \mathbf{0}, \begin{bmatrix} & & A & \\ 1 & 1, \dots, & 1 \\ -1 & -1, \dots, & -1 \end{bmatrix} \mathbf{d} \leq \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right\}.$

The set D' now looks exactly like the feasible region of a LPP.

- Since D' is like $Fea(LPP) = S$, if $D' \neq \phi$, D' must have at least one **extreme point**.

- Since D' is like $Fea(LPP) = S$, if $D' \neq \phi$, D' must have at least one extreme point.
- **Result:** \mathbf{d} is an extreme direction of S if and only if $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$ is an extreme point of D' .

- Since D' is like $Fea(LPP) = S$, if $D' \neq \phi$, D' must have at least one **extreme point**.
- **Result:** \mathbf{d} is an **extreme direction** of S if and only if $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$ is an **extreme point** of D' .
- If $D \neq \phi$, then $Fea(LPP) = S(\neq \phi)$ must have at least one **extreme direction**.

- Since D' is like $Fea(LPP) = S$, if $D' \neq \phi$, D' must have at least one **extreme point**.
- **Result:** \mathbf{d} is an **extreme direction** of S if and only if $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$ is an **extreme point** of D' .
- If $D \neq \phi$, then $Fea(LPP) = S(\neq \phi)$ must have at least one **extreme direction**.
- If $Fea(LPP) = S \neq \phi$ is **unbounded** then $D \neq \phi$ and S must have at least one **extreme direction**.

- Since D' is like $Fea(LPP) = S$, if $D' \neq \phi$, D' must have at least one **extreme point**.
- **Result:** \mathbf{d} is an **extreme direction** of S if and only if $\mathbf{d}' = \frac{\mathbf{d}}{\sum_i d_i}$ is an **extreme point** of D' .
- If $D \neq \phi$, then $Fea(LPP) = S(\neq \phi)$ must have at least one **extreme direction**.
- If $Fea(LPP) = S \neq \phi$ is **unbounded** then $D \neq \phi$ and S must have at least one **extreme direction**.
- The number of **distinct extreme directions** of S is **finite** (why?).

- The **extreme directions of S** which are **extreme points of D'** (after suitable normalization) will lie on **n LI hyperplanes** defining D' .

- The **extreme directions of S** which are **extreme points** of D' (after suitable normalization) will lie on **n LI hyperplanes** defining D' .
- Since $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ cannot be orthogonal to n **LI** vectors,

- The **extreme directions of S** which are **extreme points** of D' (after suitable normalization) will lie on **n LI hyperplanes** defining D' .
- Since $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ cannot be orthogonal to n **LI** vectors, so \mathbf{d} cannot lie on **n LI hyperplanes** of the $(m + n)$ hyperplanes given by,

- The **extreme directions of S** which are **extreme points** of D' (after suitable normalization) will lie on **n LI hyperplanes** defining D' .
- Since $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ cannot be orthogonal to n **LI** vectors, so \mathbf{d} cannot lie on **n LI hyperplanes** of the $(m + n)$ hyperplanes given by,
 $\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\}$ for $i = 1, 2, \dots, m$, and

- The **extreme directions of S** which are **extreme points** of D' (after suitable normalization) will lie on **n LI hyperplanes** defining D' .
- Since $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ cannot be orthogonal to n **LI** vectors, so \mathbf{d} cannot lie on **n LI hyperplanes** of the $(m + n)$ hyperplanes given by,

$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$

- The **extreme directions of S** which are **extreme points** of D' (after suitable normalization) will lie on **n LI hyperplanes** defining D' .
- Since $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ cannot be orthogonal to n **LI** vectors, so \mathbf{d} cannot lie on **n LI hyperplanes** of the $(m + n)$ hyperplanes given by,

$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$
- If $\mathbf{d} \in D'$, is an **extreme direction** of S then it should lie on **$(n - 1)$ LI hyperplanes** of the above mentioned **$(m + n)$ hyperplanes**,

- The **extreme directions of S** which are **extreme points** of D' (after suitable normalization) will lie on **n LI hyperplanes** defining D' .
- Since $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ cannot be orthogonal to n **LI** vectors, so \mathbf{d} cannot lie on **n LI hyperplanes** of the $(m + n)$ hyperplanes given by,

$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$
- If $\mathbf{d} \in D'$, is an **extreme direction** of S then it should lie on **$(n - 1)$ LI hyperplanes** of the above mentioned **$(m + n)$ hyperplanes**, and the hyperplane

$$\{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\},$$
 which gives a collection of **n LI hyperplanes**.

- The **extreme directions of S** which are **extreme points** of D' (after suitable normalization) will lie on **n LI hyperplanes** defining D' .
- Since $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ cannot be orthogonal to n **LI** vectors, so \mathbf{d} cannot lie on **n LI hyperplanes** of the $(m + n)$ hyperplanes given by,

$$\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\} \text{ for } i = 1, 2, \dots, m, \text{ and}$$

$$\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\} \text{ for } j = 1, 2, \dots, n.$$
- If $\mathbf{d} \in D'$, is an **extreme direction** of S then it should lie on **$(n - 1)$ LI hyperplanes** of the above mentioned **$(m + n)$ hyperplanes**, and the hyperplane

$$\{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\},$$
 which gives a collection of **n LI hyperplanes**.

- **Exercise:** Check that if a $\mathbf{d} \in D$ lies on $(n-1)$ LI hyperplanes (out of the $(m+n)$ defining hyperplanes of D) given by $\{H_1, \dots, H_{n-1}\}$, then $\{H, H_1, \dots, H_{n-1}\}$ is LI where $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$.

- **Exercise:** Check that if a $\mathbf{d} \in D$ lies on $(n - 1)$ LI hyperplanes (out of the $(m + n)$ defining hyperplanes of D) given by $\{H_1, \dots, H_{n-1}\}$, then $\{H, H_1, \dots, H_{n-1}\}$ is LI where $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$.
- Any $\mathbf{d} \in D$, which lies on $(n - 1)$ LI hyperplanes out of the $(m + n)$ hyperplanes given by

- **Exercise:** Check that if a $\mathbf{d} \in D$ lies on $(n - 1)$ LI hyperplanes (out of the $(m + n)$ defining hyperplanes of D) given by $\{H_1, \dots, H_{n-1}\}$, then $\{H, H_1, \dots, H_{n-1}\}$ is LI where $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$.
- Any $\mathbf{d} \in D$, which lies on $(n - 1)$ LI hyperplanes out of the $(m + n)$ hyperplanes given by $\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\}$ for $i = 1, 2, \dots, m$, and

- **Exercise:** Check that if a $\mathbf{d} \in D$ lies on $(n - 1)$ LI hyperplanes (out of the $(m + n)$ defining hyperplanes of D) given by $\{H_1, \dots, H_{n-1}\}$, then $\{H, H_1, \dots, H_{n-1}\}$ is LI where $H = \{\mathbf{d} \in \mathbb{R}^n : [1, 1, \dots, 1]\mathbf{d} = 1\}$.
- Any $\mathbf{d} \in D$, which lies on $(n - 1)$ LI hyperplanes out of the $(m + n)$ hyperplanes given by $\{\mathbf{d} \in \mathbb{R}^n : \mathbf{a}_i^T \mathbf{d} = 0\}$ for $i = 1, 2, \dots, m$, and $\{\mathbf{d} \in \mathbb{R}^n : -\mathbf{e}_j^T \mathbf{d} = 0\}$ for $j = 1, 2, \dots, n$, is an extreme direction of S .

- **Example 2: (revisited)** Consider the problem,
 $\text{Min } -x + 2y$
subject to

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of S is given by

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of S is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of S is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

If $\mathbf{d} \in D$ is an **extreme direction** of S then it has to lie on **exactly one** of the hyperplanes given by

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of S is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

If $\mathbf{d} \in D$ is an **extreme direction** of S then it has to lie on **exactly one** of the hyperplanes given by

- (i) $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of S is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

If $\mathbf{d} \in D$ is an **extreme direction** of S then it has to lie on **exactly one** of the hyperplanes given by

- (i) $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$
- (ii) $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\},$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of S is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

If $\mathbf{d} \in D$ is an **extreme direction** of S then it has to lie on **exactly one** of the hyperplanes given by

- (i) $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$
- (ii) $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\},$
- (iii) $\{\mathbf{d} \in \mathbb{R}^2 : d_1 = 0\},$

- **Example 2: (revisited)** Consider the problem,

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

Note that here the set of all **directions** of S is given by

- $D = \{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} \leq 0, [-1, 1]\mathbf{d} \leq 0, \mathbf{d} \geq \mathbf{0}\}.$

If $\mathbf{d} \in D$ is an **extreme direction** of S then it has to lie on **exactly one** of the hyperplanes given by

- (i) $\{\mathbf{d} \in \mathbb{R}^2 : [-1, -2]\mathbf{d} = 0\},$
- (ii) $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\},$
- (iii) $\{\mathbf{d} \in \mathbb{R}^2 : d_1 = 0\},$
- (iv) $\{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}.$

- Note that there exists no $d \geq 0, d \neq 0$ such that $[-1, -2]d = 0$.

- Note that there exists no $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ such that $[-1, -2]\mathbf{d} = 0$.
- Also if $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$, satisfies the condition $d_1 = 0$, then $[-1, 1]\mathbf{d} \leq 0$ cannot be satisfied, hence such a \mathbf{d} does not belong to D .

- Note that there exists no $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ such that $[-1, -2]\mathbf{d} = 0$.
- Also if $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$, satisfies the condition $d_1 = 0$, then $[-1, 1]\mathbf{d} \leq 0$ cannot be satisfied, hence such a \mathbf{d} does not belong to D .
- Hence if $\mathbf{d} \in D$, is an **extreme direction** of S then it lies on one of the following Hyperplanes:

- Note that there exists no $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ such that $[-1, -2]\mathbf{d} = 0$.
- Also if $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$, satisfies the condition $d_1 = 0$, then $[-1, 1]\mathbf{d} \leq 0$ cannot be satisfied, hence such a \mathbf{d} does not belong to D .
- Hence if $\mathbf{d} \in D$, is an **extreme direction** of S then it lies on one of the following Hyperplanes:
- $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = 0\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$.

- Note that there exists no $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ such that $[-1, -2]\mathbf{d} = \mathbf{0}$.
- Also if $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$, satisfies the condition $d_1 = 0$, then $[-1, 1]\mathbf{d} \leq \mathbf{0}$ cannot be satisfied, hence such a \mathbf{d} does not belong to D .
- Hence if $\mathbf{d} \in D$, is an **extreme direction** of S then it lies on one of the following Hyperplanes:
 - $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = \mathbf{0}\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$.
 - $\mathbf{d}' = [1, 1]^T$ and any **positive scalar multiple** of \mathbf{d}' , are **extreme directions** of $S = \text{Fea}(LPP)$.

- Note that there exists no $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ such that $[-1, -2]\mathbf{d} = \mathbf{0}$.
- Also if $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$, satisfies the condition $d_1 = 0$, then $[-1, 1]\mathbf{d} \leq \mathbf{0}$ cannot be satisfied, hence such a \mathbf{d} does not belong to D .
- Hence if $\mathbf{d} \in D$, is an **extreme direction** of S then it lies on one of the following Hyperplanes:
 - $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = \mathbf{0}\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$.
 - $\mathbf{d}' = [1, 1]^T$ and any **positive scalar multiple** of \mathbf{d}' , are **extreme directions** of $S = \text{Fea}(LPP)$.
 - $\mathbf{d}'' = [1, 0]^T$ and any **positive scalar multiple** of \mathbf{d}'' , are **extreme directions** of $S = \text{Fea}(LPP)$.

- Note that there exists no $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$ such that $[-1, -2]\mathbf{d} = \mathbf{0}$.
- Also if $\mathbf{d} \geq \mathbf{0}, \mathbf{d} \neq \mathbf{0}$, satisfies the condition $d_1 = 0$, then $[-1, 1]\mathbf{d} \leq \mathbf{0}$ cannot be satisfied, hence such a \mathbf{d} does not belong to D .
- Hence if $\mathbf{d} \in D$, is an **extreme direction** of S then it lies on one of the following Hyperplanes:
 - $\{\mathbf{d} \in \mathbb{R}^2 : [-1, 1]\mathbf{d} = \mathbf{0}\}, \{\mathbf{d} \in \mathbb{R}^2 : d_2 = 0\}$.
 - $\mathbf{d}' = [1, 1]^T$ and any **positive scalar multiple** of \mathbf{d}' , are **extreme directions** of $S = \text{Fea}(LPP)$.
 - $\mathbf{d}'' = [1, 0]^T$ and any **positive scalar multiple** of \mathbf{d}'' , are **extreme directions** of $S = \text{Fea}(LPP)$.
- These are the only **extreme directions** of $S = \text{Fea}(LPP)$.

- **Theorem:**

If $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty, then S has at least one **extreme point**.

- **Theorem:**

If $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty, then S has at least one **extreme point**.

- **Remark:** Note that the above result is **not necessarily true** for all **polyhedral sets**.

- **Theorem:**

If $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty, then S has at least one **extreme point**.

- **Remark:** Note that the above result is **not necessarily true** for all **polyhedral sets**.

For example take any **single half space**, or say a **straight line** in \mathbb{R}^n , which are **polyhedral sets**, but does not have any **extreme point**.

- **Theorem:**

If $S = \text{Fea}(LPP) = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty, then S has at least one **extreme point**.

- **Remark:** Note that the above result is **not necessarily true** for all **polyhedral sets**.

For example take any **single half space**, or say a **straight line** in \mathbb{R}^n , which are **polyhedral sets**, but does not have any **extreme point**.

- The theorem works for $\text{Fea}(LPP)$ because of the **non negativity constraints**, that is because $\text{Fea}(LPP)$ is given a supply of **n LI** , **defining hyperplanes**.

- **Exercise:** Can you find a nonempty polyhedral set S , $S \subset \mathbb{R}^2$ which has two defining hyperplanes but does not have any extreme point?

- **Exercise:** Can you find a nonempty polyhedral set S , $S \subset \mathbb{R}^2$ which has two defining hyperplanes but does not have any extreme point?
- **Exercise:** Can you find a nonempty polyhedral set S , $S \subset \mathbb{R}^3$ which has two LI defining hyperplanes but does not have any extreme point?

- **Exercise:** Can you find a **nonempty polyhedral** set S , $S \subset \mathbb{R}^2$ which has **two** **defining hyperplanes** but does not have any **extreme point**?
- **Exercise:** Can you find a **nonempty polyhedral** set S , $S \subset \mathbb{R}^3$ which has **two** **LI defining hyperplanes** but does not have any **extreme point**?
- **Exercise:** Is it possible to find a **nonempty polyhedral** set S , $S \subset \mathbb{R}^3$ which has three **LI defining hyperplanes** (not necessarily the nonnegativity constraints) but does not have any extreme point.

- **Exercise:** Can you find a nonempty polyhedral set S , $S \subset \mathbb{R}^2$ which has two defining hyperplanes but does not have any extreme point?
- **Exercise:** Can you find a nonempty polyhedral set S , $S \subset \mathbb{R}^3$ which has two LI defining hyperplanes but does not have any extreme point?
- **Exercise:** Is it possible to find a nonempty polyhedral set S , $S \subset \mathbb{R}^3$ which has three LI defining hyperplanes (not necessarily the nonnegativity constraints) but does not have any extreme point.
- **Definition:** Given S , a nonempty subset of \mathbb{R}^n , and $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in S$, $\sum_{i=1}^k \lambda_i \mathbf{x}_i$, is called a convex combination of $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$, where $0 \leq \lambda_i \leq 1$ for all $i = 1, 2, \dots, k$, and $\sum_{i=1}^k \lambda_i = 1$.

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.
- All possible convex combinations of **three non colinear points** gives a **triangle** with those points as corner points.

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.
- All possible convex combinations of **three non colinear points** gives a **triangle** with those points as corner points.
- All possible convex combinations of **four points** no three of which are colinear gives a **quadrilateral**.

- All possible **convex combinations** of **two distinct points** gives a **straight line segment** with those two points as boundary points.
- All possible convex combinations of **three non colinear points** gives a **triangle** with those points as corner points.
- All possible convex combinations of **four points** no three of which are colinear gives a **quadrilateral**.
- **Result:** Given $\phi \neq S \subset \mathbb{R}^n$, S is a **convex set** if and only if for all $k \in \mathbb{N}$, the **convex combination of any k elements** of S is again an element of S .

- **Theorem: (Representation Theorem)**

If $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty and if $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ are the extreme points of S and $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$ are the distinct extreme directions of S (the set of directions is empty if S is bounded) then $\mathbf{x} \in S$ if and only if

- **Theorem: (Representation Theorem)**

If $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty and if $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ are the extreme points of S and $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$ are the distinct extreme directions of S (the set of directions is empty if S is bounded) then $\mathbf{x} \in S$ if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$

- **Theorem: (Representation Theorem)**

If $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty and if $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ are the extreme points of S and $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$ are the distinct extreme directions of S (the set of directions is empty if S is bounded) then $\mathbf{x} \in S$ if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$

where $0 \leq \lambda_i \leq 1$ for all $i = 1, 2, \dots, k$, $\sum_i \lambda_i = 1$,

- **Theorem: (Representation Theorem)**

If $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty and if $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ are the extreme points of S and $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$ are the distinct extreme directions of S (the set of directions is empty if S is bounded) then $\mathbf{x} \in S$ if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$

where $0 \leq \lambda_i \leq 1$ for all $i = 1, 2, \dots, k$, $\sum_i \lambda_i = 1$,
and $\mu_j \geq 0$, for all $j = 1, 2, \dots, r$.

- **Theorem: (Representation Theorem)**

If $S = \{\mathbf{x} \in \mathbb{R}^n : A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ is nonempty and if $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ are the extreme points of S and $\mathbf{d}_1, \mathbf{d}_2, \dots, \mathbf{d}_r$ are the distinct extreme directions of S (the set of directions is empty if S is bounded) then $\mathbf{x} \in S$ if and only if

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \sum_{j=1}^r \mu_j \mathbf{d}_j$$

where $0 \leq \lambda_i \leq 1$ for all $i = 1, 2, \dots, k$, $\sum_i \lambda_i = 1$, and $\mu_j \geq 0$, for all $j = 1, 2, \dots, r$.

- That is, $\mathbf{x} \in S \Leftrightarrow \mathbf{x}$ can be written as a convex combination of the extreme points of S plus a non negative linear combination of the extreme directions of S .

- **Example 2: (revisited)** Consider the problem with $Fea(LPP)$,

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP)$,
 $x + 2y \geq 1$

- **Example 2: (revisited)** Consider the problem with

$Fea(LPP),$

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

- **Example 2: (revisited)** Consider the problem with

$Fea(LPP),$

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP),$
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- The **extreme points** are $[1, 0]^T, [0, 1]^T$ and $[0, \frac{1}{2}]^T$.

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP)$,
 $x + 2y \geq 1$
 $-x + y \leq 1$,
 $x \geq 0, y \geq 0$.
- The **extreme points** are $[1, 0]^T$, $[0, 1]^T$ and $[0, \frac{1}{2}]^T$.
- Distinct extreme directions are $[1, 0]^T$, $[1, 1]^T$.

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP),$
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- The **extreme points** are $[1, 0]^T, [0, 1]^T$ and $[0, \frac{1}{2}]^T.$
- Distinct extreme directions are $[1, 0]^T, [1, 1]^T.$
- $[0, \frac{3}{4}]^T \in Fea(LPP),$

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP),$
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- The **extreme points** are $[1, 0]^T, [0, 1]^T$ and $[0, \frac{1}{2}]^T$.
- Distinct extreme directions are $[1, 0]^T, [1, 1]^T$.
- $[0, \frac{3}{4}]^T \in Fea(LPP),$
 $[0, \frac{3}{4}]^T = \frac{1}{2}[0, \frac{1}{2}]^T + \frac{1}{2}[0, 1]^T,$

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP),$
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- The **extreme points** are $[1, 0]^T, [0, 1]^T$ and $[0, \frac{1}{2}]^T$.
- Distinct extreme directions are $[1, 0]^T, [1, 1]^T$.
- $[0, \frac{3}{4}]^T \in Fea(LPP),$
 $[0, \frac{3}{4}]^T = \frac{1}{2}[0, \frac{1}{2}]^T + \frac{1}{2}[0, 1]^T,$
 $[\frac{3}{2}, 0]^T = 1[1, 0]^T + \frac{1}{2}[1, 0]^T$

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP),$
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- The **extreme points** are $[1, 0]^T, [0, 1]^T$ and $[0, \frac{1}{2}]^T$.
- Distinct extreme directions are $[1, 0]^T, [1, 1]^T$.
- $[0, \frac{3}{4}]^T \in Fea(LPP),$
 $[0, \frac{3}{4}]^T = \frac{1}{2}[0, \frac{1}{2}]^T + \frac{1}{2}[0, 1]^T,$
 $[\frac{3}{2}, 0]^T = 1[1, 0]^T + \frac{1}{2}[1, 0]^T$

- **Example 2: (revisited)** Consider the problem with
 $Fea(LPP),$
 $x + 2y \geq 1$
 $-x + y \leq 1,$
 $x \geq 0, y \geq 0.$
- The **extreme points** are $[1, 0]^T, [0, 1]^T$ and $[0, \frac{1}{2}]^T$.
- Distinct extreme directions are $[1, 0]^T, [1, 1]^T$.
- $[0, \frac{3}{4}]^T \in Fea(LPP),$
 $[0, \frac{3}{4}]^T = \frac{1}{2}[0, \frac{1}{2}]^T + \frac{1}{2}[0, 1]^T,$
 $[\frac{3}{2}, 0]^T = 1[1, 0]^T + \frac{1}{2}[1, 0]^T$
 $[2, 3]^T = 1[0, 1]^T + 2[1, 1]^T.$

- **Example 2: (revisited)** Consider the problem with

$$\text{Fea}(LPP),$$

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- The **extreme points** are $[1, 0]^T$, $[0, 1]^T$ and $[0, \frac{1}{2}]^T$.

- Distinct extreme directions are $[1, 0]^T$, $[1, 1]^T$.

- $[0, \frac{3}{4}]^T \in \text{Fea}(LPP)$,

$$[0, \frac{3}{4}]^T = \frac{1}{2}[0, \frac{1}{2}]^T + \frac{1}{2}[0, 1]^T,$$

$$[\frac{3}{2}, 0]^T = 1[1, 0]^T + \frac{1}{2}[1, 0]^T$$

$$[2, 3]^T = 1[0, 1]^T + 2[1, 1]^T.$$

$$[3, 3]^T = 1[0, 1]^T + 2[1, 1]^T + 1[1, 0]^T$$

- **Example 2: (revisited)** Consider the problem with

$$\text{Fea}(LPP),$$

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- The **extreme points** are $[1, 0]^T$, $[0, 1]^T$ and $[0, \frac{1}{2}]^T$.

- Distinct extreme directions are $[1, 0]^T$, $[1, 1]^T$.

- $[0, \frac{3}{4}]^T \in \text{Fea}(LPP)$,

$$[0, \frac{3}{4}]^T = \frac{1}{2}[0, \frac{1}{2}]^T + \frac{1}{2}[0, 1]^T,$$

$$[\frac{3}{2}, 0]^T = 1[1, 0]^T + \frac{1}{2}[1, 0]^T$$

$$[2, 3]^T = 1[0, 1]^T + 2[1, 1]^T.$$

$$[3, 3]^T = 1[0, 1]^T + 2[1, 1]^T + 1[1, 0]^T$$

$$[3, 3]^T = \frac{1}{2}[0, 1]^T + \frac{1}{2}[1, 0]^T + \frac{3}{2}[1, 1]^T$$

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty bounded set then any $\mathbf{x} \in S$ is a convex combination of the extreme points of S .

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty bounded set then any $\mathbf{x} \in S$ is a convex combination of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty bounded set then any $\mathbf{x} \in S$ is a convex combination of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$
 $3x + 2y \leq 6$

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty bounded set then any $\mathbf{x} \in S$ is a convex combination of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$
 $3x + 2y \leq 6$
 $x + 2y \leq 4$

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty bounded set then any $\mathbf{x} \in S$ is a convex combination of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$
 $3x + 2y \leq 6$
 $x + 2y \leq 4$
 $x \geq 0, y \geq 0.$

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty bounded set then any $\mathbf{x} \in S$ is a convex combination of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$
 $3x + 2y \leq 6$
 $x + 2y \leq 4$
 $x \geq 0, y \geq 0.$
- The extreme points are $[2, 0]^T, [0, 2]^T, [0, 0]^T$ and $[1, \frac{3}{2}]^T$.

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty bounded set then any $\mathbf{x} \in S$ is a convex combination of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$
- The extreme points are $[2, 0]^T$, $[0, 2]^T$, $[0, 0]^T$ and $[1, \frac{3}{2}]^T$.

$$[1, \frac{1}{2}]^T = \frac{1}{2}[2, 0]^T + \frac{1}{4}[0, 2]^T + \frac{1}{4}[0, 0]^T.$$

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty **bounded set** then any $\mathbf{x} \in S$ is a **convex combination** of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

- The extreme points are $[2, 0]^T$, $[0, 2]^T$, $[0, 0]^T$ and $[1, \frac{3}{2}]^T$.
 $[1, \frac{1}{2}]^T = \frac{1}{2}[2, 0]^T + \frac{1}{4}[0, 2]^T + \frac{1}{4}[0, 0]^T$.
- $[1, \frac{1}{2}]^T = \frac{1}{3}[2, 0]^T + \frac{1}{3}[1, \frac{3}{2}]^T + \frac{1}{3}[0, 0]^T$.

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty **bounded set** then any $\mathbf{x} \in S$ is a **convex combination** of the extreme points of S .

- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

- The extreme points are $[2, 0]^T$, $[0, 2]^T$, $[0, 0]^T$ and $[1, \frac{3}{2}]^T$.
 $[1, \frac{1}{2}]^T = \frac{1}{2}[2, 0]^T + \frac{1}{4}[0, 2]^T + \frac{1}{4}[0, 0]^T$.
- $[1, \frac{1}{2}]^T = \frac{1}{3}[2, 0]^T + \frac{1}{3}[1, \frac{3}{2}]^T + \frac{1}{3}[0, 0]^T$.
- $[1, \frac{1}{2}]^T = \frac{5}{12}[2, 0]^T + \frac{3}{24}[0, 2]^T + \frac{1}{6}[1, \frac{3}{2}]^T + \frac{17}{24}[0, 0]^T$.

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty **bounded set** then any $\mathbf{x} \in S$ is a **convex combination** of the extreme points of S .
- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$
- The extreme points are $[2, 0]^T$, $[0, 2]^T$, $[0, 0]^T$ and $[1, \frac{3}{2}]^T$.

$$[1, \frac{1}{2}]^T = \frac{1}{2}[2, 0]^T + \frac{1}{4}[0, 2]^T + \frac{1}{4}[0, 0]^T.$$
- $$[1, \frac{1}{2}]^T = \frac{1}{3}[2, 0]^T + \frac{1}{3}[1, \frac{3}{2}]^T + \frac{1}{3}[0, 0]^T.$$
- $$[1, \frac{1}{2}]^T = \frac{5}{12}[2, 0]^T + \frac{3}{24}[0, 2]^T + \frac{1}{6}[1, \frac{3}{2}]^T + \frac{17}{24}[0, 0]^T.$$
- **Observation 7:** Since D' , the set of distinct directions of S (if it is nonempty) is a **bounded set** ($\mathbf{d} \geq \mathbf{0}$ and $\sum_{i=1}^n d_i = 1$),

- **Observation 6:** If $S = \text{Fea}(LPP)$ is a nonempty **bounded set** then any $\mathbf{x} \in S$ is a **convex combination** of the extreme points of S .

- **Example 1:(revisited)** Given the $\text{Fea}(LPP)$

$$3x + 2y \leq 6$$

$$x + 2y \leq 4$$

$$x \geq 0, y \geq 0.$$

- The extreme points are $[2, 0]^T$, $[0, 2]^T$, $[0, 0]^T$ and $[1, \frac{3}{2}]^T$.
 $[1, \frac{1}{2}]^T = \frac{1}{2}[2, 0]^T + \frac{1}{4}[0, 2]^T + \frac{1}{4}[0, 0]^T$.
- $[1, \frac{1}{2}]^T = \frac{1}{3}[2, 0]^T + \frac{1}{3}[1, \frac{3}{2}]^T + \frac{1}{3}[0, 0]^T$.
- $[1, \frac{1}{2}]^T = \frac{5}{12}[2, 0]^T + \frac{3}{24}[0, 2]^T + \frac{1}{6}[1, \frac{3}{2}]^T + \frac{17}{24}[0, 0]^T$.
- **Observation 7:** Since D' , the set of distinct directions of S (if it is nonempty) is a **bounded set** ($\mathbf{d} \geq \mathbf{0}$ and $\sum_{i=1}^n d_i = 1$),
 so any $\mathbf{d} \in D'$ is a **convex combination** of the **extreme points** of D' .

- So any direction $\mathbf{d} \in D$ is a nonnegative linear combination of the extreme directions of S .

- So any direction $\mathbf{d} \in D$ is a nonnegative linear combination of the extreme directions of S .
- **Observation 8:** Note that if there exists a $\mathbf{d} \in D$ such that $\mathbf{c}^T \mathbf{d} < 0$ then the LPP(*)

- So any direction $\mathbf{d} \in D$ is a nonnegative linear combination of the extreme directions of S .
- **Observation 8:** Note that if there exists a $\mathbf{d} \in D$ such that $\mathbf{c}^T \mathbf{d} < 0$ then the LPP(*)
 ((*) Min $\mathbf{c}^T \mathbf{x}$,

- So any direction $\mathbf{d} \in D$ is a nonnegative linear combination of the extreme directions of S .
- **Observation 8:** Note that if there exists a $\mathbf{d} \in D$ such that $\mathbf{c}^T \mathbf{d} < 0$ then the LPP(*)
 ((*) Min $\mathbf{c}^T \mathbf{x}$, subject to $A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$)

- So any direction $\mathbf{d} \in D$ is a nonnegative linear combination of the extreme directions of S .
- **Observation 8:** Note that if there exists a $\mathbf{d} \in D$ such that $\mathbf{c}^T \mathbf{d} < 0$ then the LPP(*)
 ((*) Min $\mathbf{c}^T \mathbf{x}$, subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$)
 does not have an optimal solution.

- So any direction $\mathbf{d} \in D$ is a nonnegative linear combination of the extreme directions of S .
- **Observation 8:** Note that if there exists a $\mathbf{d} \in D$ such that $\mathbf{c}^T \mathbf{d} < 0$ then the LPP(*)
 ((*) Min $\mathbf{c}^T \mathbf{x}$, subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$)
 does not have an optimal solution.
- If LPP(*) has an optimal solution then $\mathbf{c}^T \mathbf{d} \geq 0$ for all $\mathbf{d} \in D$.

- If $\mathbf{c}^T \mathbf{d}_j \geq 0$ for all extreme directions \mathbf{d}_j of the nonempty and unbounded feasible region S of a LPP, then it implies $\mathbf{c}^T \mathbf{d} \geq 0$ for all directions $\mathbf{d} \in D$, of the feasible region S .

- If $\mathbf{c}^T \mathbf{d}_j \geq 0$ for all extreme directions \mathbf{d}_j of the nonempty and unbounded feasible region S of a LPP, then it implies $\mathbf{c}^T \mathbf{d} \geq 0$ for all directions $\mathbf{d} \in D$, of the feasible region S .
- **Observation 9:** From the representation theorem the converse follows that is if $S \neq \phi$ and $\mathbf{c}^T \mathbf{d}_j \geq 0$ for all $j = 1, 2, \dots, r$, then LPP(*) has an optimal solution, and atleast one optimal solution is attained at an extreme point of S .

- **Observation 10:** From the representation theorem it follows that if $S = \text{Fea}(LPP)$ is nonempty and bounded then the $LPP(*)$ has an optimal solution and the optimal value is attained in at least one extreme point.

- **Observation 10:** From the representation theorem it follows that if $S = \text{Fea}(LPP)$ is nonempty and bounded then the $LPP(*)$ has an optimal solution and the optimal value is attained in atleast one extreme point.

From the above observations we can conclude the following:

- **Observation 10:** From the representation theorem it follows that if $S = \text{Fea}(LPP)$ is nonempty and bounded then the $LPP(*)$ has an optimal solution and the optimal value is attained in atleast one extreme point.

From the above observations we can conclude the following:

- **Conclusion 1:** If $S = \text{Fea}(LPP) \neq \phi$, then the $LPP (*)$ has an optimal solution if and only if one of the following is true:

- **Observation 10:** From the representation theorem it follows that if $S = \text{Fea}(LPP)$ is nonempty and bounded then the $LPP(*)$ has an optimal solution and the optimal value is attained in at least one extreme point.

From the above observations we can conclude the following:

- **Conclusion 1:** If $S = \text{Fea}(LPP) \neq \emptyset$, then the $LPP (*)$ has an optimal solution if and only if one of the following is true:
- (i) $S = \text{Fea}(LPP)$ is bounded (also seen before by using extreme value theorem)
- (ii) $S = \text{Fea}(LPP)$ is unbounded and $\mathbf{c}^T \mathbf{d}_j \geq 0$ for all extreme directions \mathbf{d}_j of the feasible region S .

- **Observation 10:** From the representation theorem it follows that if $S = \text{Fea}(LPP)$ is nonempty and bounded then the $LPP(*)$ has an optimal solution and the optimal value is attained in at least one extreme point.

From the above observations we can conclude the following:

- **Conclusion 1:** If $S = \text{Fea}(LPP) \neq \emptyset$, then the $LPP (*)$ has an optimal solution if and only if one of the following is true:
- (i) $S = \text{Fea}(LPP)$ is bounded (also seen before by using extreme value theorem)
- (ii) $S = \text{Fea}(LPP)$ is unbounded and $\mathbf{c}^T \mathbf{d}_j \geq 0$ for all extreme directions \mathbf{d}_j of the feasible region S .

- **Conclusion 2:** If LPP (*) has an optimal solution then there exists an extreme point of the feasible region S , which is an optimal solution.

- **Conclusion 2:** If LPP (*) has an **optimal solution** then there exists an **extreme point** of the feasible region S , which is an **optimal solution**.
- **Exercise:** Give an example of a **nonlinear** function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}$ is a **closed** and **bounded polyhedral** subset of \mathbb{R} , (what are these sets?) such that f has a **minimum value** in S but the minimum value is not attained at any **extreme point** of S .

- **Conclusion 2:** If LPP (*) has an **optimal solution** then there exists an **extreme point** of the feasible region S , which is an **optimal solution**.
- **Exercise:** Give an example of a **nonlinear** function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}$ is a **closed** and **bounded polyhedral** subset of \mathbb{R} , (what are these sets?) such that f has a **minimum value** in S but the minimum value is not attained at any **extreme point** of S .
- **Conclusion 3:** If $S = \text{Fea}(LPP)$ is nonempty,

- **Conclusion 2:** If LPP (*) has an optimal solution then there exists an extreme point of the feasible region S , which is an optimal solution.
- **Exercise:** Give an example of a nonlinear function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}$ is a closed and bounded polyhedral subset of \mathbb{R} , (what are these sets?) such that f has a minimum value in S but the minimum value is not attained at any extreme point of S .
- **Conclusion 3:** If $S = \text{Fea}(LPP)$ is nonempty, and there exists an $M \in \mathbb{R}$ such that for all $\mathbf{x} \in S$, $\mathbf{c}^T \mathbf{x} \geq M$,

- **Conclusion 2:** If LPP (*) has an optimal solution then there exists an extreme point of the feasible region S , which is an optimal solution.
- **Exercise:** Give an example of a nonlinear function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}$ is a closed and bounded polyhedral subset of \mathbb{R} , (what are these sets?) such that f has a minimum value in S but the minimum value is not attained at any extreme point of S .
- **Conclusion 3:** If $S = \text{Fea}(LPP)$ is nonempty, and there exists an $M \in \mathbb{R}$ such that for all $\mathbf{x} \in S$, $\mathbf{c}^T \mathbf{x} \geq M$, then the LPP (*) has an optimal solution.

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- The distinct extreme directions are $\mathbf{d}' = [1, 0]^T$ and $\mathbf{d}'' = [1, 1]^T$.

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- The distinct extreme directions are $\mathbf{d}' = [1, 0]^T$ and $\mathbf{d}'' = [1, 1]^T$.
- $\mathbf{c}^T \mathbf{d}' = -1$, hence this problem does not have an optimal solution.

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- The distinct extreme directions are $\mathbf{d}' = [1, 0]^T$ and $\mathbf{d}'' = [1, 1]^T$.
- $\mathbf{c}^T \mathbf{d}' = -1$, hence this problem does not have an optimal solution.
- If we change the above \mathbf{c} to $[1, -1]^T$ then $\mathbf{c}^T \mathbf{d}' = 1$ and $\mathbf{c}^T \mathbf{d}'' = 0$,

- **Example 2: (revisited)** Consider the problem:

$$\text{Min } -x + 2y$$

subject to

$$x + 2y \geq 1$$

$$-x + y \leq 1,$$

$$x \geq 0, y \geq 0.$$

- The distinct extreme directions are $\mathbf{d}' = [1, 0]^T$ and $\mathbf{d}'' = [1, 1]^T$.
- $\mathbf{c}^T \mathbf{d}' = -1$, hence this problem does not have an optimal solution.
- If we change the above \mathbf{c} to $[1, -1]^T$ then $\mathbf{c}^T \mathbf{d}' = 1$ and $\mathbf{c}^T \mathbf{d}'' = 0$, and the new problem has an optimal solution.

- To understand the significance of the previous results solve the following exercises.

- To understand the significance of the previous results solve the following exercises.
- **Exercise:** Give an example of a **linear** function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}$ is not a polyhedral subset of \mathbb{R} , such that $f(x) \geq 1$ but f **does not** have a minimum value in S .

- To understand the significance of the previous results solve the following exercises.
- **Exercise:** Give an example of a **linear** function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}$ is not a polyhedral subset of \mathbb{R} , such that $f(x) \geq 1$ but f **does not** have a minimum value in S .
- **Exercise:** Give an example of a **nonlinear** function $f : S \rightarrow \mathbb{R}$, where $S \subset \mathbb{R}$ is a **polyhedral** subset of \mathbb{R} , such that $f(x) \geq 1$ but f **does not** have a minimum value in S .

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as
(**) $\text{Max } \mathbf{c}^T \mathbf{x}$

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as
(**) $\text{Max } \mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as
(**) $\text{Max } \mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- **Conclusion 1a:** If $S = \text{Fea}(LPP) \neq \phi$, then the LPP (**) has an optimal solution if and only if one of the following is true:

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as
(**) $\text{Max } \mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- **Conclusion 1a:** If $S = \text{Fea}(LPP) \neq \phi$, then the LPP (**) has an optimal solution if and only if one of the following is true:
 - (i) $S = \text{Fea}(LPP)$ is bounded

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.

- **Conclusion 1a:** If $S = \text{Fea}(LPP) \neq \phi$, then the LPP (**) has an optimal solution if and only if one of the following is true:

(i) $S = \text{Fea}(LPP)$ is bounded

(ii) $S = \text{Fea}(LPP)$ is unbounded and $\mathbf{c}^T \mathbf{d}_j \leq 0$ for all extreme directions \mathbf{d}_j of the feasible region S .

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.

- **Conclusion 1a:** If $S = \text{Fea}(LPP) \neq \phi$, then the LPP (**) has an optimal solution if and only if one of the following is true:

(i) $S = \text{Fea}(LPP)$ is bounded

(ii) $S = \text{Fea}(LPP)$ is unbounded and $\mathbf{c}^T \mathbf{d}_j \leq 0$ for all extreme directions \mathbf{d}_j of the feasible region S .

- **Conclusion 2a:** If a LPP (**) has an optimal solution then there exists an extreme point of the feasible region S , which is an optimal solution.

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.

- **Conclusion 1a:** If $S = \text{Fea}(LPP) \neq \emptyset$, then the LPP (**) has an optimal solution if and only if one of the following is true:

(i) $S = \text{Fea}(LPP)$ is bounded

(ii) $S = \text{Fea}(LPP)$ is unbounded and $\mathbf{c}^T \mathbf{d}_j \leq 0$ for all extreme directions \mathbf{d}_j of the feasible region S .

- **Conclusion 2a:** If a LPP (**) has an optimal solution then there exists an extreme point of the feasible region S , which is an optimal solution.
- **Conclusion 3a:** If $S = \text{Fea}(LPP)$ is nonempty, and there exists an $M \in \mathbb{R}$ such that for all $\mathbf{x} \in S, \mathbf{c}^T \mathbf{x} \leq M$,

- We can come to similar conclusions if we consider a linear programming problem, LPP(**) as

$$(**) \text{Max } \mathbf{c}^T \mathbf{x}$$

subject to $\mathbf{Ax} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.

- **Conclusion 1a:** If $S = \text{Fea}(LPP) \neq \emptyset$, then the LPP (**) has an optimal solution if and only if one of the following is true:

(i) $S = \text{Fea}(LPP)$ is bounded

(ii) $S = \text{Fea}(LPP)$ is unbounded and $\mathbf{c}^T \mathbf{d}_j \leq 0$ for all extreme directions \mathbf{d}_j of the feasible region S .

- **Conclusion 2a:** If a LPP (**) has an optimal solution then there exists an extreme point of the feasible region S , which is an optimal solution.
- **Conclusion 3a:** If $S = \text{Fea}(LPP)$ is nonempty, and there exists an $M \in \mathbb{R}$ such that for all $\mathbf{x} \in S, \mathbf{c}^T \mathbf{x} \leq M$, then the LPP (**) has an optimal solution.