

The Matrix Singular Value Decomposition

Singular Value Decomposition(SVD)

The Singular Value Decomposition (SVD) of a matrix $A \in \mathbb{R}^{n \times m}$ is a decomposition of the form

$$A = U\Sigma V^T$$

where $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ are orthogonal matrices and $\Sigma := \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_p) \in \mathbb{R}^{n \times m}$ is a diagonal matrix with

$$\sigma_1 \geq \sigma_2 \geq \dots \sigma_p \geq 0$$

for $p = \min\{n, m\}$.

Singular Value Decomposition(SVD)

The Singular Value Decomposition (SVD) of a matrix $A \in \mathbb{R}^{n \times m}$ is a decomposition of the form

$$A = U \Sigma V^T$$

where $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ are orthogonal matrices and $\Sigma := \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_p) \in \mathbb{R}^{n \times m}$ is a diagonal matrix with

$$\sigma_1 \geq \sigma_2 \geq \dots \sigma_p \geq 0$$

for $p = \min\{n, m\}$.

If A is a complex $n \times m$ matrix, then the above decomposition holds with T replaced by $*$ so that U and V are unitary matrices. However Σ remains unchanged.

Singular Value Decomposition(SVD)

The Singular Value Decomposition (SVD) of a matrix $A \in \mathbb{R}^{n \times m}$ is a decomposition of the form

$$A = U\Sigma V^T$$

where $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ are orthogonal matrices and $\Sigma := \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_p) \in \mathbb{R}^{n \times m}$ is a diagonal matrix with

$$\sigma_1 \geq \sigma_2 \geq \dots \sigma_p \geq 0$$

for $p = \min\{n, m\}$.

If A is a complex $n \times m$ matrix, then the above decomposition holds with T replaced by $*$ so that U and V are unitary matrices. However Σ remains unchanged.

The numbers $\sigma_1, \sigma_2, \dots, \sigma_p$ are called the singular values of A .

Singular Value Decomposition(SVD)

Every matrix has an SVD. For example, the SVD of

$$A := \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}^T,$$

Singular Value Decomposition(SVD)

Every matrix has an SVD. For example, the SVD of

$$A := \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}^T,$$

$$B := \begin{bmatrix} 1 & 2 \end{bmatrix} = [1] \begin{bmatrix} \sqrt{5} & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{bmatrix}^T,$$

Singular Value Decomposition(SVD)

Every matrix has an SVD. For example, the SVD of

$$A := \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}^T,$$

$$B := \begin{bmatrix} 1 & 2 \end{bmatrix} = [1] \begin{bmatrix} \sqrt{5} & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{bmatrix}^T,$$

$$D := \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix} [1]^T.$$

Singular Value Decomposition(SVD)

Every matrix has an SVD. For example, the SVD of

$$A := \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}^T,$$

$$B := \begin{bmatrix} 1 & 2 \end{bmatrix} = [1] \begin{bmatrix} \sqrt{5} & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{5}} & -\frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{bmatrix}^T,$$

$$D := \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix} [1]^T.$$

Clearly if $A = U \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_p) V^T$ is the SVD of A and $\text{rank } A = r$, then the first r singular values $\sigma_1 \geq \dots \geq \sigma_r > 0$ with $\sigma_k = 0$ for $k = r + 1, \dots, p$ if $r < p$.

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Theorem Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$ with $\text{rank } A = r$.

(a) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Theorem Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$ with $\text{rank } A = r$.

(a) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$ (Exercise!)

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Theorem Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$ with $\text{rank } A = r$.

- (a) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$ (Exercise!)
- (b) $A^* = V\Sigma^T U^*$ is an SVD of A^* .

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Theorem Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$ with $\text{rank } A = r$.

(a) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$ (Exercise!)

(b) $A^* = V\Sigma^T U^*$ is an SVD of A^* . (Exercise!)

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Theorem Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$ with $\text{rank } A = r$.

- (a) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$ (Exercise!)
- (b) $A^* = V\Sigma^T U^*$ is an SVD of A^* . (Exercise!)
- (c)

$$\begin{aligned} R(A) &= \text{span}\{u_1, \dots, u_r\}, & N(A) &= \text{span}\{v_{r+1}, \dots, v_m\} \\ R(A^*) &= \text{span}\{v_1, \dots, v_r\} & N(A^*) &= \text{span}\{u_{r+1}, \dots, u_n\}. \end{aligned}$$

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Theorem Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$ with $\text{rank } A = r$.

(a) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$ (Exercise!)

(b) $A^* = V\Sigma^T U^*$ is an SVD of A^* . (Exercise!)

(c)

$$\begin{aligned} R(A) &= \text{span}\{u_1, \dots, u_r\}, & N(A) &= \text{span}\{v_{r+1}, \dots, v_m\} \\ R(A^*) &= \text{span}\{v_1, \dots, v_r\} & N(A^*) &= \text{span}\{u_{r+1}, \dots, u_n\}. \end{aligned}$$

(d) $\|A\|_2 = \sigma_1$.

Singular Value Decomposition(SVD)

If $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_m]$, then for $i = 1, \dots, p$,

$$Av_i = \sigma_i u_i \text{ and } u_i^* A = \sigma_i v_i^*$$

Hence u_i and v_i are respectively left and right singular vectors of A corresponding to σ_i .

Theorem Every matrix has a Singular Value Decomposition.

Theorem Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$ with $\text{rank } A = r$.

(a) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$ (Exercise!)

(b) $A^* = V\Sigma^T U^*$ is an SVD of A^* . (Exercise!)

(c)

$$\begin{aligned} R(A) &= \text{span}\{u_1, \dots, u_r\}, & N(A) &= \text{span}\{v_{r+1}, \dots, v_m\} \\ R(A^*) &= \text{span}\{v_1, \dots, v_r\} & N(A^*) &= \text{span}\{u_{r+1}, \dots, u_n\}. \end{aligned}$$

(d) $\|A\|_2 = \sigma_1$.

(e) $\|A\|_F = \sqrt{\sum_{k=1}^r \sigma_k^2}$.

(Here $\mathbb{F} = \mathbb{R}$ or $\mathbb{F} = \mathbb{C}$.)

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{F}^{n \times m}$.

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{R}^{n \times m}$.

- (a) If A is square and nonsingular, then $A^{-1} = (VF)(F\Sigma^{-1}F)(UF)^*$ is an SVD of A^{-1} and where F is the $n \times n$ 'flip' matrix and $\|A^{-1}\|_2 = \frac{1}{\sigma_n}$.

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{R}^{n \times m}$.

- (a) If A is square and nonsingular, then $A^{-1} = (VF)(F\Sigma^{-1}F)(UF)^*$ is an SVD of A^{-1} and where F is the $n \times n$ 'flip' matrix and $\|A^{-1}\|_2 = \frac{1}{\sigma_n}$. (Exercise!)

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{R}^{n \times m}$.

- (a) If A is square and nonsingular, then $A^{-1} = (VF)(F\Sigma^{-1}F)(UF)^*$ is an SVD of A^{-1} and where F is the $n \times n$ 'flip' matrix and $\|A^{-1}\|_2 = \frac{1}{\sigma_n}$. (Exercise!)
- (b) If $p = \min\{m, n\}$, then assuming $\kappa_2(A) = \frac{\max \text{mag } A^T}{\min \text{mag } A^T}$ if $n < m$,
- $$\kappa_2(A) = \begin{cases} \frac{\sigma_1}{\sigma_p} & \text{if rank } A = p, \\ \infty & \text{otherwise} \end{cases}$$

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{R}^{n \times m}$.

- (a) If A is square and nonsingular, then $A^{-1} = (VF)(F\Sigma^{-1}F)(UF)^*$ is an SVD of A^{-1} and where F is the $n \times n$ 'flip' matrix and $\|A^{-1}\|_2 = \frac{1}{\sigma_n}$. (Exercise!)
- (b) If $p = \min\{m, n\}$, then assuming $\kappa_2(A) = \frac{\max \text{mag } A^T}{\min \text{mag } A^T}$ if $n < m$,
- $$\kappa_2(A) = \begin{cases} \frac{\sigma_1}{\sigma_p} & \text{if rank } A = p, \\ \infty & \text{otherwise} \end{cases} \quad \text{(Exercise!)}$$

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{R}^{n \times m}$.

- (a) If A is square and nonsingular, then $A^{-1} = (VF)(F\Sigma^{-1}F)(UF)^*$ is an SVD of A^{-1} and where F is the $n \times n$ 'flip' matrix and $\|A^{-1}\|_2 = \frac{1}{\sigma_n}$. (Exercise!)
- (b) If $p = \min\{m, n\}$, then assuming $\kappa_2(A) = \frac{\max \text{mag } A^T}{\min \text{mag } A^T}$ if $n < m$,
$$\kappa_2(A) = \begin{cases} \frac{\sigma_1}{\sigma_p} & \text{if rank } A = p, \\ \infty & \text{otherwise} \end{cases} \quad \text{(Exercise!)}$$
- (c) Assuming, $\sigma_k = 0$ for $k > \min\{m, n\}$, $A^*Av_i = \sigma_i^2v_i$, $i = 1, \dots, m$, and $AA^*u_j = \sigma_j^2u_j$, $j = 1, \dots, n$.

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{R}^{n \times m}$.

- (a) If A is square and nonsingular, then $A^{-1} = (VF)(F\Sigma^{-1}F)(UF)^*$ is an SVD of A^{-1} and where F is the $n \times n$ 'flip' matrix and $\|A^{-1}\|_2 = \frac{1}{\sigma_n}$. (Exercise!)
- (b) If $p = \min\{m, n\}$, then assuming $\kappa_2(A) = \frac{\max \text{mag } A^T}{\min \text{mag } A^T}$ if $n < m$,
$$\kappa_2(A) = \begin{cases} \frac{\sigma_1}{\sigma_p} & \text{if rank } A = p, \\ \infty & \text{otherwise} \end{cases} \quad \text{(Exercise!)}$$
- (c) Assuming, $\sigma_k = 0$ for $k > \min\{m, n\}$, $A^*Av_i = \sigma_i^2v_i$, $i = 1, \dots, m$, and $AA^*u_j = \sigma_j^2u_j$, $j = 1, \dots, n$. (Exercise!)

Singular Value Decomposition(SVD)

Corollary Let $A = U\Sigma V^*$ be an SVD of $A \in \mathbb{R}^{n \times m}$.

- (a) If A is square and nonsingular, then $A^{-1} = (VF)(F\Sigma^{-1}F)(UF)^*$ is an SVD of A^{-1} and where F is the $n \times n$ 'flip' matrix and $\|A^{-1}\|_2 = \frac{1}{\sigma_n}$. (Exercise!)
- (b) If $p = \min\{m, n\}$, then assuming $\kappa_2(A) = \frac{\max \text{mag } A^T}{\min \text{mag } A^T}$ if $n < m$,
$$\kappa_2(A) = \begin{cases} \frac{\sigma_1}{\sigma_p} & \text{if rank } A = p, \\ \infty & \text{otherwise} \end{cases} \quad \text{(Exercise!)}$$
- (c) Assuming, $\sigma_k = 0$ for $k > \min\{m, n\}$, $A^*Av_i = \sigma_i^2v_i$, $i = 1, \dots, m$, and $AA^*u_j = \sigma_j^2u_j$, $j = 1, \dots, n$. (Exercise!)
- (d) If $n = m$ and A is a singular matrix, then for any $\epsilon > 0$, there exists a nonsingular matrix $B \in \mathbb{R}^{n \times n}$ such that $\|A - B\|_2 < \epsilon$.