CSIT113 Problem Solving

Workshop - Week 9

Sorting Practice

Consider the following list of numbers:





- Sort it using:
 - 1. Selection sort
 - 2. Insertion sort
 - 3. Bubble sort
- Count the number of operations each time.
- Which is the best sort?

Selection Sort.

- Selection sort uses the following strategy:
 - 1. Start with the whole list.
 - 2. Find the smallest element in the list.
 - 3. Swap the first element with the smallest.
 - 4. Shorten the list by ignoring its first element (because it is in the right place).
 - 5. If the list has only one element, stop.
 - 6. Otherwise, go to step 1. with the shorter list.







Start	9	11	8	9	13	18	3	10	
1 st cycle	3	11	8	9	13	18	9	10	1 swap, 7 comparisons
2 nd cycle	3	8	11	9	13	18	9	10	1 swap, 6 comparisons
3 rd cycle	3	8	9	11	13	18	9	10	1 swap, 5 comparisons
4 th cycle	3	8	9	9	13	18	11	10	1 swap, 4 comparisons
5 th cycle	3	8	9	9	10	18	11	13	1 swap, 3 comparisons
6 th cycle	3	8	9	9	10	11	18	13	1 swap, 2 comparisons
7 th cycle	3	8	9	9	10	11	13	18	1 swap, 1 comparisons
	We have o	done!							
	3	8	9	9	10	11	13	18	7 swaps, 28 comparisons

Insertion Sort.

- Insertion sort uses the following strategy:
 - 1. Start with the second element in the list.
 - 2. Insert it in the right place in the preceding list.
 - 3. Repeat with the next unsorted element.
 - 4. Keep going until we have placed the last element in the list.



Start with the second element



- Compare to the preceding list (only one element 9): 1 comparison
- It is in the right position. So do nothing!



• Finish 1st cycle






Start with the next element



- Compare to the preceding list: 2 comparisons
- Insert into the right position: 2 swaps



• Finish 2st cycle





 8
 9
 11
 9
 13
 18
 3
 10

Compare to the preceding list: 2 comparisons

• Insert into the right position: 1 swap

 8
 9
 9
 11
 13
 18
 3
 10


• Finish 3rd cycle


 8
 9
 9
 11
 13
 18
 3
 10


Start	9	11	8	9	13	18	3	10	
1st cycle	9	11	8	9	13	18	3	10	0 swap, 1 comparison
2 nd cycle	8	9	11	9	13	18	3	10	2 swaps, 2 comparisons
3 rd cycle	8	9	9	11	13	18	3	10	1 swap, 2 comparisons
4 th cycle	8	9	9	11	13	18	3	10	0 swap, 1 comparisons
5 th cycle	8	9	9	11	13	18	3	10	0 swap, 1 comparisons
6 th cycle	3	8	9	9	11	13	18	10	6 swaps, 6 comparisons
7 th cycle	3	8	9	9	10	11	13	18	3 swaps, 4 comparisons
	We have	done!							
	3	8	9	9	10	11	13	18	12 swaps, 17 comparisons

Bubble Sort.

- Bubble sort uses the following strategy
- 1. Compare the first element with the next one.
- 2. If they are in the wrong order swap them.
- 3. Continue comparing until the end of the list.
- 4. Shorten the list by one the last element is now in the right place.
- 5. Repeat from step 1.







Start	9	11	8	9	13	18	3	10	
1 st cycle	9	8	9	11	13	3	10	18	4 swap, 7 comparisons
2 nd cycle	8	9	9	11	3	10	13	18	3 swaps, 6 comparisons
3 rd cycle	8	9	9	3	10	11	13	18	2 swap, 5 comparisons
4 th cycle	8	9	3	9	10	11	13	18	1 swap, 4 comparisons
5 th cycle	8	3	9	9	10	11	13	18	1 swap, 3 comparisons
6 th cycle	3	8	9	9	10	11	13	18	1 swap, 2 comparisons
	3	8	9	9	10	11	13	18	0 swap, 1 comparison
	We have o	done!							
	3	8	9	9	10	11	13	18	12 swaps, 28 comparisons

Summary

Start

• Finish

9	11	8	9	13	18	3	10
2	0	0	0	10	4.4	42	18
3	8	9	9	10	11	13	18

	Number of Swaps	Number of Comparisons	Total number of operations
Selection sort	7	28	35
Insertion sort	12	17	29
Bubble sort	12	28	39

Sorting Practice

Consider the following list of numbers:



- Sort it using:
 - 1. Selection sort
 - 2. Insertion sort
 - 3. Bubble sort
- Count the number of operations each time.
- Does this change the best sort?

Selection sort

Start 5 6 7 8 4 12 15 17	
1 st cycle 4 6 7 8 5 12 15 17 1 swap, 7 co	mparisons
2 nd cycle 4 5 7 8 6 12 15 17 1 swap, 6 co	mparisons
3 rd cycle 4 5 6 8 7 12 15 17 1 swap, 5 co	mparisons
4 th cycle 4 5 6 7 8 12 15 17 1 swap, 4 co	mparisons
5 th cycle 4 5 6 7 8 12 15 17 0 swap, 3 co	mparisons
6 th cycle 4 5 6 7 8 12 15 17 0 swap, 2 co	mparisons
7 th cycle 4 5 6 7 8 12 15 17 0 swap, 1 co	•
We have done!	·
4 5 6 7 8 12 15 17 4 swaps, 28 c	comparisons

Insertion sort

Start	5	6	7	8	4	12	15	17	
1 st cycle	5	6	7	8	4	12	15	17	0 swap, 1 comparison
2 nd cycle	5	6	7	8	4	12	15	17	0 swap, 1 comparison
3 rd cycle	5	6	7	8	4	12	15	17	0 swap, 1 comparison
4 th cycle	4	5	6	7	8	12	15	17	4 swaps, 4 comparisons
5 th cycle	4	5	6	7	8	12	15	17	0 swap, 1 comparison
6 th cycle	4	5	6	7	8	12	15	17	0 swap, 1 comparison
7 th cycle		5	6	7	8	12	15	17	0 swap, 1 comparison
Ve have do	ne!								
	4	5	6	7	8	12	15	17	4 swaps, 10 comparisons

Bubble sort

Start	5	6	7	8	4	12	15	17	
1 st cycle	5	6	7	4	8	12	15	17	1 swap, 7 comparisons
2 nd cycle	5	6	4	7	8	12	15	17	1 swap, 6 comparisons
3 rd cycle	5	4	6	7	8	12	15	17	1 swap, 5 comparisons
4 th cycle	4	5	6	7	8	12	15	17	1 swap, 4 comparisons
5 th cycle	4	5	6	7	8	12	15	17	0 swap, 3 comparisons



4 5 6	7	8	12	15	17	4 swaps, 25 comparisons
-------	---	---	----	----	----	-------------------------

Summary

Start

• Finish

5	6	7	8	4	12	15	17
4	5	6	7	8	12	15	17

	Number of Swaps	Number of Comparisons	Total number of operations
Selection sort	4	28	32
Insertion sort	4	10	14
Bubble sort	4	25	29

Sorting Practice

Consider the following list of numbers:

15 13 12	11 8	5	3	1
----------	------	---	---	---

- Sort it using:
 - 1. Selection sort
 - 2. Insertion sort
 - 3. Bubble sort
- Count the number of operations each time.
- Does this change the best sort?

Selection sort

Start	15	13	12	11	8	5	3	1	
1 st cycle	1	13	12	11	8	5	3	15	1 swap, 7 comparisons
2 nd cycle	1	3	12	11	8	5	13	15	1 swap, 6 comparisons
3 rd cycle	1	3	5	11	8	12	13	15	1 swap, 5 comparisons
4 th cycle	1	3	5	8	11	12	13	15	1 swap, 4 comparisons
5 th cycle	1	3	5	8	11	12	13	15	0 swap, 3 comparisons
6 th cycle	1	3	5	8	11	12	13	15	0 swap, 2 comparisons
7 th cycle	1	3	5	8	11	12	13	15	0 swap, 1 comparison
Ve have dor	ne!								
	1	3	5	8	11	12	13	15	4 swaps, 28 comparisons

Insertion sort

Start	15	13	12	11	8	5	3	1	
1 st cycle	13	15	12	11	8	5	3	1	1 swap, 1 comparison
2 nd cycle	12	13	15	11	8	5	3	1	2 swaps, 2 comparisons
3 rd cycle	11	12	13	15	8	5	3	1	3 swaps, 3 comparisons
4 th cycle	8	11	12	13	15	5	3	1	4 swaps, 4 comparisons
5 th cycle	5	8	11	12	13	15	3	1	5 swaps, 5 comparisons
6 th cycle	3	5	8	11	12	13	15	1	6 swaps, 6 comparisons
7 th cycle	1	3	5	8	11	12	13	15	7 swaps, 7 comparisons
We have dor	ne!								
	1	3	5	8	11	12	13	15	28 swaps, 28 comparisons

Bubble sort

Start	15	13	12	11	8	5	3	1	
1 st cycle	13	12	11	8	5	3	1	15	7 swaps, 7 comparisons
2 nd cycle	12	11	8	5	3	1	13	15	6 swaps, 6 comparisons
3 rd cycle	11	8	5	3	1	12	13	15	5 swaps, 5 comparisons
4 th cycle	8	5	3	1	11	12	13	15	4 swaps, 4 comparisons
5 th cycle	5	3	1	8	11	12	13	15	3 swaps, 3 comparisons
6 th cycle	3	1	5	8	11	12	13	15	2 swaps, 2 comparisons
7 th cycle	1	3	5	8	11	12	13	15	1 swap, 1 comparisons
We have don		2	-	o	11	12	12	15	20 20
	1	3	5	8	11	12	13	15	28 swaps, 28 comparisons

Summary

Start

• Finish

15	13	12	11	8	5	3	1
			8				

	Number of Swaps	Number of Comparisons	Total number of operations
Selection sort	4	28	32
Insertion sort	28	28	56
Bubble sort	28	28	56

Ball breaking.

- We have two bowling balls and a multi storey building.
- We know that if we drop a ball from a sufficiently high window it will break.
- Let us define a test as the process of dropping a ball from a specific floor.
- Our aim is to find out which is the lowest floor from which dropping a ball breaks it. Let's call this the critical floor.
- We also want to do this with the smallest number of tests.

The questions.

- What is the maximum number of tests that I need to conduct to find the critical floor in a 100 floor building?
- What order should I test the floors in?
- What is the highest floor you can reach with n tests?

A simpler case.

- Before thinking about the problem with two balls start with a single ball.
- What strategy makes sure that you exactly identify the critical floor?
- If I am allowed to conduct *n* tests what is the highest floor I can test and still guarantee that I find the critical floor.
- If my building is 100 floors high what is the worst case value for the number of tests?

- Drop the ball from each floor in turn: 1, 2, 3 ... until the ball breaks.
- If the critical floor is floor n you conduct n tests to find it.
- A 100 floor building will need at most 100 tests.

The real problem.

- Now that you have the answers for a single ball think about how you can make use of a second ball to increase the floor coverage for a given number of floors.
- We still have to exactly identify the critical floor.
- If we are allowed to perform at most *n* tests, what is the highest floor we can test and still be sure of finding the critical floor?
- Hint: there are two cases to consider:
 - Before the first ball breaks;
 - After the first ball breaks.

The strategy to follow is as follows:

- Test floor *n* with ball 1 (1 test)
- if the ball breaks test floors 1 up to n-1 with the second ball (n-1) tests for a total of n)
- if the ball survives we can use it to test floor 2n 1 (this equals n + (n 1) (2 tests)
- if it breaks we test floors n + 1 up to 2n 2 with the second ball (n 2) tests for a total of n
- if it survives use ball 1 to test floor 3*n* -3 (3 tests so far)
- If we repeat this process, reducing the added number of floors by one each time ball 1 survives we can test a total of n + (n 1) + (n 2) + ... + 2 + 1
- =n(n + 1)/2 floors with n tests.

- We need to solve n(n + 1)/2 > 100
- The smallest value of n which this is true for is 14
- Test floors in the order 14, 27, 39, 50, 60, 69, 77, 84, 90, 95, 99, 100.
- If the first ball breaks, test the floors between the last two floors tested in order.