

# The CNOT Gate 1: Elementary Properties

```
In[*]:= Quit[]
```


```
In[*]:= Let[Qubit, S]
```

---

## Elementary Properties

```
In[*]:= cnot = QuantumCircuit[CNOT[S[1], S[2]]]
```

Out[\*]=



```
In[*]:= Matrix[cnot] // MatrixForm
```

Out[\*]//MatrixForm=

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

```
In[*]:= in = Basis[S@{1, 2}];  
out = cnot ** in
```

Out[\*]=

$$\{ |0_{S_1}0_{S_2}\rangle, |0_{S_1}1_{S_2}\rangle, |1_{S_1}1_{S_2}\rangle, |1_{S_1}0_{S_2}\rangle \}$$

```
In[*]:= ProductForm[Thread[in → out], S@{1, 2}] // TableForm
```

Out[\*]//TableForm=


$$\begin{array}{l} |0\rangle \otimes |0\rangle \rightarrow |0\rangle \otimes |0\rangle \\ |0\rangle \otimes |1\rangle \rightarrow |0\rangle \otimes |1\rangle \\ |1\rangle \otimes |0\rangle \rightarrow |1\rangle \otimes |1\rangle \\ |1\rangle \otimes |1\rangle \rightarrow |1\rangle \otimes |0\rangle \end{array}$$

---

```
In[*]:= Let[Binary, x]
```

```
In[*]:= qc = QuantumCircuit[in = Ket[S@{1, 2} → x@{1, 2}], cnot]
```

Out[\*]=



```
In[*]:= out = Elaborate[qc]
```

```
Out[*]:=
```

$$|x_{1S_1} x_1 \oplus x_{2S_2}\rangle$$

```
In[*]:= ProductForm[in → out, S@{1, 2}]
```


```
Out[*]:=
```

$$|x_1\rangle \otimes |x_2\rangle \rightarrow |x_1\rangle \otimes |x_1 \oplus x_2\rangle$$

In summary,

```
In[*]:= qc = QuantumCircuit[in = Ket[S@{1, 2} → x@{1, 2}], cnot,
  Ket[S[1] → x[1], S[2] → Mod[x[1] + x[2], 2]],
  "PortSize" → {0.7, 1.5}]
```

```
Out[*]:=
```




## CNOT copies *basis* states

```
In[*]:= Let[Binary, x]
```

```
In[*]:= in = Ket[S@{1, 2} → {x, 0}, S@{1, 2}];
qc = QuantumCircuit[in, cnot]
```

```
Out[*]:=
```



```
In[*]:= out = Elaborate[qc]
```

```
Out[*]:=
```

$$|x_{S_1} x_{S_2}\rangle$$

```
In[*]:= ProductForm[in → out, S@{1, 2}]
```


```
Out[*]:=
```

$$|x\rangle \otimes |0\rangle \rightarrow |x\rangle \otimes |x\rangle$$

In summary,

```
In[*]:= qc = QuantumCircuit[in = Ket[S@{1, 2} → {x, 0}, S@{1, 2}], ,
  Ket[S[1] → x, S[2] → x]]
```

```
Out[*]:=
```



## No-Cloning Theorem

Quantum mechanics does not allow to copy an unknown state.


$$\left( \left| 0 \right\rangle_{c_0} + \left| 1 \right\rangle_{c_1} \right) \otimes \left| 0 \right\rangle \mapsto \left( \left| 0 \right\rangle_{c_0} + \left| 1 \right\rangle_{c_1} \right) \otimes \left( \left| 0 \right\rangle_{c_0} + \left| 1 \right\rangle_{c_1} \right) : \text{NOT ALLOWED!}$$

The result above was possible because the input state was one of the basis states.

## So, what happens to a superposition state?

```
In[*]:= qc1 = QuantumCircuit[Ket[S@{1, 2}], S[1, 6]]
```

Out[\*]=




```
In[*]:= out1 = Elaborate[qc1];
out1 // KetFactor
```

Out[\*]=

$$\frac{(\left| 0_{S_1} \right\rangle + \left| 1_{S_1} \right\rangle) \otimes \left| 0_{S_2} \right\rangle}{\sqrt{2}}$$

```
In[*]:= qc2 = QuantumCircuit[qc1, "Separator", CNOT[S[1], S[2]]]
```

Out[\*]=



```
In[*]:= out2 = Elaborate[qc2]
```

Out[\*]=

$$\frac{\left| 0_{S_1} 0_{S_2} \right\rangle}{\sqrt{2}} + \frac{\left| 1_{S_1} 1_{S_2} \right\rangle}{\sqrt{2}}$$

```
In[*]:= out1 → out2
```

Out[\*]=

$$\frac{\left| 0_{S_1} 0_{S_2} \right\rangle}{\sqrt{2}} + \frac{\left| 1_{S_1} 0_{S_2} \right\rangle}{\sqrt{2}} \rightarrow \frac{\left| 0_{S_1} 0_{S_2} \right\rangle}{\sqrt{2}} + \frac{\left| 1_{S_1} 1_{S_2} \right\rangle}{\sqrt{2}}$$

## Application: GHZ State

```

In[*]:= $n = 3;
        CC = S[{0}, $]
        TT = S[Range@$n, $]


Out[*]=
        {S0}

Out[*]=
        {S1, S2, S3}

In[*]:= cnot = Map[CNOT[CC, #] &, TT]
Out[*]=
        {CNOT[{S0} → {1}, {S1}], CNOT[{S0} → {1}, {S2}], CNOT[{S0} → {1}, {S3}] }

In[*]:= qc0 = QuantumCircuit[Ket[CC], Ket[TT], S[0, 6]]
Out[*]=

```



```

In[*]:= out0 = Elaborate[qc0];
        out0 // KetFactor


```

$$\text{Out[*]} = \frac{(|0_{S_0}\rangle + |1_{S_0}\rangle) \otimes |0_{S_1}0_{S_2}0_{S_3}\rangle}{\sqrt{2}}$$

```

In[*]:= qc = QuantumCircuit[qc0, "Separator", Sequence @@ cnot]
Out[*]=

```



```

In[*]:= out = Elaborate[qc]
Out[*]=

```

$$\frac{|0_{S_0}0_{S_1}0_{S_2}0_{S_3}\rangle}{\sqrt{2}} + \frac{|1_{S_0}1_{S_1}1_{S_2}1_{S_3}\rangle}{\sqrt{2}}$$

---

# Summary

## Functions

- CNOT
- Hadamard

## Related Links

- Chapters 2 of the Quantum Workbook (2022, 2023).
- Tutorial: “Quantum Computation: Overview”