

In [34]:

```
import pandas as pd
import scipy.stats as stats
import matplotlib.pyplot as plt
import numpy as np
```

In [4]:

```
# define matrix containing the values
rows=int(input("Enter number of rows"))
cols=int(input("Enter number of Columns"))
A=np.zeros((rows,cols))

# Enter values

for i in range(rows):
    for j in range(cols):
        A[i,j]=float(input(f"Enter the value of {i}th row and {j}th column"))

print("Matrix : ",A)
```

```
Matrix :  [[0.78 0.8  0.81 0.75 0.77 0.78]
 [0.85 0.85 0.92 0.86 0.81 0.83]
 [0.93 0.92 0.95 0.89 0.89 0.83]
 [1.14 0.97 0.98 0.88 0.86 0.83]
 [0.97 0.86 0.78 0.76 0.76 0.75]]
```

In [21]:

```
# Calculate the sum of rows
Sum_of_rows=np.sum(A,axis=1)
Sum_of_rows
```

Out[21]:

```
array([4.69, 5.12, 5.41, 5.66, 4.88])
```

In [22]:

```
# Calculate the sum of Columns
Sum_of_cols=np.sum(A,axis=0)
Sum_of_cols
```

Out[22]:

```
array([4.67, 4.4 , 4.44, 4.14, 4.09, 4.02])
```

In [23]:

```
Total_sum_of_elements=np.sum(Sum_of_cols,axis=0)
Total_sum_of_elements
```

Out[23]:

```
25.759999999999998
```

In [24]:

```
# Calculate Sum of Squares total
SST=0
for i in range(rows):
    for j in range(cols):
        SST+=A[i,j]**2

print(SST)
SST-=Total_sum_of_elements**2/(rows*cols)
print("Sum of total :",SST)
```

22.3416

Sum of total : 0.222346666666667025

In [25]:

```
# Calculation on treatments
SStreatments=0
print(Sum_of_rows.shape)
for i in range(Sum_of_rows.shape[0]):
    SStreatments+=Sum_of_rows[i]**2
SStreatments/=Sum_of_cols.shape[0]
SStreatments-=Total_sum_of_elements**2/(rows*cols)
print("Sum of Treatments :",SStreatments)
```

(5,)

Sum of Treatments : 0.102180000000000416

In [26]:

```
# Calculation of Block
SSblock=0
print(Sum_of_cols.shape)
for i in range(Sum_of_cols.shape[0]):
    SSblock+=Sum_of_cols[i]**2
SSblock/=Sum_of_rows.shape[0]
SSblock-=Total_sum_of_elements**2/(rows*cols)
print("Sum of Blocks :",SSblock)
```

(6,)

Sum of Blocks : 0.062866666666666818

In [27]:

```
# Error
SSerror=SST-SStreatments-SSblock
print("Error",SSerror)
```

Error 0.057299999999999791

In [28]:

```
# Calculation of degrees of Freedom
dft=Sum_of_rows.shape[0]-1
dfb=Sum_of_cols.shape[0]-1
dfw=dft*dfb
dftotal=rows*cols-1
```

In [29]:

```
# Calculation of Mean squares
mst=(SStreatments/dft)
msb=SSblock/dfb
mse=SSerror/dfw
```

In [30]:

```
Ftreat=mst/mse
Fblock=msb/mse
```

In [31]:

```
# create the ANOVA table
anova_table = pd.DataFrame({'Source of Variation': ['Between Treatments', 'Blocks', 'Error', 'Total'],
    'Sum of Squares': [SStreatments, SSblock, SSerror, SST],
    'Degrees of Freedom': [dft, dfb, dfw, dftotal],
    'Mean Square': [mst, msb, mse, np.nan],
    'F value': [Ftreat, Fblock, np.nan, np.nan],
})

# set the index to the source column
anova_table.set_index('Source of Variation', inplace=True)

# display the ANOVA table
print(anova_table)
```

	Sum of Squares	Degrees of Freedom	Mean Square	F value
Between Treatments	0.102180	4	0.025545	8.91
Blocks	0.062867	5	0.012573	4.38
Error	0.057300	20	0.002865	
Total	0.222347	29	NaN	


In [36]:

```
# create the residual plot
residuals = []
for i in range(rows):
    for j in range(cols):
        residuals.append(A[i,j]-Sum_of_rows[i]/Sum_of_cols.shape[0]) # Sum_of_rows[i]/
Sum_of_cols.shape[0]
#stats.probplot(residuals, plot=plt)
stats.probplot(residuals, dist="norm", plot=plt)

# Calculate normal scores

# set the title and labels
plt.title('Normal Probability Plot of Residuals')
plt.xlabel('Residuals')
plt.ylabel('Normal % probability')

# show the plot
plt.show()
```



## COMMENTS :

- As the value of the F value found (8.9162) is less than the value found from F from the table with degrees of freedom 4,20 which is (2.866) so we can say that the Nozzle design affects the shape factor. Thus, we rejectd the NULL Hypothesis and accept Alternate Hypothesis.
- The probability plot don't show any abnormalities in the data. Though some outliers can be seen at the ends.