

MAT315 Combinatorial Enumeration

Monsoon 2024

Instructor: Manjil Saikia

TA: Kanak Dhotre

MATHEMATICAL AND PHYSICAL SCIENCES DIVISION, SCHOOL OF ARTS AND
SCIENCES, AHMEDABAD UNIVERSITY, AHMEDABAD 380009, GUJARAT, INDIA
Email address: `manjil.saikia@ahduni.edu.in`

2020 *Mathematics Subject Classification.* Primary 05-01

Key words and phrases. Combinatorics, Discrete Mathematics.

Contents

Preface	vii
Chapter 1. What is Combinatorics?	1
1. Introduction	1
2. Counting Principles	8
3. The Pigeon-hole Principle	13
4. The Principle of Inclusion-Exclusion	15
Chapter 2. Generating Functions	19
1. Ordinary Generating Functions	19
2. Exponential Generating Functions	20
Chapter 3. The Art of Combinatorial Thinking	21
1. Binomial Coefficients	21
2. Catalan Numbers	23
Chapter 4. Partitions	29
1. Set Partitions	29
2. Integer Partitions	30
Chapter 5. Lattice Path Combinatorics	33
1. Dyck Paths Revisited	33
2. Motzkin and Schröder Paths	33
3. Non-intersecting lattice paths	33
4. q -Counting of lattice paths	33
Chapter 6. A Combinatorial Miscellany	35
1. Domino Tilings	35
2. Permutations	35
3. Symmetric Functions	35
4. Graphs and Trees	35

Preface

These are the lecture notes of MAT315 Combinatorial Enumeration, offered in Monsoon 2024 semester at Ahmedabad University, India. The notes were written down by the TA for the course, Kanak Dhotre and is as close to the classroom teaching as possible.

There are several very good textbooks in combinatorics, however the material that I wish to cover in this course is not available in a single source to my liking. So, I decided to make my own notes for this iteration, as well as for any future iterations of this course.

The background required to enroll for the course is very minimal, in fact, it is not even mandatory for a student to have done a first course in Discrete Mathematics. So, we introduce several basic concepts along the way and if there is a scope for some digression then we will take it. The course is supplemented by some homework assignments, some of the problems in the text were set in those assignments, some even appeared in the examinations.

I am thankful to Kanak Dhotre for typing these notes. Any errors that remain are mine. If there are any errors, comments, or corrections, please write to me via email.

Manjil Saikia

CHAPTER 1

What is Combinatorics?

1. Introduction

This course aims to delve into the study of discrete mathematical structures, a field which traces its roots back to the 1700s with the work of Leonhard Euler and has gained much attention between the 1960s and 1970s with the advent of computer science. Notably, Euler answered the following question posed by Philip Naude in the year 1741: “In how many ways can the number 50 be written as a sum of seven different positive integers?”. We shall understand the outline of Euler’s solution to the problem later in this course. A few important personalities (some of whose work we will study eventually) in the subject include Gian Carlo Rota, Donald Knuth, Richard Stanley, Srinivasa Ramanujan, and Pinagala.

Combinatorics is the science of patterns and arrangements. More concretely, it deals with the study of the existence and the number of arrangements possible for a given mathematical structure. We start our discussion with a few motivating questions which will make our statement clearer.

QUESTION 1.1. *In how many ways can you arrange the elements of the set $[n] := \{1, 2, 3, \dots, n\}$ such that the first entry in the arrangement is an even number?*

SOLUTION. Notice how when n is an even number we have $n/2$ choices of even numbers to make for the first entry in our arrangement. Once a choice for the said even number is made the remaining $n - 1$ choices can be made in $(n - 1)!$ ways. Hence, in the case where n is an even number we have $n/2 \cdot (n - 1)!$ possible arrangements. Can you see why we will have $(n - 1)/2 \cdot (n - 1)!$ arrangements for the case where n is an odd number?

QUESTION 1.2. *In how many ways can you arrange elements from the set $[n] := \{1, 2, 3, \dots, n\}$ on a grid with n columns and n rows?*

SOLUTION. Notice how for each one of the n^2 spaces we have n choices to make. Hence, there are a total of $\underbrace{n \cdots n}_{n^2 \text{ times}} = n^{n^2}$ possible arrangements.

QUESTION 1.3. *In how many ways can you arrange elements from the set $[n]$ (as defined in the previous two examples) on a grid with n columns and n rows such that each element appears atleast(/exactly) once in each row?*

SOLUTION. Since for each row in the grid we have $n!$ possible arrangements and the grid has n rows there are a total of $n \cdot n!$ possible arrangements.

QUESTION 1.4. *How many matrices of order $n \times n$ exist given that the entries must be from the set $\{0, 1\}$?*

SOLUTION. Since we have 2 choices for each one of the n^2 entries there are a total of 2^{n^2} such matrices.

QUESTION 1.5 (Permutation Matrices). *How many matrices of order $n \times n$ exist given that the entries must be from the set $\{0, 1\}$ and each row and column must have exactly one 1.*

SOLUTION. Notice how in the first row of our matrix we have n ways to fix the occurrence of 1. This forces $n - 1$ ways to fix the occurrence of 1 in the second row and so on. Hence, in all there are a total of $n \cdot (n - 1) \cdots 1 = n!$ such matrices.

REMARK 1.1. *Question 1.5 can also be restated as counting the number of order $n \times n$ matrices which have row-sum and column-sum equal to 1.*

With the following definition we shall now look at a generalization of sorts of the kind of matrices we were dealing with in Question 1.5.

DEFINITION 1.1 (Alternating Sign Matrix (ASM)). *A matrix of order $n \times n$ is called an alternating sign matrix if the following conditions hold:*

- (1) *All the entries of the matrix come from the set $\{-1, 0, 1\}$.*
- (2) *Each row-sum and column-sum is 1.*
- (3) *The non-zero entries (both row-wise and column-wise) alternate in sign.*

A result first proved by Doron Zeilberger in the year 1992 states that there are precisely

$$\prod_{k=0}^{n-1} \frac{(3k+1)!}{(n+k)!}$$

number of ASMs of order $n \times n$. A proof of this result is beyond the scope of these lectures and is mentioned only for the sake of completeness. We shall, however, count ASMs of order 3 now.

QUESTION 1.6. *How many ASMs of order 3 exist?*

SOLUTION. Notice how the set of matrices we counted in Question 1.5 are a subset of the set of ASMs of order n (verify each one of the three defining properties of an ASM). Next, we notice a pattern; an ASM (of any order) can't have a -1 in the first row. Why? To the contrary, assume there is an ASM with a -1 in the first row. Since the immediate non-zero entry below it must be a 1, the column sum cannot be 1 without violating the alternativity condition. A similar argument shows that ASMs cannot have a -1 in the last row, the first column, or the last column either. This pattern allows us to easily list all ASMs of order 3. First we list all the ASMs counted in Question 1.5:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

Now, the only ASM of order 3 with a negative entry is

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & -1 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

We shall now explore a combinatorial problem which involves counting the number of checkerboard tilings (also known as dimer models) which is of great interest to physicists.

QUESTION 1.7. *Consider an $m \times n$ board. In how many ways can you cover all the squares of the said board with no overlaps, no diagonal placements, and no protrusions off the board, using dominoes (blocks of order 2×1)?*

It is clear that the existence of atleast one tiling is guaranteed if and only if m and n are not simultaneously odd. However, to count the number of such tilings is not a trivial task. A result due to the famous Dutch physicist Pieter Kasteleyn in the 1960s states that the number of such tilings is

$$\prod_{v=1}^m \prod_{h=1}^n \left(4 \cos^2 \left(\frac{v\pi}{m+1} \right) + 4 \cos^2 \left(\frac{h\pi}{n+1} \right) \right)^{\frac{1}{4}}.$$

Once again, a proof of the result is beyond the scope of these lectures and is stated only for the sake of completeness.




FIGURE 1. 4 out of the 36 possible domino-tilings of a 4×4 board.

Next, we state an equivalent formulation of Question 1.7.

DEFINITION 1.2 (Graph). *A graph is a pair $G = (V, E)$ where V is a set whose elements are called vertices and E is a set of unordered pairs of vertices whose elements are called edges.*

DEFINITION 1.3 (Perfect Matching). *Let $G = (V, E)$ be a graph. An $M \subseteq E$ is called a perfect matching of G if no two edges in M share a common vertex and every vertex of G is incident to atleast one edge in M .*






FIGURE 2. The only possible perfect matchings of the grid graph of order 2


Now consider the following construction. To each square in a given a board of order $m \times n$, we assign a vertex. Additionally, there is an edge between the said vertices if and only if the corresponding squares are adjacent to one another. Figure 3 is an example of this construction.

FIGURE 3. A board of order 4×4 and it's corresponding graph

It is clear that our construction gives a one-to-one correspondence between the set of possible domino-tilings of a board and the set of perfect matchings of it's corresponding graph.

FIGURE 4. A domino tiling of a board of order 4×4 and the perfect matching that it admits

We shall revisit this formulation of the problem much later in the course. For now, restrict ourselves to a boards of order $2 \times n$ and count the number of possible domino tilings, say t_n as n varies over the set of natural numbers. It is not too difficult to convince yourself of the fact that $t_0 = 1$, $t_1 = 1$, $t_2 = 2$, $t_3 = 3$, and $t_4 = 5$. Since the first five terms in the sequence t_n are the same as the five terms of the Fibonacci sequence (say F_n), one might guess that $t_n = F_n$. This is indeed true, and we shall now give a proof of our claim using a counting argument.

FIGURE 5. Two ways to tile the first column of board of order $2 \times n$

PROOF. Notice how there are exactly two ways to tile the first column of any board of order $2 \times n$ (refer to Figure 5). It can either be tiled using a single vertical domino, in which case it remains to tile the sub-board of order $2 \times (n-1)$, or using two horizontal dominos, in which case it remains to tile the sub-board of order $2 \times (n-2)$. Since both of these cases are valid, it follows that

$$t_n = t_{n-1} + t_{n-2}.$$

Which is the same as the Fibonacci recurrence. \square

Now that we have $F_n = t_n$, we prove two standard Fibonacci identities using counting arguments similar to the ones used in the above proof.

CLAIM 1.1. $F_{m+n} = F_m F_n + F_{m-1} F_{n-1}$




FIGURE 6. The only 3 ways to tile the m -th row of a $2 \times (m+n)$ board.

PROOF. Notice how we have exactly 3 ways to tile the m -th row of a $2 \times (m+n)$ board (refer to Figure 6). Cases (a) and (b) account for when we have F_m ways to tile the sub-block of order $2 \times m$ **and** F_n ways to tile the sub-board which occurs immediately after. Case (c), on the other hand, accounts for when we have F_{m-1} ways to tile the sub-board of order $2 \times (m-1)$ which occurs before the already tiled m -th row **and** F_{n-1} ways to tile the sub-board of order $2 \times (n-1)$ which occurs after the already tiled m -th row. This proves the required identity. \square

CLAIM 1.2. $F_0 + \dots + F_n = F_{n+2} - 1$

PROOF. Notice how every board of order $2 \times k$ has a trivial tiling; the one which only uses vertical dominos and no horizontal ones. On a board of order $2 \times (n+2)$ if this trivial tiling is ignored, every other possible tiling must have the occurrence of atleast one pair of horizontal dominos. If the last such pair occurs at the $n+1$ -th column, the sub-board of order $2 \times n$ preceeding it can be tiled in F_n ways. If the last such pair occurs at the n -th column, the sub-board of order $2 \times (n-1)$ can be tiled in F_{n-1} ways, and so on. This gives us the required identity. \square




FIGURE 7. Examples of occurrences of the last pair of horizontal dominos in 3 non-trivial tilings of a board of order $2 \times (n+2)$

Recall how $\binom{n}{k}$ counts the number of ways one can choose k elements from a set of n elements. We shall now state a rather interesting re-interpretation of binomial coefficients.




FIGURE 8. 3 examples of lattice paths from $(0,0)$ to $(4,3)$

QUESTION 1.8. *Given that the only moves allowed are North (N), corresponding to $(i,j) \rightarrow (i,j+1)$, or East (E), corresponding to $(i,j) \rightarrow (i+1,j)$, count the number of paths from $(0,0)$ to (m,n) on a grid of order $m \times n$ (refer to Figure 8 for examples of such paths).*

SOLUTION. Notice how we require exactly m E-moves and n N-moves to reach (m,n) from $(0,0)$. However, every lattice path is determined completely by a choice of m E-moves (or equivalently n N-moves) which can be made in $\binom{m+n}{m}$ ways (or equivalently $\binom{m+n}{n}$ ways).

REMARK 1.2. *As a consequence of this counting exercise we have not only given a re-interpretation of binomial coefficients, but have also proved that they are symmetric, i.e., $\binom{n}{k} = \binom{n}{n-k}$.*

Per usual, we are now interested in giving a counting argument for a standard identity concerning binomial coefficients called Pascal's identity.

$$\text{CLAIM 1.3. } \binom{n}{k} + \binom{n}{k+1} = \binom{n+1}{k+1}$$

PROOF. Notice how there are $\binom{n+1}{k+1}$ choices of a lattice paths from $(0,0)$ to $(n-k, k+1)$. The last step in each one of these paths is either an N-move or an E-move. If it is an N-move then it suffices to choose one lattice path from $(0,0)$ to $(n-k-1, k+1)$ out of the

$$\binom{(n-k-1) + (k+1)}{k+1} = \binom{n}{k+1}$$

choices. Alternatively, if it is an E-move then it suffices to choose one lattice path from $(0,0)$ to $(n-k, k)$ out of the

$$\binom{(n-k) + k}{k} = \binom{n}{k}$$

choices. □

Let's remind ourselves that the central objective of this course is to count. From Question 1.1 through Question 1.6, we encountered examples where we derived exact closed-form expressions for counting. During our discussion of the dimer model,

we first stated a closed-form expression and then constructed a bijection between the tiling configuration and the perfect matchings configuration. By restricting ourselves to boards of size $2 \times n$, we demonstrated that finding a recurrence relation is also a valid counting technique. However, as we will see, the methods we have discussed so far are not always applicable. In such cases, we turn to the idea of generating functions, which we will introduce next.

DEFINITION 1.4 (Ordinary Generating Function). *Given a sequence $\{a_n\}_{n \geq 0}$ the formal power series, $\sum_{k=0}^{\infty} a_k x^k$, in an indeterminate x is called the generating function of $\{a_n\}_{n \geq 0}$.*

Since our definition works with a *formal* power series we need not worry about the divergence of the involved infinite sum. That being said, we can always be cautious and assume $|x| < 1$ to guarantee convergence.

QUESTION 1.9. *Find the generating function for the sequence of Fibonacci numbers.*

SOLUTION. Let $\{f_n\}_{n \geq 0}$ denote the sequence of Fibonacci numbers and let $F(x)$ be the corresponding generating function. Notice how

$$\begin{aligned}
 F(x) &= \sum_{k=0}^{\infty} f_k x^k \\
 &= f_0 + f_1 x + \sum_{k=2}^{\infty} f_k x^k \\
 &= 1 + x + \sum_{k=2}^{\infty} f_k x^k \\
 &= 1 + x + \sum_{k=2}^{\infty} (f_{k-1} + f_{k-2}) x^k \\
 &= 1 + x + \sum_{k=2}^{\infty} f_{k-1} x^k + \sum_{k=2}^{\infty} f_{k-2} x^k \\
 &= 1 + x + x(F(x) - 1) + x^2 F(x) \\
 &= 1 + xF(x) + x^2 F(x)
 \end{aligned}$$

Solving which we obtain

$$F(x) = \frac{1}{1 - x - x^2}$$

Throughout this course, we will explore how and why generating functions are extensively used in combinatorics. We introduce one such instance now, which will be discussed in greater depth later.

DEFINITION 1.5. *An integer partition of a natural number n is a non-increasing sequence of natural numbers, $\{\lambda_i\}_{i \geq 1}$, whose sum is n .*

QUESTION 1.10. *In how many ways can a natural number n be partitioned?*

Let $p(n)$ denote the number of partitions of n . Interestingly, no "nice" closed-form expression for $p(n)$ has been found. However, we will derive the generating function

$$\prod_{n=1}^{\infty} (1 + x^n + x^{2n} + x^{3n} + \cdots) = \sum_{n=0}^{\infty} p(n) x^n$$

which was given by Euler, later in the course.

REMARK 1.3. *In the year 1918, G.H Hardy and Srinivasa Ramanujan obtained an asymptotic expression (an expression which describes the limiting behaviour) for $p(n)$ which is given by*

$$p(n) \sim \frac{1}{4n\sqrt{3}} \exp\left(\pi\sqrt{\frac{2n}{3}}\right).$$

Oftentimes stating an asymptotic expression is also a valid counting technique.

2. Counting Principles

We state the fundamental counting principles now.

- (1) (Addition Principle): If $\{A_k\}_{k \geq 1}$ is a family of finite and pairwise disjoint sets then $|\bigcup_{k \geq 1} A_k| = \sum_{k \geq 1} |A_k|$.
- (2) (Subtraction Principle): If A and B are finite sets such that $B \subseteq A$ then $|A \setminus B| = |A| - |B|$.
- (3) (Product Rule): If A and B are finite sets, then $|A \times B| = |A| \cdot |B|$.
- (4) (Division Rule): For finite sets A and B if there exists a d -to-many function $f : A \rightarrow B$ then $|B| = |A|/d$.

In some sense we've already been using these principles in disguise thus far. For instance;

QUESTION 2.1. *How many k -digit positive numbers are there?*

SOLUTION. Since we have 10 choices for the first $(k-1)$ digits and 9 choices for the last digit, by the product rule there are $9 \cdot 10^{k-1}$ k digit numbers.

This is a good point to recall the binomial theorem and prove it using a counting argument instead of the standard induction argument.

THEOREM 2.1 (Binomial Theorem).

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k.$$

PROOF. Consider the expansion of

$$(x + y)^n = \underbrace{(x + y) \cdots (x + y)}_{n \text{ factors}}.$$

In this expansion, there are $\binom{n}{k}$ ways to choose x from exactly k of the n factors (which forces the choice of y from the remaining $(n-k)$ factors). Since k can range from 0 to n , applying the addition principle gives us the binomial theorem. \square

Now we state the multinomial theorem which generalizes the binomial theorem to expressions with more than two terms, allowing us to expand any power of a sum of multiple terms as a sum of products of those terms, each multiplied by a what is called a multinomial coefficient which is denoted and defined as

$$\binom{n}{k_1, \dots, k_m} := \frac{n!}{k_1! \dots k_m!}.$$

Next, we state an identity concerning multinomial coefficients which would help us prove the multinomial theorem.

CLAIM 2.1.

$$\binom{n}{k_1, \dots, k_r} = \binom{n}{k_1} \binom{n-k_1}{k_2} \dots \binom{n-k_1-\dots-k_{r-1}}{k_r}$$

PROOF. The proof follows simply by expanding the R.H.S out.

$$\begin{aligned} \binom{n}{k_1} \dots \binom{n-k_1-\dots-k_{r-1}}{k_r} &= \frac{n!}{(n-k_1)!k_1!} \frac{(n-k_1)!}{(n-k_1-k_2)!k_2!} \dots \frac{(n-k_1-\dots-k_{r-1})!}{(n-k_1-\dots-k_r)!k_r!} \\ &= \frac{n!}{k_1! \dots k_r!} \frac{(n-k_1)! \dots (n-k_1-\dots-k_{r-1})!}{(n-k_1)! \dots (n-k_1-\dots-k_{r-1})!} \\ &= \frac{n!}{k_1! \dots k_r!} \\ &= \binom{n}{k_1, \dots, k_r} \end{aligned}$$

□

We would expect the multinomial coefficient to count something. That is indeed the case, it counts the number of ways to arrange n distinct objects into m distinct bins such that the i -th bin contains k_i elements. With this interpretation of multinomial coefficients, and Claim 2.1, mimicing the steps involved in the proof of the binomial theorem we get the multinomial theorem which is stated below.

THEOREM 2.2 (Multinomial Theorem).

$$(x_1 + \dots + x_r)^n = \sum_{k_1 + \dots + k_r = n} \binom{n}{k_1, \dots, k_r} x_1^{k_1} \dots x_r^{k_r}.$$

It is natural to expect a Pascal-like identity for multinomial coefficients as well. In this spirit, we state the following claim.

CLAIM 2.2.

$$\binom{n}{k_1, \dots, k_r} = \binom{n-1}{k_1-1, k_2, \dots, k_r} + \dots + \binom{n-1}{k_1, k_2, \dots, k_r-1}.$$

REMARK 2.1. Notice how when $k_1 = k$, $k_2 = n - k$ and all the other k_i s are 0, we retrieve the Pascal's identity (Claim 1.3)

This is a good point to introduce ourselves to yet another generalization of the binomial theorem which deals with the expansion of $(x + y)^n$ where n is allowed to take complex values.

THEOREM 2.3 (Generalized Binomial Theorem). *For all $n \in \mathbb{C}$ we have*

$$(1+x)^n = \sum_{k \geq 0} \binom{n}{k} x^k$$

REMARK 2.2. *There are two well-known proofs of Theorem 2.3. One uses the idea of Taylor's series expansion, and the other uses the solution to a differential equation called the Legendre's equation. We state neither one of them here.*

In addition to the fundamental counting principles, there are a few more principles/techniques which we will use quite often in this course. One of them is the bijection principle. We state the principle formally and then give an example.

THEOREM 2.4 (Bijection Principle). *If S and T are finite sets then $|S| = |T|$ if and only if there exists a bijection $f : S \rightarrow T$.*

QUESTION 2.2. *Let S be a finite set. A **permutation** of S is a bijection from and to S . How many permutations of $[n] = \{1, 2, 3, \dots, n\}$ exist?*

We introduce the cycle notation before outlining a solution. In the case where $n = 1$, it is clear that there is only possible bijection, namely, $f_1 : \{1\} \rightarrow \{1\}$ such that $f_1(1) = 1$. We shall write f_1 as

$$\begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

In the case where $n = 2$, it is clear that there are two possible bijections. These are $g_1 : \{1, 2\} \rightarrow \{1, 2\}$ such that $g_1(1) = 1$, $g_1(2) = 2$, and $g_2 : \{1, 2\} \rightarrow \{1, 2\}$ such that $g_2(1) = 2$, $g_2(2) = 1$. We shall write g_1 and g_2 as

$$\begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$$

respectively. In essence, the first row in our notation lists the elements of the domain, and the second row lists where the bijection sends these elements. Do you see why in the case where $n = 3$ the bijections can be written as

$$\begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}.$$

Since the first row of any bijection in our notation is fixed and we have

$$n \cdot (n-1) \cdots 1 = n!$$

ways to fill out the second row, it is clear that there are $n!$ permutations of $[n]$. Recall from Question 1.5 that we also have exactly $n!$ permutation matrices. By the bijection principle we must have a bijection between the set of permutation matrices of order $n \times n$ and the set of permutations of $[n]$. The bijection is as follows. Corresponding to every permutation

$$\begin{pmatrix} 1 & 2 & \cdots & n \\ f(1) & f(2) & \cdots & f(n) \end{pmatrix}$$

of $[n]$ we assign an $n \times n$ matrix where the entry at $(i, f(i))$ for $1 \leq i \leq n$ is 1, and all other entries are 0. More explicitly,

$$\begin{aligned} \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} &\longleftrightarrow \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}, \\ \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix} &\longleftrightarrow \begin{pmatrix} 1 & & \\ & & 1 \\ & 1 & \end{pmatrix}, \\ &\vdots \\ \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} &\longleftrightarrow \begin{pmatrix} & & 1 \\ & 1 & \\ 1 & & \end{pmatrix}. \end{aligned}$$

We outline a proof of an identity involving binomial coefficients using the bijection principle now.

CLAIM 2.3.

$$\sum_{i=0}^n \binom{n}{i} = 2^n$$

PROOF. Let $S = \{a_1, \dots, a_n\}$ be a set with n elements. For each of the 2^n possible subsets $T = \{a'_1, \dots, a'_k\}$ of S (where k can be any value from 0 to n), there is a corresponding binary string of length n . In this string, the i th bit is 1 if the element a_i is in the subset T , and 0 if it is not. For instance in the case where $S = \{0, 1\}$ we have the following correspondence

$$\begin{aligned} \emptyset &\longleftrightarrow 00 \\ \{0\} &\longleftrightarrow 10 \\ \{1\} &\longleftrightarrow 01 \\ \{0, 1\} &\longleftrightarrow 11 \end{aligned}$$

□

Another counting technique which we've seen before in the proof of Pascal's identity (Claim 1.3) is called double counting. The broad idea is to count the same combinatorial object in two different ways. As an example, we give a proof of Claim 2.3 using the double counting technique.

PROOF. It is clear that 2^n counts the number of n -bit binary strings. To prove the claimed identity, we count the number of n -bit binary strings once again in a different way. Notice how an arbitrary n -bit binary string has $0 \leq i \leq n$ number of 0s. Since there are $\binom{n}{0}$ choices when $i = 0$, $\binom{n}{1}$ choices when $i = 1$, and so on, by the addition principle we have proved the required identity. □

We have all seen a proof by induction of the following identity. Now we give a proof using a double counting argument.

$$\text{CLAIM 2.4. } 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2} = \binom{n+1}{2}$$

PROOF. Consider the problem of counting the number of handshakes in a group of $n + 1$ people, where each person shakes hands with every other person. The first person in the group shakes hands with everyone except themselves; this gives n handshakes. The second person in the group shakes hands with everyone else; this gives $(n - 1)$ handshakes (not n because we've already counted the handshake between the first and second person). The third person in the group shakes hands with everyone; this gives $(n - 2)$, and so on. In all we have $n + (n - 1) + (n - 2) + \cdots + 1$ handshakes. Let's count again in a different way now. Notice how for each pair of people there is one handshake. Hence, there are $\binom{n+1}{2}$ handshakes in total. Putting the two observations together proves the claim. \square

Next, we introduce the "Stars-and-Bars" method. We shall outline how this method works with the help of a standard example.

QUESTION 2.3. For a given $n \in \mathbb{N}$ find the number of solutions to the equation $x_1 + \cdots + x_k = n$ where $n \in \mathbb{N}$ and $x_i \in \mathbb{Z}$?

SOLUTION. Recall how in the definition of partition (Definition 1.5), the order of the parts did not matter, i.e, the partitions $2 + 1$ and $1 + 2$ of 3 are the same to us. Partitions where the order does matter are called compositions. Compositions where 0s are allowed to be a part are called weak compositions. With these two definitions at hand, notice how there are two cases to deal with in Question 2.3.

- (1) All $x_i > 0$; which is equivalent to counting the number of compositions of n into k parts.
- (2) All $x_i \geq 0$; which is equivalent to counting the number of weak compositions of n into k parts.

Let $x_1 + \cdots + x_k = n$ be a composition (may or may not be weak) of n into k parts. Now we replace each x_i with x_i consecutive stars, and each $+$ with a bar. For instance, the composition $5 + 3 + 1$ would be written as

***** | *** | *

and the (weak) composition $5 + 3 + 1 + 0$ would be written as

***** | *** | * | |.

Since every composition (may or may not be weak) of n into k parts is completely determined by an arrangement of n stars and $k - 1$ bars, it is clear that we have $\binom{n+k-1}{k-1}$ weak compositions and $\binom{n-1}{k-1}$ compositions.

REMARK 2.3. Notice how every weak composition $n = d_1 + \cdots + d_k$ of n into k parts naturally corresponds to the monomial degree n in k variables given by $x_1^{d_1} \cdots x_k^{d_k}$.

We conclude this section by stating an informal technique called proof by pictures (or proof without words). This technique involves the use of visual representations to demonstrate the validity of a mathematical statement. Although in the strictest sense of things a proof by pictures is not a counting principle, it often helps develop intuition. For instance, we state such a proof for Claim 2.4.




FIGURE 9

PROOF. In Figure 9 notice how there are $1 + 2 + 3 + \cdots + n$ dots colored in red, $1 + 2 + 3 + \cdots + n$ dots colored in blue, and hence $2(1 + 2 + 3 + \cdots + n)$ total. It is also true that in our arrangement there are $n(n + 1)$ dots in total. Now the identity follows. \square

Yet another fundamental counting principle is the pigeon-hole principle. Given its importance, we will dedicate the upcoming section to explore it in detail.

REMARK 2.4. *We have seen how the principles we've discussed so far can be useful in proving combinatorial identities. Determining how to derive these identities is not always straightforward. However, a common technique involves running computer simulations to generate a sequence of values corresponding to the first few natural numbers. Tools like OEIS, RATE (in Mathematica), and Guess (in Sage-Math and Maple) can then be used to conjecture a closed form for the sequence.*

3. The Pigeon-hole Principle

We state what is known as the pigeon-hole principle.

THEOREM 3.1 (Pigeon-hole Principle). *Let A_1, \dots, A_k be pair-wise disjoint finite sets. If*

$$|A_1 \cup A_2 \cup \cdots \cup A_k| > kr$$

then there exists an $1 \leq i \leq k$ such that $|A_i| > r$.

Even though the statement sounds trivial, it has immense applications. We state a few examples now.

CLAIM 3.1. *Given $b \in \mathbb{Z}$, the decimal expansion of $1/b$ is either finite or eventually periodic.*

PROOF. Each step in the long division of 1 by b involves subtracting multiples of b from powers of 10 until the remainder is 0 or repeats itself. Since each one of these remainders must be strictly less than b , we have b possible remainders, namely, $0, 1, 2, \dots, b - 1$. If the remainder becomes 0 at some step, the division terminates, in which case the decimal expansion is finite. Next, consider the case where the division does not terminate. By the pigeon-hole principle, since there are

only b possible remainders and more than b steps, at least one of the remainders must repeat. This proves our claim because when a remainder repeats, the sequence of digits that follow in the decimal expansion must repeat as well. \square

CLAIM 3.2. *Let $S = \{1, 2, \dots, 2n\}$. Any $n + 1$ element subset K of S has two co-prime numbers, and two numbers such that one is an integer multiple of the other.*

PROOF. Consider the n -element subset $\{(1, 2), (3, 4), \dots, (2n - 1, 2)\}$ of $S \times S$ and notice how if we choose $n + 1$ numbers from $\{0, \dots, 2n\}$ then by the pigeon-hole principle we are guaranteed to pick two numbers from the same pair in $S \times S$. Since each element in the said subset is a pair of consecutive numbers, their gcd is necessarily 1. Next, notice how every element in the set $\{1, 2, \dots, 2n\}$ can be written in the form $2^a b$ where b is odd by factoring out as many 2s as possible. Since there are only n odd numbers in these set $\{1, 2, \dots, 2n\}$, we are left with n choices of b . Next, we construct b subsets of S , say $S_k \subseteq S$ such that $i \in S_k$ if and only if $i = 2^k b$. Finally, by the pigeon-hole principle, if one were to pick $n + 1$ elements from S at least 2 of them would be in the same S_k , i.e., there always exists a pair of numbers such that one divides the other. \square

CLAIM 3.3. *If S be an $n + 1$ -element set of natural numbers, then there exists a pair of numbers in S whose difference is divisible by n .*

PROOF. Notice how the remainder obtained upon dividing any element $a_i \in S = \{a_1, \dots, a_{n+1}\}$ by n must be one of $0, 1, \dots, n - 1$. Since there are $n + 1$ a_i s but only n possible remainders, by the pigeon-hole principle, at least two members (say a_i and a_j) in S must have the same remainder, i.e., $a_i = nq_1 + r$ and $a_j = nq_2 + r$ for some natural numbers q_1 and q_2 and $0 \leq r \leq n - 1$. Clearly, $a_i - a_j = n(q_1 - q_2)$ is divisible by n . \square

CLAIM 3.4. *Given m integers say a_1, \dots, a_m there exists k and l with $0 \leq k \leq l \leq m$ such that m divides $a_{k+1} + a_{k+2} + \dots + a_l$*

PROOF. Consider sums of the form $\sum_{i=1}^k a_i$ for $1 \leq k \leq m$. If one of these sums is divisible by m , i.e., their remainder upon division by m is 0 then we are done. If not, we have $m - 1$ possible remainders for the said sums, namely, $1, 2, \dots, m - 1$. Since there are m possible sums and only $m - 1$ possible remainders at least two of these sums share the same remainder. More specifically, for some $1 \leq r < s \leq m$ and integers q_1, q_2 we have $\sum_{i=1}^r a_i = nq_1 + r$ and $\sum_{i=1}^s a_i = nq_2 + r$. But then clearly, $\sum_{i=r+1}^s a_i = n(q_2 - q_1)$ is divisible by n and we are done. \square

QUESTION 3.1. *In a party of n people, where some (maybe all, maybe none) shake hands with one another, show that there are at least two people who shake the same number of hands.*

SOLUTION. Consider the case where each one of the n people shake hands at least once. The number of handshakes possible for everybody then is $n - 1$. Since there are n people in the party, by the pigeon-hole principle our claim holds true. If we don't assume that everyone shakes at least one hand then we can separate the n people between k people who don't shake any hands and m people who do, and the problem is reduced to the last case with $n = m$.

QUESTION 3.2. *In a group of 6 people where each pair of people are either friends or enemies, show that there are 3 people who are either all friends or all enemies.*

SOLUTION. Let A, B, C, D, E and F be a group of 6 friends. Pick A , and notice how each one of the remaining 5 people is either a friend of A or an enemy of A . Now, by the pigeon-hole at least three of them (say B, C and D) are either friends of A , or enemies of A . We consider the case where they are friends of A and the other case will follow similarly. If any one of the pairs B -and- C , C -and- D , or, B -and- D are friends, then we have found a group of 3 people who are all friends. If neither one of the pairs B -and- C , C -and- D , or, B -and- D are friends, then we have found a group of 3 people who are all enemies.

We state an interesting graph-theoretic (recall Definition 1.2) restatement of Question 3.2 now. Let the group of 6 be represented as vertices in a graph. For each pair of these vertices, let there be an edge connecting them, which is colored red if the two represented people are enemies and blue if they are friends. Now Question 3.2 asserts that now matter how you color the (edges of the) graph, there is always a “triangle” of the same color in the graph.




FIGURE 10. 3 randomly colored examples, each of which has either a red triangle or a blue triangle.

REMARK 3.1. *Figure 10 makes it clear that in some sense we are looking for appearance of a uniformly colored triangle sitting inside the (bigger) graph. Ramsey theory, a branch in the field of combinatorics, focuses precisely on questions of these kind.*

We conclude this section by stating two definitions which will turn out to be useful in the upcoming sections.

DEFINITION 3.1 (k -Permutation). *A k -permutation on $[n] = \{1, 2, \dots, n\}$ is a bijection between two k -element subsets of $[n]$.*

DEFINITION 3.2 (Circular k -Permutation). *k -permutations π and σ are called circular equivalents if there exists a shift $s \in [k]$ such that $i + s \equiv j \pmod{k}$.*

4. The Principle of Inclusion-Exclusion

[This section is a work in progress!!!] Note to scribe:

- (1) Finish the proof of Theorem 4.2

(2) Rewrite the proof of Claim 4.1

(3) Write and prove Claim 4.2

Recall that for finite sets A and B , the cardinality of their union is given by $|A \cup B| = |A| + |B| - |A \cap B|$. This formula, which adds (includes) the sizes of A and B while subtracting (excluding) the overcount in $|A \cap B|$ encapsulates the essence of the inclusion-exclusion principle. We start with a useful generalization of the motivating formula we just discussed.

THEOREM 4.1 (Inclusion-Exclusion Principle). *Given finite sets A_1, \dots, A_k ,*

$$\left| \bigcup_{i=1}^k A_i \right| = \sum_{n=1}^k (-1)^{n+1} \left(\sum_{1 \leq i_1 < \dots < i_n \leq k} |A_{i_1} \cap \dots \cap A_{i_n}| \right).$$

To see how useful this result is, we state two examples.

QUESTION 4.1. *How many of the first 1000 natural numbers are not divisible by 2, 3 or 6?*

SOLUTION. Out of the first 1000 natural numbers, notice how, there are $\lfloor 1000/n \rfloor$ numbers which are divisible by n . Now, by the inclusion exclusion principle we have

$$1000 - 500 - 333 - 200 + \lfloor 1000/6 \rfloor + \lfloor 1000/15 \rfloor + \lfloor 1000/10 \rfloor - \lfloor 1000/30 \rfloor = 266$$

natural numbers which are not divisible by 2, 3 or 6.

Question 4.1 motivates an equivalent version of Theorem 4.1 which we shall now state and prove.

THEOREM 4.2 (Inclusion-Exclusion Principle). *For $1 \leq i \leq n$, let A_i be subsets of a set X . Then the number of elements of X which do not lie in any of the A_i s is given by*

$$\sum_{I \subseteq \{1, \dots, n\}} (-1)^{|I|} |A_I|$$

where $A_I = A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}$ for $I = \{i_1, \dots, i_k\}$.

PROOF. □

Now we state a corollary of Theorem 4.2.

CLAIM 4.1. *The number of surjective mappings from $[n] = \{1, 2, 3, \dots, n\}$ to $[k] = \{1, 2, 3, \dots, k\}$ is given by*

$$\sum_{i=0}^k (-1)^i \binom{k}{i} (k-i)^n$$

PROOF. We mimic the setting of Theorem 4.2. Choose X to be the set of all mappings from $[n]$ to $[k]$, and A_i to be the set of all maps from $[n]$ to $[k]$ such that i is not in the range. Notice how $|X| = k^n$, $|A_i| = (k-1)^n$, and $|A_I| = |A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_j}| = (k-j)^n$. Now the claim follows. □

REMARK 4.1. *Since every surjection from $[n]$ to $[n]$ is necessarily a bijection, and by Question 1.5 we know that there are $n!$ bijections from $[n]$ to $[n]$, we've proved a combinatorial identity, namely that*

$$\sum_{i=0}^n \binom{n}{i} (n-i)^n = n!.$$

We now give an application of the inclusion-exclusion principle in number theory. More specifically, consider the Euler's φ function which is defined as

$$\begin{aligned}\varphi : \mathbb{N} &\rightarrow \mathbb{N} \\ n &\rightsquigarrow |\{x \in \mathbb{N} : x < n, \gcd(x, n) = 1\}|.\end{aligned}$$

For instance, $\varphi(1) = |\{1\}| = 1$, $\varphi(2) = |\{1\}| = 1$, $\varphi(3) = |\{1, 2\}| = 2$, and so on. In essence, $\varphi(n)$ counts the positive integers which are less n and are co-prime to n . It is clear that for a prime p , $\varphi(p) = p - 1$. More generally for a natural number $r > 0$ we also have

$$\varphi(p^r) = p^r - \left\lfloor \frac{p^r}{p} \right\rfloor = p^r - p^{r-1}.$$

THEOREM 4.3. For $n = p_1^{a_1} \cdots p_k^{a_k}$ we have

$$\varphi(n) = n \left(1 - \frac{1}{p_1}\right) \cdots \left(1 - \frac{1}{p_k}\right)$$

PROOF. We are interested in counting all the numbers a in the set $[n] = \{1, 2, 3, \dots, n\}$ is relatively prime to n . This happens if and only if a is not divisible by any of the primes p_1, \dots, p_k . Now, let A_i be the subset of $[n]$ that contains all numbers which are divisible p_i . By our setup it is clear that

$$\begin{aligned}\varphi(n) &= |[n] \setminus \bigcup_{i=1}^k A_i| \\ &= n - \left| \bigcup_{i=1}^k A_i \right| \\ &= n - \sum_{I \neq \emptyset \subseteq [k]} (-1)^{|I|+1} \left| \bigcap_{i \in I} A_i \right| \quad (\text{By Theorem 4.2})\end{aligned}$$

Notice how $\bigcap_{i \in I} A_i$ contains elements of $[n]$ which are not divisible by any p_i where $i \in I$, and hence also not divisible by $\prod_{i \in I} p_i$. It follows that

$$\begin{aligned}|\bigcap_{i \in I} A_i| &= \frac{n}{\prod_{i \in I} p_i} \\ \implies \varphi(n) &= n - \sum_{I \neq \emptyset \subseteq [k]} (-1)^{|I|+1} \frac{n}{\prod_{i \in I} p_i} \\ \implies \varphi(n) &= n \left(1 + \sum_{I \neq \emptyset \subseteq [k]} (-1)^{|I|} \frac{1}{\prod_{i \in I} p_i} \right) \\ \implies \varphi(n) &= n \left(1 + \sum_{I \neq \emptyset \subseteq [k]} \prod_{i \in I} \frac{-1}{p_i} \right) \\ \implies \varphi(n) &= n \prod_{i=1}^k \left(1 - \frac{1}{p_i} \right)\end{aligned}$$

□

Now that we have counted the number of elements which are co-prime to n , we are interested in finding their sum. In this spirit consider the following claim.

CLAIM 4.2.

We conclude this section by introducing derangements.

QUESTION 4.2. *Suppose n people go a party and each person is wearing a different hat. In how many ways can the n people return from the party such that nobody is wearing their original hat?*

SOLUTION. It is clear that we are counting the number of permutations on $[n]$ with no fixed points. Let A_j be the set of permutations which consist of all permutations such that j is fixed point. Now the inclusion-exclusion principle the number of permutations with no fixed points are

$$D_n := \left| \underbrace{A_1^c}_{1 \text{ is not fixed}} \cup \underbrace{A_2^c}_{2 \text{ is not fixed}} \cup \underbrace{A_3^c}_{3 \text{ is not fixed}} \cup \cdots \cup \underbrace{A_n^c}_{n \text{ is not fixed}} \right|.$$

Since it is clear that $|A_j| = (n-1)!$, we have

$$|A_{i_1} \cup A_{i_2} \cup \cdots \cup A_{i_k}| = (n-k)!.$$

Once again by the inclusion-exclusion principle

$$\begin{aligned} D_n &= n! - \binom{n}{1} (n-1)! + \binom{n}{2} (n-2)! - \cdots + (-1)^n \binom{n}{n} 0! \\ &= n! - \frac{n!}{1!} + \frac{n!}{2!} - \cdots + (-1)^n \frac{n!}{n!} \end{aligned}$$

CHAPTER 2

Generating Functions

This chapter builds on our earlier discussion of generating functions from Chapter 1 (see Definition 1.4). Among the various types used in combinatorics and other fields like number theory, we will focus on two: ordinary generating functions and exponential generating functions

1. Ordinary Generating Functions

We start with a few examples.

EXAMPLE 1.1. *Corresponding to the constant sequence of 1s, the generating function is $1 + x + x^2 + x^3 + \cdots = \frac{1}{1-x}$*

EXAMPLE 1.2. *Corresponding to the sequence $1, 2, 3, \dots$, the generating function is $1 + 2x + 3x^2 + 4x^3 + \cdots = 1/(1-x)^2$. Notice how equality follows from the fact that $1 + 2x + 3x^2 + 4x^3 + \cdots$ is the formal derivative of the generating function we obtained in the previous example.*

More often than not generating functions are used to solve recurrences. For instance consider the following question.

QUESTION 1.1. *Find a closed form expression for the recurrence given by $a_{n+1} = 2a_n + 1$ where $a_0 = 0$.*

SOLUTION. Let $A(x) = a_0 + a_1x + a_2x^2 + \cdots$ be the generating function corresponding to the sequence. Notice how

$$\begin{aligned} \sum_{n \geq 0} a_{n+1}x^n &= 2 \sum_{n \geq 0} a_nx^n + \sum_{n \geq 0} x^n \\ \implies \frac{A(x)}{x} &= 2A(x) + \frac{1}{1-x} \\ \implies A(x) &= \frac{x}{(1-x)(1-2x)} \\ \implies A(x) &= \frac{1}{1-2x} - \frac{1}{1-x} \\ \implies A(x) &= (1 + 2x + 4x^2 + 8x^3 + \cdots) - (1 + x + x^2 + x^3 + \cdots) \\ \implies A(x) &= x + 3x^2 + 7x^3 + \cdots \end{aligned}$$

Now a_n is just the coefficient of x^n in $A(x)$.

We state one more example. Recall how with Question 1.9, we found the generating function for the sequence of Fibonacci numbers. We are now interested in finding a closed form of numbers in this sequence.

SOLUTION. Let $r_1, r_2 = (-1 \pm \sqrt{5})/2$ be the roots of the polynomial $1 - x - x^2$ and notice how

$$\begin{aligned}
 F(x) &= \frac{1}{1 - x - x^2} \\
 &= \frac{1}{(r_1 - x)(r_2 - x)} \\
 &= \frac{1}{(r_1 - x)(r_2 - r_1)} + \frac{1}{(r_2 - x)(r_1 - r_2)} \\
 &= \frac{1}{\sqrt{5}} \left(\frac{1}{r_2 - x} - \frac{1}{r_1 - x} \right) \\
 &= \frac{1}{\sqrt{5}} \left(\frac{1/r_2}{1 - (x/r_2)} - \frac{1/r_1}{1 - (x/r_1)} \right) \\
 &= \frac{1}{\sqrt{5}} \left(\frac{1}{r_2} \left(1 + \frac{x}{r_2} + \frac{x^2}{r_2^2} + \cdots \right) - \frac{1}{r_1} \left(1 + \frac{x}{r_1} + \frac{x^2}{r_1^2} + \cdots \right) \right) \\
 &= \frac{1}{\sqrt{5}} \left(\frac{1}{r_2} + \frac{x}{r_2^2} + \frac{x^2}{r_2^3} + \cdots - \frac{1}{r_1} - \frac{x}{r_1^2} - \frac{x^2}{r_1^3} - \cdots \right) \\
 &= \frac{1}{\sqrt{5}} \left(\left(\frac{1}{r_2} - \frac{1}{r_1} \right) + \left(\frac{1}{r_2^2} - \frac{1}{r_1^2} \right) x + \left(\frac{1}{r_2^3} - \frac{1}{r_1^3} \right) x^2 + \cdots \right)
 \end{aligned}$$

Now, the n -th Fibonacci number is just the coefficient of x^n in $F(x)$.

Often-times we are interested in computing the product of two generating functions. To this end, consider the following result due to Cauchy.

CLAIM 1.1 (Cauchy Product). *Let $A(x) = \sum_{n \geq 0} a_n x^n$ and $B(x) = \sum_{n \geq 0} b_n x^n$ be two ordinary generating functions. Their product $C(x)$, is then given by $A(x)B(x) = \sum_{n \geq 0} c_n x^n$ where*

$$c_n = \sum_{k=0}^n a_k b_{n-k}$$

2. Exponential Generating Functions

CHAPTER 3

The Art of Combinatorial Thinking

1. Binomial Coefficients

This section aims to push-forward our familiarity with binomial coefficients which we have gained from the previous chapter. We do so by outlining proofs of a few standard combinatorial identities.

CLAIM 1.1.

$$\sum_{k=1}^n k \binom{n}{k}^2 = n \binom{2n-1}{n-1}$$

PROOF. Let J be a collection of $2n$ objects which is partitioned into 2 equal sub-collections J_1 and J_2 . Now, we want to make a choice of n elements from J which includes a distinguished element, say x from J_1 (or equivalently J_2). Since there are n ways to choose the said distinguished element, and $\binom{2n-1}{n-1}$ ways to choose the rest of the $n-1$ elements, in all we have

$$n \binom{2n-1}{n-1}$$

ways to make the said choice. Alternatively, we can also choose k elements from J_1 in $\binom{n}{k}$ ways, $n-k$ elements from J_2 in $\binom{n}{n-k}$ ways, and then choose the distinguished element from J_1 (or equivalently J_2) in $1 \leq k \leq n$ ways. Thus far, we have proved

$$\sum_{k=1}^n k \binom{n}{k} \binom{n}{n-k} = n \binom{2n-1}{n-1}.$$

Since by Remark 1.2

$$\binom{n}{k} = \binom{n}{n-k},$$

we are done. □

CLAIM 1.2.

$$\sum_{k=1}^n k \binom{n}{k} = n2^{n-1}$$

PROOF. In a class of n students, a football coach wants to choose students to form a football team of $1 \leq k \leq n$ players. Additionally, the coach also wants one of k students to be the captain of the team. Since the representative can be chosen in n ways and the remaining set of students can be chosen in 2^{n-1} ways, the team can be chosen in $n2^{n-1}$ ways. Alternatively, the coach can select k players out of

n in $\binom{n}{k}$ ways, and then choose the captain in k ways. Since $0 < k \leq n$, we are done. \square

CLAIM 1.3.

$$\sum_{i=1}^n i(n-1) = \sum_{i=1}^n \binom{i}{2} = \sum_{i=0}^{n-2} \binom{n-i}{2} = \binom{n+1}{3}$$

PROOF. It is clear that $\binom{n+1}{3}$ is the number of ways to choose 3 numbers, say $a_1 < a_2 < a_3$, from the $n+1$ -element set $\{a_1, \dots, a_{n+1}\}$. Now we argue based on which one of these three is chosen first. \square

CLAIM 1.4. For all $n \geq r \geq 0$ we have

$$\binom{r}{r} + \binom{r+1}{r} + \dots + \binom{n}{r} = \binom{n+1}{r+1}.$$

PROOF. We give a double counting argument. Notice how $\binom{n+1}{r+1}$ counts the number of ways to choose $(r+1)$ -element subsets of $[n+1] = \{1, \dots, n+1\}$. A choice of the said subsets can also be made by first fixing the largest of the $r+1$ numbers. This is given by $\binom{r}{r} + \dots + \binom{n}{r}$. \square

CLAIM 1.5. $\binom{2n}{n} < 4^n$

PROOF. The claim follows from the simple observation that

$$4^n = (1+1)^{2n} = \sum_{i=0}^{2n} \binom{2n}{i} = \binom{2n}{n} + \underbrace{\sum_{\substack{0 \leq i \leq 2n, i \neq n \\ > 0}} \binom{2n}{i}}_{>0}.$$

\square

CLAIM 1.6 (Chu-Vandermonde Identity).

$$\binom{2n}{n} = \sum_{k=0}^n \binom{n}{k}^2$$

PROOF. Once again, by Remark 1.2, notice how

$$\sum_{k=1}^n \binom{n}{k}^2 = \sum_{k=1}^n \binom{n}{k} \binom{n}{n-k}.$$

Now we argue combinatorially. Let J be a collection of $2n$ objects which is partitioned into 2 equal sub-collections J_1 and J_2 . Any choice of n elements from J involves choosing k elements from J_1 and $n-k$ elements from J_2 where $0 \leq k \leq n$. \square

CLAIM 1.7. For all $n > 0$.

$$\sum_{k=0}^n 2^k \binom{n}{k} = 3^n$$

PROOF. We give a double counting argument. Notice how 3^n is the number of n -bit ternary strings. Each such string will have $n - k$ -many 0s in it where $0 \leq k \leq n$. Such a string can be chosen in $\binom{n}{n-k} = \binom{n}{k}$ ways, and the remaining k bits (which can now only either be 1s or 2s) can be chosen in 2^k ways. This completes the proof. \square

2. Catalan Numbers

[This section is a work in progress!!!] Note to scribe:

- (1) Add more details to the proof of Claim 2.1.
- (2) Explain what rooted binary trees are before counting them.
- (3) Finish proof of Theorem 2.2.

This section introduces a set of problems which are seemingly distinct but share a common underlying sequence of numbers called the Catalan numbers. We shall show this commonality by setting up appropriate bijections between the said problems.

QUESTION 2.1. Recall how with Question 1.8 we counted the number of lattice paths from $(0,0)$ to (m,n) . Count the number of lattice paths, $a_1(n)$, from $(0,0)$ to (n,n) which never go below the $x = y$ diagonal.




FIGURE 1. All possible lattice paths which don't go below the diagonal for cases $n = 1, 2, 3$

REMARK 2.1. Notice that reflecting each path in Figure 1 across the main diagonal produces a lattice path that remains strictly below it. This bijection clearly shows that $a_1(n)$, the number of lattice paths that do not fall below the diagonal is the same as $a_2(n)$, the number of those that do not rise above it.

QUESTION 2.2. Count the number of ways $a_3(n)$, of filling a $2 \times n$ grid with elements from the set $\{0, 1, 2, 3, \dots, 2n\}$ such that all elements are unique, increasing row-wise, and decreasing column-wise.

1	
2	

1	2
3	4

1	3
2	4

1	2	3
4	5	6

1	2	4
3	5	6

1	3	4
2	5	6

1	2	5
3	4	6

1	3	5
2	4	6

FIGURE 2. All possible such arrangements of order $2 \times n$ for cases $n = 1, 2, 3$

SOLUTION. A bijection presents itself when Figure 1 and Figure 2 are compared. Namely, the entries in the first row of the $2 \times n$ grid correspond to when an N -step in our lattice path occurs, and the entries in the second row correspond to when an E -step in our lattice path occurs.

QUESTION 2.3. Count the number of Dyck paths $a_4(n)$, from $(0, 0)$ to $(2n, 0)$. Where, by a Dyck path we refer to the path admitted by a sequence of up-moves, corresponding to $(i, j) \rightarrow (i + 1, j + 1)$ and down-moves, corresponding to $(i, j) \rightarrow (i + 1, j - 1)$, which does not go below the x -axis.





FIGURE 3. All possible Dyck paths from $(0, 0) \rightarrow (2n, 0)$ for $n = 1, 2, 3$

SOLUTION. Once again, a bijection presents itself when Figure 1 and Figure 3 are compared. Namely, the entries in the first row of the $2 \times n$ grid correspond to when an up-step in our Dyck path occurs, and the entries in the second row correspond to when a down-step in our Dyck path occurs.

QUESTION 2.4. Count the number of ways $a_5(n)$, of joining n non-intersecting chords on a circle marked with $2n$ points.

FIGURE 4. All possible such configurations for cases $n = 1, 2, 3$

SOLUTION. Once again, a bijection presents itself when Figure 2 and Figure 4 are compared. Namely, if p_1, \dots, p_{2n} are the marked points on the circle, then we join p_i and p_j if and only if i occurs below j in the arranged grid of numbers.

QUESTION 2.5. Count $a_6(n)$, the number of legal sequences of $2n$ parentheses. Where, by a legal sequence of parentheses we mean one in which the parentheses can be properly matched, i.e., each opening parenthesis should be matched to a closing one that lies further to its right. We count these for $n = 1, 2, 3$.

$$\begin{aligned} n = 1 : & \quad () \\ n = 2 : & \quad (()), ()() \\ n = 3 : & \quad ((())), ()(()), (())(), ()()() \end{aligned}$$

Once again, notice how corresponding to each grid in Figure 2 there is a legal sequence of parentheses. Namely, for each entry in the first row we open a parenthesis, and close it for each entry in the second row.

QUESTION 2.6. Count $a_7(n)$, the number of non-crossing partitions on the set $[2n] := \{1, 2, 3, \dots, 2n\}$. Where by a non-crossing partition on $[2n]$ we refer to an arrangement of $2n$ points on a line, with n non-intersecting arcs joining them.




FIGURE 5. All possible non-crossing partitions on $[2n]$ for $n = 1, 2, 3$

SOLUTION. Once again, a bijection presents itself when Figure 2 and Figure 5 are compared. Namely, if p_1, \dots, p_{2n} are the marked points on the line, then we join p_i and p_j if and only if i occurs below j in the arranged grid of numbers.

To summarize, by the many bijections we have set up, $C(n) := a_1(n) = a_2(n) = \dots = a_7(n)$ for all $n \geq 0$. Additionally, we have also seen that $A(n)$ takes values 1, 1, 2 and 5 for when $n = 0, 1, 2$ and 3 respectively. Motivated readers may check why and how $C(4) = 14$, $C(5) = 42$, and so on, but this is not how we want to proceed. Consider the following definitions.

DEFINITION 2.1 (Triangulation). *A triangulation of a convex polygon with $n+2$ vertices \mathcal{P}_{n+2} , is a set of $n-1$ diagonals which do not cross each other in the interior of \mathcal{P}_{n+2} .*

REMARK 2.2. *For instance, the 3-gon (triangle) has no triangulations, but the 4-gon (square/rectangle) has 2.*

DEFINITION 2.2 (Catalan Numbers). *The sequence $C(n)$ which counts the the number of triangulations of \mathcal{P}_{n+2} is called the sequence of Catalan numbers.*

THEOREM 2.1. *For all $n \geq 0$ we have*

$$C(n+1) = \sum_{k=0}^n C(k)C(n-k)$$




FIGURE 6

PROOF. Let \mathcal{P}_{n+3} be an $n + 3$ convex polygon with vertices A_1, \dots, A_{n+3} . Next, pick an arbitrary vertex A_k and consider the triangle $\Delta A_1 A_{n+3} A_k$. See Figure 6 and notice how this choice splits \mathcal{P}_{n+3} into two convex polygons. One with k vertices (marked red) and the one with $n - k + 4$ vertices (marked blue). Since k is allowed to vary from 2 to $n + 2$, by Definition 2.2 we have $\sum_{k=2}^{n+2} C(k-2)C(n-k+2)$ ways to triangulate \mathcal{P}_{n+3} . Finally, shifting the index of summation by 2 grants the required result. \square

CLAIM 2.1.

$$C(n) = \frac{1}{n+1} \binom{2n}{n}$$

PROOF. Let

$$f(x) = \sum_{n=0}^{\infty} C(n)x^n$$

be the generating function corresponding to the Catalan numbers. By Theorem 2.1,

$$\begin{aligned} f(x)^2 &= C(0)^2 + (C(0)C(1) + C(1)C(0))x + \dots + (C(0)C(n) + C(1)C(n-1) + \dots + C(n)C(0))x^n + \dots \\ &= C(1) + C(2)x + \dots + C(n+1)x^n + \dots \\ &= \frac{C(x) - C(0)}{x}. \end{aligned}$$

Solving for $C(x)$ gives

$$C(x) = \frac{1 \pm \sqrt{1-4x}}{2x}.$$

Since $C(n)$ is always positive, we discard

$$C(x) = \frac{1 + \sqrt{1-4x}}{2x}.$$

and expand $\sqrt{1-4x}$ using Theorem 2.3 to see why

$$C(x) = \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{2n}{n} x^n.$$

\square

REMARK 2.3. *Catalan numbers come up in a huge class of counting problems. In fact, Richard Stanley, in his book, “Catalan Numbers” outlines 214 such problems.*

QUESTION 2.7. *Count the number of rooted binary trees*

We are now interested in stating a recurrence for Catalan numbers which is different from the one we already stated. To this end consider the following theorem and its corollary.

THEOREM 2.2. *Let T_n be the number of triangulations of an n -gon, then for $n \geq 4$,*

$$(2n - 6)T_n = n \sum_{k=3}^{n-1} T_k T_{n-k+2}$$

PROOF.

□

COROLLARY 2.1.

$$(2n - 2)C_n = (n + 2) \sum_{k=1}^{n-1} C_k C_{n-k}$$

CHAPTER 4

Partitions

A first course in combinatorics typically focuses on two types of partitions: set partitions and integer partitions. We will begin with a brief discussion of set partitions, followed by a more in-depth exploration of integer partitions.

1. Set Partitions

[This section is a work in progress!!!] Note to scribe:

(1) Finish proofs of Claim 1.1-1.6.

Notice how there are 6 ways to partition the set $\{1, 2, 3, 4\}$ into 3 blocks. These are,

- (1) $[1], [2], [3, 4]$
- (2) $[1], [2, 3], [4]$
- (3) $[1, 2], [3], [4]$
- (4) $[1, 4], [2], [3]$
- (5) $[1, 3], [2], [4]$
- (6) $[2, 4], [1], [3]$

This kind of counting is generalized by what are called a Stirling numbers of the 2nd kind. More formally,

DEFINITION 1.1. A set partition of a finite set B into k “blocks” is a collection of k subsets of B say B_1, \dots, B_k such that

- (1) $\bigcup_{i=1}^k B_i = B$,
- (2) $B_i \cap B_j = \emptyset$ for all $i \neq j$,
- (3) and none of the B_i ’s are empty.

DEFINITION 1.2. If $B = [n] = \{1, \dots, n\}$ then a Stirling number of the 2nd kind $S(n, k)$, is the number of set partitions of B into k blocks.

We take $S(0, 0)$ to be 1 by convention. Additionally, the fact that $S(n, n) = 1$, $S(n, 0) = 0$, and $S(n, 1) = 1$ follow immediately. Per usual we state a few identities concerning these numbers.

CLAIM 1.1. For all $n, k \geq 0$, with $n \geq k$ we have $S(n, k) = S(n-1, k-1) + kS(n-1, k)$

CLAIM 1.2. For all $n, k \geq 0$, with $n \geq k$ we have $S(n+1, k) = \sum_{i=0}^n S(n-i, k-1)$

CLAIM 1.3. $S(n, 2) = 2^{n-1} - 1$

CLAIM 1.4. $S(n, n-1) = \binom{n}{2}$

CLAIM 1.5.

$$S(n, k) = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$$

DEFINITION 1.3 (Bell Numbers). *The number of all partitions of $[n]$ is called a bell number and is denoted by $B(n)$. More specifically,*

$$B(n) = \sum_{k=1}^n S(n, k)$$

We are interested in coming up with a recurrence of $B(n)$ which is independent of any $S(n, k)$ s.

CLAIM 1.6.

$$B(n+1) = \sum_{k=0}^n \binom{n}{n-k} B(k)$$

2. Integer Partitions

Recall how with Definition 1.5 we defined integer partitions. The study of these partitions dates back to the work of Leonard Euler in the 18th century, who introduced generating functions to analyze them. Their study was then developed further through the work of mathematicians like Srinivasa Ramanujan and Major MacMachon, who revealed deep arithmetic and combinatorial properties. For instance, if $p(n)$ denotes the number of partitions of n , then the celebrated Ramanujan congruences state that

$$P(5n+4) \equiv 0 \pmod{5}$$

$$P(7n+5) \equiv 0 \pmod{7}$$

$$P(11n+6) \equiv 0 \pmod{11}$$

In the absence of a closed form (recall Remark 1.3), we are interested in finding the generating function of $p(n)$.

CLAIM 2.1.

$$\prod_{n \geq 1} (1 + q^n + q^{2n} + \cdots) = \sum_{n \geq 0} p(n) q^n.$$

PROOF. We can expand the product on the L.H.S,

$$(1 + q + q^2 + q^3 + \cdots)(1 + q^2 + q^4 + q^6 + \cdots)(1 + q^3 + q^6 + q^9 + \cdots) \cdots,$$

out by choosing one term from each factor in all possible ways. If we then collect like terms, the coefficient of q^k will be the number of ways to choose one term from each factor so that the exponents of the said terms sum up to k . This is also what the R.H.S counts. For instance q^3 can be obtained in the following ways

- (1) Choose q^3 from the first bracket and 1 from every other bracket. This corresponds to the partition $3 = 1 + 1 + 1$.
- (2) Choose q from the first bracket, q^2 from the second bracket, and 1 from every other bracket. This corresponds to the partition $3 = 2 + 1$.
- (3) Chose 1 from the first two brackets and q^3 from the third bracket. This corresponds to the partition $3 = 3$.

□

Next, we are interested in looking at a two refinements of partitions which follow quite naturally by the arguments we used in the previous proof.

CLAIM 2.2. *Let $p_E(n)$ and $p_O(n)$ denote the number of partitions of n into even and odd parts respectively. Then,*

$$\prod_{n=1,3,5,\dots} (1 + q^n + q^{2n} + \dots) = \sum_{n \geq 0} p_O(n) q^n$$

$$\prod_{n=2,4,6,\dots} (1 + q^n + q^{2n} + \dots) = \sum_{n \geq 0} p_E(n) q^n$$

CHAPTER 5

Lattice Path Combinatorics

1. Dyck Paths Revisited
2. Motzkin and Schröder Paths
3. Non-intersecting lattice paths
4. q -Counting of lattice paths

CHAPTER 6

A Combinatorial Miscellany

1. Domino Tilings
2. Permutations
3. Symmetric Functions
4. Graphs and Trees

