

# Applications of MoS<sub>2</sub> as a Two-Dimensional Material Beyond Graphene

Kraig Andrews\*

Wayne State University Department of Physics and Astronomy

(Dated: March 23, 2015)

An article usually includes an abstract, a concise summary of the work covered at length in the main body of the article.

## I. INTRODUCTION

The study of nanomaterials is wide-reaching and overlaps many different disciplines of scientific research, from physics to chemistry, materials science, and biomedical engineering, for example. In the past decade research focusing on two-dimensional materials has rapidly increased compared to previous decades. This swift jump in the amount literature and studies being done on various two-dimensional materials is due to several breakthroughs which occurred in the mid 2000s, the resulting properties of such materials, and their possible and nu-

merous applications which are many sweeping across a wide-range of disciplines both academic and commercial. The major breakthrough that spurred the plethora of research and interest that now exists on two-dimensional materials and its uses was isolation of a monolayer of graphene. In 2004 Geim et al. were able to isolate a graphene sheet for the first time [3, 4]. Graphite's existence had been known for several centuries prior and was applied in some mechanical settings [2]. However, it was not until 1985 and the discovery of fullerenes (C<sub>60</sub>) that there was thought to be the possibility of interesting and beneficial properties of this structure and its subsequent derivatives, assuming that it could be synthesized in large quantities [1].

- 
- [1] H. W. Kroto, J. R. Heath, S. C. O'Brien, R. F. Curl, and R. E. Smalley. C<sub>60</sub>: Buckminsterfullerene. *Nature*, 318:162–163, 1985.
  - [2] Ruben Mas-Balleste, Cristina Gomez-Navarro, Julio Gomez-Herrero, and Felix Zamora. 2d materials: to graphene and beyond. *Nanoscale*, 3:20–30, 2011.
  - [3] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A.

- Firsov. Electric field effect in atomically thin carbon films. *Science*, 306(5696):666–669, 2004.
- [4] K. S. Novoselov, D. Jiang, F. Schedin, T. J. Booth, V. V. Khotkevich, S. V. Morozov, and A. K. Geim. Two-dimensional atomic crystals. *Proceedings of the National Academy of Sciences of the United States of America*, 102(30):10451–10453, 2005.

---

\* kraig.andrews@wayne.edu