



## MDS 6106 – Introduction to Optimization

### Solutions 2

#### Exercise E2.1 (Optimization Problem I):

Consider the function  $f_\alpha : \mathbb{R}^2 \rightarrow \mathbb{R}$ ,

$$f_\alpha(x) := \alpha x_1^2 + x_2^2 - 2x_1x_2 - 2x_2,$$

where  $\alpha \in \mathbb{R}$  is a scalar.

- Find the stationary points (in case they exist) of  $f_\alpha$  for each value of  $\alpha$ .
- For each stationary point  $x^*$  in part a), determine whether  $x^*$  is a local maximizer or a local minimizer or a saddle point of  $f_\alpha$ .
- For which values of  $\alpha$  can  $f_\alpha$  have a global minimizer?

#### *Solution :*

- a) The gradient and Hessian of  $f_\alpha$  are given by

$$\nabla f_\alpha(x) = \begin{pmatrix} 2\alpha x_1 - 2x_2 \\ 2x_2 - 2x_1 - 2 \end{pmatrix}, \quad \nabla^2 f_\alpha(x) = \begin{pmatrix} 2\alpha & -2 \\ -2 & 2 \end{pmatrix}$$

and it holds that

$$\nabla f_\alpha(x) = 0 \iff x_2 = x_1 + 1 \quad \text{and} \quad 2\alpha x_1 - 2x_1 = 2$$

which implies  $(\alpha - 1)x_1 = 1$ . This equation only has a solution if  $\alpha \neq 1$ . In this case, we obtain

$$x_1^* = \frac{1}{\alpha - 1}, \quad x_2^* = 1 + \frac{1}{\alpha - 1} = \frac{\alpha}{\alpha - 1}.$$

This is also the unique stationary point of  $f_\alpha$ .

- b) We have

$$\text{tr}(\nabla^2 f_\alpha(x)) = 2\alpha + 2 = 2(\alpha + 1), \quad \det(\nabla^2 f_\alpha(x)) = 4\alpha - 4 = 4(\alpha - 1) \quad (\forall x).$$

Consequently, if  $\alpha < 1$ , the Hessian is indefinite and  $x^*$  is a saddle point. If  $\alpha > 1$ , then  $\nabla^2 f_\alpha(x)$  is positive definite and  $x^*$  is a local minimizer of  $f_\alpha$ .

- c) The function  $f_\alpha$  can only have a global minimizer if  $\alpha > 1$ . In the case  $\alpha \leq 1$ ,  $f_\alpha$  is unbounded and it does not possess a global minimizer (all stationary points are saddle points). Moreover, since the mapping  $f_\alpha$  is quadratic (i.e., the Hessian  $\nabla^2 f_\alpha(x)$  does not depend on  $x$ ), the positive definiteness of the Hessian implies that  $f_\alpha$  is strongly convex for all  $\alpha > 1$ . Hence,  $x^*$  is the unique global solution of  $\min_x f_\alpha(x)$  in this situation.

**Exercise E2.2 (Optimization Problem II):**

We consider the optimization problem

$$\min_{x \in \mathbb{R}^2} f(x) = \frac{1}{2}x_1^2x_2^2 + \frac{1}{2}x_1^2 + \frac{1}{2}x_2^2 - 2x_1.$$

- Is the function  $f$  coercive?
- Calculate the gradient and Hessian of  $f$  and determine all stationary points of  $f$ .
- Show that  $f$  has a unique global minimizer.
- Is the mapping  $f$  convex?

**Solution :**

- Yes, the function  $f$  is coercive. To see this, we use the estimate

$$f(x) \geq \frac{1}{2}(x_1^2 - 4x_1 + 4) + \frac{1}{2}x_2^2 - 2 = \frac{1}{2}(x_1 - 2)^2 + \frac{1}{2}x_2^2 - 2.$$

Obviously the latter lower bound converges to  $+\infty$  if either  $|x_1| \rightarrow \infty$  or  $|x_2| \rightarrow \infty$ . This establishes coercivity of  $f$ .

- It holds that

$$\nabla f(x) = \begin{pmatrix} x_1x_2^2 + x_1 - 2 \\ x_1^2x_2 + x_2 \end{pmatrix}, \quad \nabla^2 f(x) = \begin{pmatrix} x_2^2 + 1 & 2x_1x_2 \\ 2x_1x_2 & x_2^2 + 1 \end{pmatrix}.$$

Furthermore, we have  $\nabla f(x) = 0$  if and only if  $x_2(1 + x_1^2) = 0$  which implies  $x_2 = 0$  and  $x_1 = 2$ . Thus,  $x^* = (2, 0)^\top$  is the single stationary point of  $f$ .

- Since  $f$  is coercive, it possesses a global solution. Every global solution of  $\min_x f(x)$  is a stationary point of  $f$ . Since  $f$  has only one stationary point, this implies that  $x^*$  is the unique global solution of the problem  $\min_x f(x)$ .
- We have  $\text{tr}(\nabla^2 f(x)) = x_1^2 + x_2^2 + 2 \geq 0$  and

$$\det(\nabla^2 f(x)) = (x_1^2 + 1)(x_2^2 + 1) - 4x_1^2x_2^2 = -3x_1^2x_2^2 + x_1^2 + x_2^2 + 1.$$

Consequently, setting  $x_1 = x_2 = \sqrt{2}$ , we obtain  $\det(\nabla^2 f(x)) = -3 \cdot 4 + 2 + 2 + 1 = -7$  which shows that  $\nabla^2 f(x)$  is indefinite at  $x = (\sqrt{2}, \sqrt{2})^\top$ . Hence,  $f$  can not be convex on  $\mathbb{R}^2$ .

**Exercise E2.3 (Convex Sets and Convex Functions):**

In this exercise, we investigate convexity of sets and functions.

- Let  $A \in \mathbb{R}^{n \times n}$  be a given symmetric and positive semidefinite matrix and consider the set  $X := \{x \in \mathbb{R}^n : x^\top A x \leq 0\}$ . Show that the set  $X$  is convex.
- Show that the function  $f : \mathbb{R}_{++}^2 \rightarrow \mathbb{R}$ ,  $f(x) = x_1^2 - 2x_1x_2 + x_2^2 - \ln(x_1x_2)$  is strictly convex on  $\mathbb{R}_{++}^2 := \{x \in \mathbb{R}^2 : x > 0\}$ .
- Determine whether the function  $f(x) = -x_1^2 - x_2^2 - 2x_3^2 + x_1x_2$  is convex or concave.
- Show that the hyperbolic set  $\{x \in \mathbb{R}_+^2 : x_1x_2 \geq 1\}$  is convex, where  $\mathbb{R}_+^2 := \{x \in \mathbb{R}^2 : x \geq 0\}$ .

**Hint:** Rewrite the condition " $x_1x_2 \geq 1$ " in a suitable way.

**Solution :**

- a) Since the matrix  $A$  is symmetric and positive semidefinite, the mapping  $f(x) = x^\top Ax$  is convex. As a consequence the level set  $L_{\leq 0} := \{x \in \mathbb{R}^n : f(x) \leq 0\} = X$  is convex. (See lecture L-05, slide 15).
- b) We calculate the gradient and Hessian of  $f$ . It holds that

$$\nabla f(x) = \begin{pmatrix} 2x_1 - 2x_2 - \frac{1}{x_1} \\ 2x_2 - 2x_1 - \frac{1}{x_2} \end{pmatrix}, \quad \nabla^2 f(x) = \begin{pmatrix} 2 + \frac{1}{x_1^2} & -2 \\ -2 & 2 + \frac{1}{x_2^2} \end{pmatrix}$$

Furthermore, we have  $\text{tr}(\nabla^2 f(x)) = 4 + \frac{1}{x_1^2} + \frac{1}{x_2^2} > 4$  for all  $x \in \mathbb{R}_{++}^2$  and

$$\det(\nabla^2 f(x)) = \left(2 + \frac{1}{x_1^2}\right) \left(2 + \frac{1}{x_2^2}\right) - 4 = \frac{2}{x_1^2} + \frac{2}{x_2^2} + \frac{1}{x_1^2 x_2^2} > 0$$

for all  $x \in \mathbb{R}_{++}^2$ . This implies that  $\nabla^2 f(x)$  is positive definite for all  $x \in \mathbb{R}_{++}^2$  and hence,  $f$  is (strictly) convex on  $\mathbb{R}_{++}^2$ .

- c) As in part b), we utilize definiteness of the Hessian to determine convexity. We have

$$\nabla f(x) = \begin{pmatrix} -2x_1 + x_2 \\ -2x_2 + x_1 \\ -4x_3 \end{pmatrix}, \quad \nabla^2 f(x) = \begin{pmatrix} -2 & 1 & 0 \\ 1 & -2 & 0 \\ 0 & 0 & -4 \end{pmatrix}.$$

We calculate the eigenvalues of  $\nabla^2 f(x)$ :

$$\det \left( \begin{bmatrix} -2 - \lambda & 1 & 0 \\ 1 & -2 - \lambda & 0 \\ 0 & 0 & -4 - \lambda \end{bmatrix} \right) = -(4 + \lambda)[(2 + \lambda)^2 - 1] = -(4 + \lambda)[3 + 4\lambda + \lambda^2].$$

This yields  $\lambda_1 = -4$ ,  $\lambda_{2/3} = \frac{1}{2}(-4 \pm \sqrt{16 - 12}) = \frac{1}{2}(-4 \pm 2) = -3 / -1$ . Consequently,  $\nabla^2 f(x)$  is negative definite for all  $x$  which shows that  $f$  is a concave mapping.

- d) We first notice  $X := \{x \in \mathbb{R}_+^2 : x_1 x_2 \geq 1\} = \{x \in \mathbb{R}_{++}^2 : x_1 x_2 \geq 1\}$ , where  $\mathbb{R}_{++}^2 = \{x \in \mathbb{R}^2 : x > 0\}$ . Since the (natural) logarithm  $\ln(\cdot)$  is monotonically increasing, the condition  $x_1 x_2 \geq 1$  is equivalent to

$$f(x_1, x_2) := -\ln(x_1) - \ln(x_2) \leq 0.$$

The gradient and Hessian of  $f$  are given by:

$$\nabla f(x_1, x_2) = \begin{pmatrix} -x_1^{-1} \\ -x_2^{-1} \end{pmatrix}, \quad \nabla^2 f(x_1, x_2) = \begin{pmatrix} x_1^{-2} & 0 \\ 0 & x_2^{-2} \end{pmatrix}$$

Since  $\nabla^2 f$  is positive definite on the open and convex set  $\mathbb{R}_{++}^2$ , it follows that  $f$  is convex on  $\mathbb{R}_{++}^2$ . This shows that the set  $X$  is convex.

**Exercise E2.4 (Multiple Choice):**

Answer the following multiple choice questions and decide whether the statements are true or false. Try to give short explanations of your answer.

- a) *A point is a stationary point of  $f$  if and only if it is a local minimum or maximum of  $f$ .*  
☐ True. ☐ False.
- b) *The point  $x^*$  is a stationary point of  $f$  if and only if  $\nabla f(x^*)^\top h = 0$  for all  $h \in \mathbb{R}^n \setminus \{0\}$ .*  
☐ True. ☐ False.
- c) *Let  $X \subset \mathbb{R}^n$  be nonempty and let  $x^* \in X$  be a global solution of  $\min_{x \in X} f(x)$ . Then, it holds that  $\nabla f(x^*) = 0$ .*  
☐ True. ☐ False.
- d) *The function  $f(x) = x^4$  is strongly convex.*  
☐ True. ☐ False.
- e) *Let  $I := [a, b]$  be given with  $a, b \in \mathbb{R}$ ,  $a < b$ . Suppose that  $g : I \rightarrow \mathbb{R}$  is a convex function. Then,  $g^2$  is strictly convex.*  
☐ True. ☐ False.
- f) *Suppose that  $g : \mathbb{R} \rightarrow \mathbb{R}$  is convex but not strictly convex. Then,  $g$  has either no global minimum or infinitely many.*  
☐ True. ☐ False.
- g) *Suppose we have  $\lambda_{\min}(\nabla^2 f(x)) > 0$  for all  $x \in \mathbb{R}^n$ , then  $f$  is strongly convex.*  
☐ True. ☐ False.

**Solution :**

- a) *False.* Consider the function  $f(x) = x^3$ . Then, the point  $x^* = 0$  is a stationary point, i.e., it satisfies  $f'(x^*) = 3(x^*)^2 = 0$ . However,  $x^*$  is a saddle-point, i.e., it's neither a minimum nor a maximum.
- b) *True.* If  $x^*$  is a stationary point then we have  $\nabla f(x^*) = 0$  and the condition is obviously satisfied. On the other hand, let us assume  $\nabla f(x^*)^\top h = 0$  for all  $h \in \mathbb{R}^n \setminus \{0\}$ . In the case  $\nabla f(x^*) = 0$ , there is nothing to show. Hence, let us assume  $\nabla f(x^*) \neq 0$ . But then setting  $h = \nabla f(x^*) \neq 0$  implies  $\|\nabla f(x^*)\|^2 = 0$  which is a contradiction to  $\nabla f(x^*) \neq 0$  and thus, this case cannot occur. Together, we can infer  $\nabla f(x^*) = 0$  which finishes the proof.
- c) *False.* Consider the problem  $\min_{x \in X} f(x) = x$  with  $X = [1, \infty)$ . Obviously,  $x^* = 1$  is the global solution of this problem, but we have  $f'(x^*) = 1 \neq 0$ .
- d) *False.* It holds that  $f''(x) = 12x^2$  and  $f''(0) = 0$ . Hence, by Theorem 4.15,  $f(x) = x^4$  cannot be strongly convex.
- e) *False.* Take  $I = [0, 1]$  and  $g(x) = 1$ . This function is convex, but  $g(x)^2 = 1 = g(x)$  is not strictly convex.
- f) *False.* The function  $g(x) = |x|$  is convex but not strictly convex. Moreover, it has a unique global minimum at  $x = 0$ .

- g) *False*. For strong convexity a uniform lower bound for the minimal eigenvalue of the Hessian is required. A possible counterexample is the function  $f(x) = e^x$ .

**Assignment A2.1 (Optimization Problem):**

(approx. 20 points)

Consider the unconstrained optimization problem

$$\min_{x \in \mathbb{R}^2} f(x) = \frac{1}{3}x_1^3 - x_1 \left( \frac{3}{2} + x_2^2 \right) + x_2^4.$$

- Is the function  $f$  coercive?
- Compute the gradient and Hessian of  $f$  and calculate all stationary points.
- For each stationary point  $x^*$  found in part c) investigate whether  $x^*$  is a local maximizer, local minimizer, or saddle point and explain your answer.
- Does the mapping  $f$  possess any strict local or global minimizer?

**Solution :**

- a) No, the function  $f$  is not coercive. We have

$$f(x_1, 0) = \frac{1}{6}x_1(2x_1^2 - 9) \rightarrow \pm\infty \quad \text{if } x_1 \rightarrow \pm\infty.$$

- b) The gradient and Hessian of  $f$  are given by

$$\nabla f(x) = \begin{pmatrix} x_1^2 - x_2^2 - \frac{3}{2} \\ -2x_1x_2 + 4x_2^3 \end{pmatrix} \quad \text{and} \quad \nabla^2 f(x) = \begin{pmatrix} 2x_1 & -2x_2 \\ -2x_2 & -2x_1 + 12x_2^2 \end{pmatrix}.$$

Moreover, it holds that  $\nabla f(x) = 0$  if and only if  $2x_2(2x_2^2 - x_1) = 0$ . Let us first consider the case  $x_2 = 0$ . Then, it follows  $x_1 = \pm\sqrt{3}/2$ . Otherwise, we have  $x_1 = 2x_2^2$  which implies  $4x_2^4 - x_2^2 - \frac{3}{2} = 0$ , i.e.,

$$x_2^2 = \frac{1 \pm \sqrt{1 + 6 \cdot 4}}{2 \cdot 4} = \frac{1 \pm 5}{8} = \frac{3}{4} \text{ or } -\frac{1}{2}$$

This yields  $x_2 = \pm\sqrt{3}/2$  and  $x_1 = 3/2$ . In total,  $f$  has the following for stationary points:

$$\bar{x}_1 = \left( \frac{\sqrt{6}}{2}, 0 \right), \quad \bar{x}_2 = \left( -\frac{\sqrt{6}}{2}, 0 \right), \quad \bar{x}_3 = \left( \frac{3}{2}, \frac{\sqrt{3}}{2} \right), \quad \bar{x}_4 = \left( \frac{3}{2}, -\frac{\sqrt{3}}{2} \right).$$

- c) We have

$$\nabla^2 f(\bar{x}_1) = \begin{pmatrix} \sqrt{6} & 0 \\ 0 & -\sqrt{6} \end{pmatrix} \quad \text{and} \quad \nabla^2 f(\bar{x}_2) = \begin{pmatrix} -\sqrt{6} & 0 \\ 0 & \sqrt{6} \end{pmatrix}.$$

Both Hessians are diagonal matrices with eigenvalues  $-\sqrt{6}$  and  $\sqrt{6}$  and, hence  $\nabla^2 f(\bar{x}_1)$  and  $\nabla^2 f(\bar{x}_2)$  are indefinite and the stationary points  $\bar{x}_1$  and  $\bar{x}_2$  are saddle points. Furthermore, it holds that

$$\nabla^2 f(\bar{x}_3) = \begin{pmatrix} 3 & -\sqrt{3} \\ -\sqrt{3} & 6 \end{pmatrix} \quad \text{and} \quad \nabla^2 f(\bar{x}_4) = \begin{pmatrix} 3 & -\sqrt{3} \\ -\sqrt{3} & 6 \end{pmatrix}$$

and  $\text{tr}(\nabla^2 f(\bar{x}_3)) = \text{tr}(\nabla^2 f(\bar{x}_4)) = 9 > 0$  and  $\det(\nabla^2 f(\bar{x}_3)) = \det(\nabla^2 f(\bar{x}_4)) = 18 - 3 > 0$ . This shows that  $\nabla^2 f(\bar{x}_3)$  and  $\nabla^2 f(\bar{x}_4)$  are positive definite. Thus, by the second order sufficient conditions,  $\bar{x}_3$  and  $\bar{x}_4$  are strict local minimizers.

- d) Our results in a) and c) show that  $\bar{x}_3$  and  $\bar{x}_4$  are strict local solutions and that  $f$  does not possess a global minimum.

**Assignment A2.2 (Another Optimization Problem):**

(approx. 15 points)

Consider the unconstrained optimization problem

$$\min_{x \in \mathbb{R}^3} f(x) = x_1^4 - 2x_1^2 + x_2^2 + 2x_2x_3 + 2x_3^2.$$

- Is the mapping  $f$  coercive?
- Verify whether the function  $f$  is convex on  $\mathbb{R}^3$ .
- Find all stationary points of  $f$  and classify them according to whether they are saddle points, strict / non-strict, local / global, minimum / maximum points.

**Solution :**

- Yes, the mapping  $f$  coercive. In particular, we have

$$\begin{aligned} f(x) &= x_1^2(x_1^2 - 2) + \frac{1}{3}x_2^2 + \frac{1}{2}x_3^2 + \left(\frac{2}{3}x_2^2 + 2x_2x_3 + \frac{3}{2}x_3^2\right) \\ &= x_1^2(x_1^2 - 2) + \frac{1}{3}x_2^2 + \frac{1}{2}x_3^2 + \left(\sqrt{\frac{2}{3}}x_2 + \sqrt{\frac{3}{2}}x_3\right)^2 \geq x_1^2(x_1^2 - 2) + \frac{1}{3}x_2^2 + \frac{1}{2}x_3^2. \end{aligned}$$

Since  $x_1^4$  dominates  $x_1^2$  (i.e., it grows faster than  $x_1^2$  as  $|x_1| \rightarrow \infty$ , this establishes coercivity of  $f$ .

- We calculate the gradient and Hessian of  $f$ . It holds that

$$\nabla f(x) = \begin{pmatrix} 4x_1^3 - 4x_1 \\ 2x_2 + 2x_3 \\ 2x_2 + 4x_3 \end{pmatrix}, \quad \nabla^2 f(x) = \begin{pmatrix} 12x_1^2 - 4 & 0 & 0 \\ 0 & 2 & 2 \\ 0 & 2 & 4 \end{pmatrix},$$

and

$$e_1^\top \nabla^2 f(x) e_1 = 12x_1^2 - 4.$$

For all  $x_1 \in (-1/\sqrt{3}, 1/\sqrt{3})$ , this term is negative which implies that  $f$  can not be convex on  $\mathbb{R}^3$ .

- We have  $4x_1(x_1^2 - 1) = 0$  and  $x_2 = x_3 = 0$  which yields the stationary points

$$\bar{x}_1 = (0, 0, 0), \quad \bar{x}_2 = (1, 0, 0), \quad \bar{x}_3 = (-1, 0, 0).$$

Furthermore, it holds that  $\det(\nabla^2 f(\bar{x}_1) - \lambda I) = (-4 - \lambda)[(2 - \lambda)(4 - \lambda) - 4] = -(4 + \lambda)(4 - 6\lambda + \lambda^2)$ . Hence, the eigenvalues of  $\nabla^2 f(\bar{x}_1)$  are given by  $-4, 3 \pm \sqrt{5}$ . This shows that  $\nabla^2 f(\bar{x}_1)$  is indefinite and thus,  $\bar{x}_1$  is a saddle point. Similarly, we obtain  $\det(\nabla^2 f(\bar{x}_{2/3}) - \lambda I) = (8 - \lambda)(4 - 6\lambda + \lambda^2)$  and consequently,  $\nabla^2 f(\bar{x}_{2/3})$  is positive definite and the stationary points  $\bar{x}_2$  and  $\bar{x}_3$  are strict local minimizer. Since the function  $f$  is coercive, it possesses at least one global minimizer that needs to be a stationary point. Due to  $f(\bar{x}_2) = f(\bar{x}_3) = -1$ , all local minimizer have the same function value and we can infer that  $\bar{x}_2$  and  $\bar{x}_3$  are global solutions of the problem  $\min_x f(x)$ .

**Assignment A2.3 (Convex Sets):**

(approx. 20 points)

In this exercise, we study convexity of various sets.

- a) Verify whether the following sets are convex or not and explain your answer!

$$X_1 = \{x \in \mathbb{R}^n : \alpha \leq (a^\top x)^2 \leq \beta\}, \quad \alpha, \beta \in \mathbb{R}, \quad \alpha \leq \beta, \quad a \in \mathbb{R}^n,$$

$$X_2 = \{x \in \mathbb{R}^n : \|x - a\|_2 \leq \|x - b\|_2\}, \quad a, b \in \mathbb{R}^n, \quad a \neq b.$$

- b) Let the set of all positive semidefinite and symmetric  $n \times n$  matrices be denoted by  $\mathbb{S}_+^n$ . Show that the set  $X := \{A \in \mathbb{R}^{n \times n} : A \in \mathbb{S}_+^n \text{ and } \text{tr}(A) = 1\}$  is a convex subset of  $\mathbb{R}^{n \times n}$ .
- c) Decide whether the following statements are true or false. Explain your answer and either present a proof / verification or a counter-example.
- The union of two convex sets  $X_1, X_2 \subset \mathbb{R}^n$ ,  $X_1 \neq X_2$  is never a convex set.
  - Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a concave. Then, the set  $X := \{x \in \mathbb{R}^n : f(x) \geq 0\}$  is convex.

**Solution :**

- a) The set  $X_1$  is not convex. We can choose  $n = 1$  and  $\alpha = \beta = a = 1$ . Then, it follows  $X_1 = \{x \in \mathbb{R} : x^2 = 1\} = \{\pm 1\}$ . This set is obviously not convex.

The condition  $\|x - a\| \leq \|x - b\|$  in the definition of  $X_2$  is equivalent to

$$\|x - a\|^2 \leq \|x - b\|^2 \iff -2a^\top x + \|a\|^2 \leq -2b^\top x + \|b\|^2 \iff 2(b - a)^\top x \leq \|b\|^2 - \|a\|^2.$$

Hence,  $X_2$  is a convex half space.

- b) Let  $A, B \in X$  be two symmetric, positive semidefinite matrices and let  $\lambda \in [0, 1]$  be arbitrary. Then, the matrix  $C = \lambda A + (1 - \lambda)B$  is symmetric and it follows

$$x^\top C x = \lambda[x^\top A x] + (1 - \lambda)[x^\top B x] \geq 0, \quad \forall x \in \mathbb{R}^n,$$

where we used the positive semidefiniteness of  $A$  and  $B$ . Hence, we have  $C \in \mathbb{S}_+^n$  and by the linearity of the trace operator, it follows  $\text{tr}(C) = \lambda \text{tr}(A) + (1 - \lambda) \text{tr}(B) = \lambda + 1 - \lambda = 1$ . This shows that  $X$  is a convex subset of  $\mathbb{R}^{n \times n}$ .

- c) The first statement is wrong: consider  $X_1 = [0, 1]$  and  $X_2 = [-1, 0]$ . Then  $X_1 \neq X_2$  and  $X_1 \cup X_2 = [-1, 1]$  is a convex set.

We verify the second statement briefly. Let  $x, y \in X$  and  $\lambda \in [0, 1]$  be arbitrary. To establish convexity of  $X$ , we need to show that  $\lambda x + (1 - \lambda)y \in X$ . Utilizing the concavity of  $f$  and  $x, y \in X$ , we have

$$g(\lambda x + (1 - \lambda)y) \geq \lambda g(x) + (1 - \lambda)g(y) \geq 0.$$

Hence,  $X$  is a convex set.

**Assignment A2.4 (Convex Functions):**

(approx. 25 points)

In this exercise, convexity properties of different functions are investigated.

a) Verify that the following functions are convex over the specified domain:

- $f : \mathbb{R}_{++} \rightarrow \mathbb{R}$ ,  $f(x) := \sqrt{1+x^{-2}}$ , where  $\mathbb{R}_{++} := \{x \in \mathbb{R} : x > 0\}$ .
- $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f(x) := \frac{1}{2}\|Ax - b\|^2 + \mu\|Lx\|^2$ , where  $A \in \mathbb{R}^{m \times n}$ ,  $L \in \mathbb{R}^{p \times n}$ ,  $b \in \mathbb{R}^m$ , and  $\mu > 0$  are given.
- $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ ,  $f(x, y) := \frac{\lambda}{2}\|x\|^2 + \sum_{i=1}^m \max\{0, 1 - b_i(a_i^\top x + y)\}$ , where  $a_i \in \mathbb{R}^n$  and  $b_i \in \{-1, 1\}$  are given data points for all  $i = 1, \dots, m$  and  $\lambda > 0$  is a parameter.

b) Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a convex mapping and set  $g(x) := (f(x))^2$ . Is the function  $g$  convex? Explain your answer and either present a brief verification or a counter-example.

Is the mapping  $x \mapsto \frac{1}{2}(\|x\|^2 - 1)^2$  convex?

**Solution :**

a) The mapping  $f(x) = \sqrt{1+x^{-2}}$  is twice continuously differentiable on the convex set  $\mathbb{R}_{++}$  and it holds that

$$f'(x) = \frac{1}{2\sqrt{1+x^{-2}}} \frac{-2}{x^3} = \frac{-1}{x^2\sqrt{1+x^2}}, \quad f''(x) = \frac{2x\sqrt{1+x^2} + \frac{x^3}{\sqrt{1+x^2}}}{x^4(1+x^2)} = \frac{3x^2+2}{x^3(1+x^2)\sqrt{1+x^2}}.$$

Notice that we have  $f''(x) > 0$  for all  $x > 0$  and thus,  $f$  is (strictly) convex on  $\mathbb{R}_{++}$ .

In the second example, we obtain

$$\nabla f(x) = A^\top(Ax - b) + \mu L^\top Lx \quad \text{and} \quad \nabla^2 f(x) = A^\top A + \mu L^\top L$$

Thus, it follows  $h^\top \nabla^2 f(x) h = h^\top A^\top A h + \mu h^\top L^\top L h = \|Ah\|^2 + \mu\|Lh\|^2 \geq 0$  for all  $h \in \mathbb{R}^n$ . This establishes convexity of  $f$  on  $\mathbb{R}^n$ .

Finally, let us define  $g(x, y) = \frac{\lambda}{2}\|x\|^2$  and  $g_i(x, y) = \max\{0, 1 - b_i(a_i^\top x + y)\}$ . Then,  $f$  can be interpreted as the sum of the functions  $g$  and  $g_i$ ,  $i = 1, \dots, m$  and convexity follows if each of the functions  $g$ ,  $g_i$ ,  $i = 1, \dots, m$  is convex. The mapping  $g_i$  is the composition of the max-function  $z \mapsto \max\{0, z\}$  and the affine-linear function  $(x, y) \mapsto h_i(x, y) := 1 - b_i(a_i^\top x + y)$ . Since  $h_i$  is convex (as a linear mapping), the max-function  $g_i$  also is convex! Finally, the Hessian of  $g$  is given by

$$\mathbb{R}^{(n+1) \times (n+1)} \ni \nabla^2 g(x, y) = \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} \succeq 0.$$

Combining these different results, we see that  $f$  is a convex function.

b) No, the mapping  $g$  generally does not need to be convex. We can consider the example  $g(x) = \frac{1}{2}(x^2 - 1)^2 = \frac{1}{2}x^4 - x^2 + \frac{1}{2}$ . The derivatives of  $g$  are given by  $g'(x) = 2x^3 - 2x$  and  $g''(x) = 6x^2 - 2$ . Since  $g''$  takes negative values around  $x = 0$ , the mapping  $g$  can not be convex. This also verifies that  $\frac{1}{2}(\|x\|^2 - 1)^2$  is not a convex function.

**Remark:** Since the function  $x \mapsto x^2$  is monotonically increasing on  $[0, \infty)$ , we can infer convexity in the case  $f(x) \geq 0$  for all  $x \in \mathbb{R}^n$ .



**Assignment A2.5 (A Penalty Problem):**

(approx. 20 points)

We consider the parametrized optimization problem

$$\min_x f_\beta(x) := \frac{1}{2}\|x - b\|^2 + \frac{\beta}{2} \left( \sum_{i=1}^n x_i \right)^2, \quad x \in \mathbb{R}^n, \quad (1)$$

where  $b \in \mathbb{R}^n$  is given and  $\beta \geq 0$  is a parameter.

- Calculate the gradient and Hessian of  $f_\beta$ .
- Show that the mapping  $f_\beta$  is strongly convex for all  $\beta \geq 0$ .
- Show that  $f_\beta$  has a unique stationary point  $x_\beta^*$  and compute it explicitly. Determine whether  $x_\beta^*$  is a local minimizer, a local maximizer, or a saddle point of problem (1).
- For  $\beta \rightarrow \infty$ , the solutions  $x_\beta^*$  converge to a point  $x^*$ . Calculate the limit  $x^* = \lim_{\beta \rightarrow \infty} x_\beta^*$  explicitly and show that  $x^*$  satisfies the constraint  $\mathbf{1}^\top x^* = \sum_{i=1}^n x_i^* = 0$ .

**Solution :**

- The objective function is given by  $f_\beta(x) = \frac{1}{2} \sum_{i=1}^n (x_i - b_i)^2 + \frac{\beta}{2} (\sum_{i=1}^n x_i)^2$ . Hence, we obtain

$$\frac{\partial f_\beta}{\partial x_i}(x) = (x_i - b_i) + \beta \left( \sum_{i=1}^n x_i \right) \cdot 1 \quad \text{and} \quad \frac{\partial^2 f_\beta}{\partial x_i \partial x_j}(x) = \begin{cases} 1 + \beta & \text{if } i = j, \\ \beta & \text{if } i \neq j. \end{cases}$$

Due to  $\mathbf{1}^\top x = \sum_{i=1}^n x_i$  and  $(\mathbf{1}\mathbf{1}^\top)_{ij} = 1$  for all  $i, j$ , this shows that  $\nabla f_\beta$  and  $\nabla^2 f_\beta$  are given by:

$$\nabla f_\beta(x) = x - b + \beta \cdot (\mathbf{1}^\top x) \mathbf{1}, \quad \nabla^2 f_\beta(x) = I + \beta \cdot \mathbf{1}\mathbf{1}^\top.$$

- We have

$$d^\top \nabla^2 f_\beta(x) d = d^\top d + \beta (\mathbf{1}^\top d)^2 \geq \|d\|^2 \quad \forall d \in \mathbb{R}^n$$

and hence, the Hessian  $\nabla^2 f_\beta$  is uniformly positive definite for all  $x$ . This implies that  $f_\beta$  is 1-strongly convex for all  $\beta \geq 0$ .

- Multiplying the condition  $\nabla f_\beta(x) = 0$  with  $\mathbf{1}^\top$ , we obtain

$$\mathbf{1}^\top x - \mathbf{1}^\top b + \beta n \cdot (\mathbf{1}^\top x) = 0 \quad \implies \quad \mathbf{1}^\top x = \frac{\mathbf{1}^\top b}{1 + \beta n}.$$

Using this formula, it now follows  $\nabla f_\beta(x) = 0$  if and only if  $x = b - \frac{\beta}{1 + \beta n} \cdot (\mathbf{1}^\top b) \mathbf{1}$ . Since all stationary points are uniquely characterized in this way, this implies that  $x_\beta^* = b - \frac{\beta}{1 + \beta n} \cdot (\mathbf{1}^\top b) \mathbf{1}$  is the unique stationary point of problem (1). Moreover, due to part b), we know that  $x^*$  is a strict global minimizer of problem (1).

- We have  $x_\beta^* = b - \frac{1}{1/\beta + n} \cdot (\mathbf{1}^\top b) \mathbf{1} \rightarrow b - \frac{1}{n} \cdot (\mathbf{1}^\top b) \mathbf{1} =: x^*$  as  $\beta \rightarrow \infty$ . Moreover, we obtain

$$\mathbf{1}^\top x^* = \mathbf{1}^\top b - \frac{\mathbf{1}^\top \mathbf{1}}{n} \mathbf{1}^\top b = 0.$$