

# Test a Perceptual Phenomenon

January 28, 2018

## 0.0.1 Analyzing the Stroop Effect

Perform the analysis in the space below. Remember to follow [the instructions](#) and review the [project rubric](#) before submitting. Once you've completed the analysis and write up, download this file as a PDF or HTML file and submit in the next section.

- (1) What is the independent variable? What is the dependent variable?

The independent variable is the test given (congruent or incongruent). The dependent variable is the participant's reaction time.

- (2) What is an appropriate set of hypotheses for this task? What kind of statistical test do you expect to perform? Justify your choices.

In this task, we are examining a small sample of a population in order to determine whether or not the stroop effect is present. We do this by comparing the time differences of the two tests set forth - the first being the congruent test, and the second being the incongruent test.

Null Hypothesis (H0) The null hypothesis states that there is no difference in the mean (average) time it took to complete both the incongruent (i) and congruent (c) tests and, therefore, that the Stroop effect is not present.

H0:  $c = i$

Alternative Hypothesis (H1) The alternative hypothesis is that there is a statistically significant difference in the response times of the congruent and incongruent tests, thereby proving that the Stroop effect had occurred. The incongruent test should take more time than the congruent test.

H1:  $c < i$

Since we are comparing two sample sets of data, we will use a paired t-Test. A paired t-Test compares the means of two related groups to see if there are statistically significant differences between them.

since each entry of i and c times is from same indiv we use pttest to see if mean of diff is 0, if it is not, null hypo refused

we suppose that in the population the variable- time of response- is normally distributed it means that for small samples will be distributed as a student t distribution

Our reasoning for using the paired t-Test is as follows: 1. We have a small sample size (there are only 24 unique tests). 2. We assume that the sample is normal and representative of the population as a whole. 3. The way the data is structured: 24 individuals submitted two test results each. A paired t\_test is the best way to compare these. We're measuring each person on the same dependent variable- the time it takes to complete each test. We are looking for differences in the mean response time for each test.

- (3) Report some descriptive statistics regarding this dataset. Include at least one measure of central tendency and at least one measure of variability. The name of the data file is 'stroop-data.csv'.

```
In [2]: # Perform the analysis here
```

```
import pandas as pd

stroop = pd.read_csv('stroopdata.csv')

print(stroop)

stroop.describe()
```

	Congruent	Incongruent
0	12.079	19.278
1	16.791	18.741
2	9.564	21.214
3	8.630	15.687
4	14.669	22.803
5	12.238	20.878
6	14.692	24.572
7	8.987	17.394
8	9.401	20.762
9	14.480	26.282
10	22.328	24.524
11	15.298	18.644
12	15.073	17.510
13	16.929	20.330
14	18.200	35.255
15	12.130	22.158
16	18.495	25.139
17	10.639	20.429
18	11.344	17.425
19	12.369	34.288
20	12.944	23.894
21	14.233	17.960
22	19.710	22.058
23	16.004	21.157

```
Out[2]:
```

	Congruent	Incongruent
count	24.000000	24.000000
mean	14.051125	22.015917
std	3.559358	4.797057
min	8.630000	15.687000
25%	11.895250	18.716750
50%	14.356500	21.017500

75%	16.200750	24.051500
max	22.328000	35.255000

For the congruent set: The mean is 14.05 and the median is 14.35. The standard deviation is 3.559. For the incongruent set: the mean is 22.02 and the median is 21.02. The standard deviation is 4.797.

In both, the standard deviation is relatively smaller than the mean.

Since the mean and median in both are very close, we know that the distribution will be symmetrical.

Finally, we can see that there is a distinct difference in the timing for each of these datasets- as we initially believed, the Stroop effect could be reasonably occurring as the congruent data set's average is 7 seconds faster than the average of the incongruent data set.


- (4) Provide one or two visualizations that show the distribution of the sample data. Write one or two sentences noting what you observe about the plot or plots.

```
In [5]: # Build the visualizations here
import numpy as np
import matplotlib.pyplot as plt

x = stroop['Congruent']
y = stroop['Incongruent']


n, bins, patches = plt.hist(x, 24, alpha=0.25)
n, bins, patches = plt.hist(y, 24, alpha=0.25)
plt.title('Bar chart comparing Congruent (blue) to Incongruent (orange)')
plt.show
```

Out[5]: <function matplotlib.pyplot.show>



```
In [6]: fig, ax = plt.subplots()
        ax.boxplot(stroop.T)

        plt.ylabel('Time in Seconds')
        plt.xticks([1,2], ['Congruent Test', 'Incongruent Test'])
        plt.show()
```



The first thing I noticed is that there's more overlap than I initially expected between the congruent and incongruent sets (between 15-20 seconds in the first bar chart).

For this reason, I created a box plot, to try to see how much overlap there really was- in fact, the box plot shows they don't overlap that much- more than 75% doesn't overlap with other 75%.

- (5) Now, perform the statistical test and report the results. What is the confidence level and your critical statistic value? Do you reject the null hypothesis or fail to reject it? Come to a conclusion in terms of the experiment task. Did the results match up with your expectations?

```
In [7]: #check if variance equal or not
        import scipy.stats as stats

        stats.bartlett(x, y)
```

```
Out[7]: BartlettResult(statistic=1.9756393730727815, pvalue=0.15985054445707891)
```

```
In [8]: #same variance
        #p-val is over 0.05
```

```
In [16]: import math
import numpy as np
```

```
In [17]: diff = x - y
```

```
In [32]: #check work / compute without scipy
#t-stat = (mean(a) - mean(b)) / (standard_deviation(a - b)/square_root(N))

mean_c, mean_i = pd.DataFrame.mean(stroop)

std_diff = np.std(diff)

N = len(stroop)

paired_t = (mean_c - mean_i) / (std_diff / math.sqrt(N))

print(paired_t)
```

```
-8.19321500097
```

```
In [25]: #using SciPy to compute:
```

```
stats.ttest_1samp(diff, 0)
```

```
Out[25]: Ttest_1sampResult(statistic=-8.020706944109957, pvalue=4.1030005857111781e-08)
```

```
In [28]: #The above results are different, because the SciPy formula uses N-1 instead of just N,
```

```
paired_t_check = (mean_c - mean_i) / (std_diff / math.sqrt(N-1))

print(paired_t_check)
```

```
-8.02070694411
```

In both cases, the p-value is under 0.05, thereby disproving the null hypothesis. The p-value for this test is very low (-8.1932). Since the p-value is well below 0.05, our null hypothesis is rejected- the stroop effect is occurring.

6. Optional Question: What is responsible for the effects observed? What is a similar task that would have the same effect?

These effects are caused by the Stroop Effect. It's taking participants a longer time to correctly identify the colors in the incongruent test.

A similar task could be trying to identify anything incorrectly labelled- perhaps an apple labeled as 'pear' (though, we'd have to account for other indicators present, as the smell and taste could also affect results).