

## Direct Sums and Products

In the next few sections we will introduce several ways to construct new vector spaces from ones we already have. We then study carefully how these new vector spaces depend on the old ones. At the end of the discussion we will have found a powerful new way to think about such constructions. We now start with direct sums and direct products. All vector spaces in a given construction will be over the same field. We begin with the simplest case of two vector spaces.

**Definition 1.** Let  $V_1$  and  $V_2$  be two vector spaces over the same field  $F$ . Their *(external) direct sum*

$$V_1 \oplus V_2 = \{ (v_1, v_2) \mid v_i \in V_i \}$$

as a set is simply the cartesian product, the set of all ordered pairs. The set  $V_1 \oplus V_2$  is given the structure of a vector space over  $F$  by defining the sum by

$$(u_1, u_2) + (v_1, v_2) = (u_1 + v_1, u_2 + v_2)$$

and scalar multiplication by

$$a(v_1, v_2) = (av_1, av_2) .$$

It is easy to check that  $V_1 \oplus V_2$  is a vector space with respect to these operations. The zero vector is just  $(0, 0)$  and the negative of a vector is given by

$$-(v_1, v_2) = (-v_1, -v_2) .$$

Two elements of  $V_1 \oplus V_2$  are equal (by definition) if and only if their first coordinates are equal and their second coordinates are equal. Hence each of the equations one has to verify for  $V_1 \oplus V_2$  naturally breaks into a pair of equations, one for each coordinate. It should now be clear that  $V_1 \oplus V_2$  is a vector space as each of  $V_1$  and  $V_2$  are.

There are several natural linear transformations associated with  $V_1 \oplus V_2$ . First there are

$$\begin{aligned} i_1 : V_1 &\longrightarrow V_1 \oplus V_2 \\ i_2 : V_2 &\longrightarrow V_1 \oplus V_2 \end{aligned}$$

which are given by

$$\begin{aligned} i_1(v_1) &= (v_1, 0) \\ i_2(v_2) &= (0, v_2) . \end{aligned}$$

These are clearly injective (one-to-one) and thus show that there is a subspace of  $V_1 \oplus V_2$  which “looks just like  $V_1$ ” and one that “looks just like  $V_2$ ”.

There are also linear transformations

$$\begin{aligned} p_1 : V_1 \oplus V_2 &\longrightarrow V_1 \\ p_2 : V_1 \oplus V_2 &\longrightarrow V_2 \end{aligned}$$

given by

$$\begin{aligned} p_1(v_1, v_2) &= v_1 \\ p_2(v_1, v_2) &= v_2. \end{aligned}$$

These are clearly surjective (onto) and are sometimes referred to as the natural (or canonical) *projections*.

Note that

$$\ker p_1 = \operatorname{im} i_2 = \{ (0, v_2) \mid v_2 \in V_2 \}$$

and

$$\ker p_2 = \operatorname{im} i_1 = \{ (v_1, 0) \mid v_1 \in V_1 \}.$$

One has the following equations

$$\begin{aligned} p_1 \circ i_1 &= I_{V_1} \\ p_2 \circ i_2 &= I_{V_2} \\ p_1 \circ i_2 &= 0 \\ p_2 \circ i_1 &= 0 \end{aligned}$$

and further

$$i_1 \circ p_1 + i_2 \circ p_2 = I_{V_1 \oplus V_2}$$

(see Exercise 3).

Let  $W_1 = \operatorname{im} i_1$  and  $W_2 = \operatorname{im} i_2$  be the subspaces of  $V = V_1 \oplus V_2$  just discussed. Note that these satisfy

$$W_1 + W_2 = V$$

and

$$W_1 \cap W_2 = 0.$$

**Definition 2.** Let  $V$  be a vector space over a field  $F$ . Let  $V_1$  and  $V_2$  be subspaces.  $V$  is called the (*internal*) *direct sum* of the subspaces  $V_1$  and  $V_2$  if

$$V_1 + V_2 = V$$

and

$$V_1 \cap V_2 = 0.$$

**Lemma 3.** Let  $V$  be a vector space over the field  $F$  which is the internal direct sum of the subspaces  $V_1$  and  $V_2$ . There is a natural isomorphism of  $V$  with the external direct sum of  $V_1$  and  $V_2$

$$\psi: V_1 \oplus V_2 \longrightarrow V$$

given by  $\psi(v_1, v_2) = v_1 + v_2$ .

*Proof.* See Exercise 8. □

**Remark 4.** Let  $V$  be a vector space with subspaces  $V_i$ ,  $1 \leq i \leq k$ . In order that  $V$  be the internal direct sum of the subspaces  $V_i$  one needs that  $V$  is the sum of all the subspaces

$$V = \sum_i V_i$$

and that for each  $i$

$$V_i \cap \sum_{j \neq i} V_j = 0.$$

It is not sufficient that the subspaces have pairwise trivial intersections for  $k > 2$ .

One may define the external direct sum of a finite number of vector spaces either inductively, or by using  $k$ -tuples. There is an analogous isomorphism between the internal and external versions here as well (see Exercise 9).

**Remark 5.** We will, as is commonly done, use the term *direct sum* and the symbol  $\oplus$  to mean either the internal or external direct sum. It should be clear from context which is meant. As there is a natural isomorphism between the two, it is easy to convert from a statement about one case to the corresponding statement about the other.

**Definition 6.** Let  $V_i$ ,  $i \in I$ , be an arbitrary collection of vector spaces over a field  $F$ . The *direct sum* of the collection of vector spaces  $\{V_i \mid i \in I\}$  is the set of all functions

$$f: I \longrightarrow \bigcup_{i \in I} V_i$$

which have the property that  $f(i) \in V_i$  and for which  $f(i) \neq 0$  for only finitely many values of  $i$  in  $I$ . This set becomes a vector space over  $F$  by defining operations in a pointwise fashion

$$\begin{aligned} (f_1 + f_2)(i) &= f_1(i) + f_2(i) \\ (af)(i) &= af(i) \end{aligned}$$

for  $i \in I$ ,  $f, f_1, f_2$  functions, and  $a \in F$ . We denote this vector space by

$$\bigoplus_{i \in I} V_i.$$

**Remark 7.** a. For  $I = \{1, 2\}$  there is an isomorphism between this definition and the earlier one given by

$$\phi : \bigoplus_{i \in I} V_i \longrightarrow V_1 \oplus V_2$$

where  $\phi(f) = (f(1), f(2))$ . That is, one of the functions  $f$  under consideration is completely determined by its values at 1 and 2. A similar statement holds for a finite number  $k$  of vector spaces  $V_i$ .

- b. It should now be clear why  $\bigoplus_{i \in I} V_i$  is a vector space over  $F$ : the sum and scalar multiple of functions with finite support have finite support, the zero vector is the function that has the value 0 in  $V_i$  for each  $i \in I$ , the negative of a function  $f$  has  $(-f)(i) = -f(i)$  and to verify that the definitions above make this a vector space requires checking equations for each  $i \in I$ , which are valid because  $V_i$  is a vector space over  $F$ .
- c. As before, the description of  $V$  as an internal direct sum is given by a collection of subspaces  $V_i$ ,  $i \in I$ , such that

$$V = \sum_i V_i$$

and that for each  $i$

$$V_i \cap \sum_{j \neq i} V_j = 0.$$

In general here although  $I$  may be infinite, an element is just a finite sum of elements from various  $V_i$ .

**Definition 8.** Let  $V_i$ ,  $i \in I$  be an arbitrary collection of vector spaces over a field  $F$ . The *direct product* of the collection of vector spaces  $\{V_i \mid i \in I\}$  is the set of all functions

$$f : I \longrightarrow \bigcup_{i \in I} V_i$$

which have the property that  $f(i) \in V_i$ . This set becomes a vector space over  $F$  by defining operations in a pointwise fashion

$$\begin{aligned} (f_1 + f_2)(i) &= f_1(i) + f_2(i) \\ (af)(i) &= af(i) \end{aligned}$$

for  $i \in I$ ,  $f, f_1, f_2$  functions, and  $a \in F$ . We denote this vector space by

$$\prod_{i \in I} V_i.$$

**Remark 9.** a. Note that the only difference between the definition of  $\prod_{i \in I} V_i$  and  $\bigoplus_{i \in I} V_i$  is the latter has the extra condition requiring all functions to have finite support (be non-zero for only finitely many  $i \in I$ ). Hence  $\bigoplus_{i \in I} V_i$  is a subset of  $\prod_{i \in I} V_i$ , and, in fact, is a subspace since the operations are defined by the same formulas.

- b. It is thus clear that the earlier remark about why the direct sum is a vector space applies in the case of direct product as well.
- c. If  $I$  is a finite set, then the two are identical. For that reason many times one will sometimes see the two concepts referred to by either term, and denoted with either symbol.
- d. For  $I$  infinite one should be extremely careful to distinguish between the two as they are definitely different. In the case where  $I$  is countably infinite (the same size as the set  $\mathbb{Z}$ ), an exercise in the section "Dual Spaces" outlines a proof.
- e. In either case one can define linear transformations

$$i_j : V_j \longrightarrow \prod_{i \in I} V_i$$

$$i_j : V_j \longrightarrow \bigoplus_{i \in I} V_i$$

which are given by

$$i_j(v_j) = f$$

where  $f$  is the function with values  $f(j) = v_j$  and  $f(i) = 0$  for  $i \neq j$ . There are also linear transformations

$$p_j : \prod_{i \in I} V_i \longrightarrow V_j$$

$$p_j : \bigoplus_{i \in I} V_i \longrightarrow V_j$$

given by

$$p_j(f) = f(j) .$$

The  $i_j$  are injective and the  $p_j$  are surjective as before and there is an analogous list of subspaces, formulas, etc. as in the earlier discussion. However, not every such formula necessarily makes sense (see exercises).

- f. Finally, when we begin to talk about what are called *universal mapping properties* we will introduce new definitions for sum and product for which the two (even in the finite case) will appear to be quite different.

We end this section with a question:

**Question 10.** Let  $W$ ,  $V_1$ ,  $V_2$  be vector spaces over the field  $F$ :

- a. How does one determine all linear transformations

$$W \longrightarrow V_1 \oplus V_2 ?$$

- b. How does one determine all linear transformations

$$V_1 \oplus V_2 \longrightarrow W ?$$

## Exercises

**SumProd 1.** Verify that addition and scalar multiplication as given in Definition 1 indeed defines a vector space.

**SumProd 2.** Verify that the maps  $p_1, p_2, i_1$  and  $i_2$  in the discussion following Definition 1 are linear transformations, that the given equations hold, that  $p_1$  and  $p_2$  are surjective and  $i_1$  and  $i_2$  are injective.

**SumProd 3.** Verify that  $i_1 \circ p_1 + i_2 \circ p_2$  is the identity map on  $V_1 \oplus V_2$ . What is the analogous equation for the direct sum  $V_1 \oplus V_2 \oplus \cdots \oplus V_k$ ? What about an infinite direct sum? Is there such an equation which is valid for an arbitrary direct product?

**SumProd 4.** Let  $T : V \rightarrow V$  be a linear transformation from a vector space  $V$  to itself, where  $V$  is a vector space over the field  $F$ . Assume  $T^2 = T$ . Show that  $V = \text{im } T \oplus \ker T$ .

**SumProd 5.** Let  $V$  be the vector space of all functions from  $\mathbb{R}$  to  $\mathbb{R}$ . Let  $V_e$  be the subset of even functions,  $f(-x) = f(x)$  and let  $V_o$  be the subset of odd functions,  $f(-x) = -f(x)$ .

- a. Prove that  $V_e$  and  $V_o$  are subspaces of  $V$ .
- b. Prove that  $V_e + V_o = V$ .
- c. Prove that  $V_e \cap V_o = \{0\}$ .
- d. What conclusion can you now make?
- e. Let  $F$  be an arbitrary field and  $V$  the vector space of all functions from  $F$  to  $F$ . Define  $V_e$  and  $V_o$  exactly the same way as above. Determine precisely when the exact same conclusions hold as held for  $\mathbb{R}$ .

**SumProd 6.** Let  $F$  be a field with characteristic unequal to 2 and let  $V$  be a vector space over  $F$ . Let  $T : V \rightarrow V$  be a linear transformation which satisfies  $T^2 = I$ , where  $I$  denotes the identity linear transformation. Define  $V^+ = \{v \in V \mid T(v) = +v\}$  and  $V^- = \{v \in V \mid T(v) = -v\}$ . Show that  $V = V^+ \oplus V^-$ . [Hint: Note the characteristic of  $F$ . Nothing other than elementary manipulations is required, nor allowed, to solve this exercise.]

**SumProd 7.** a. Show that the operation of direct sum is “commutative”: that is, there is a natural isomorphism

$$V_1 \oplus V_2 \approx V_2 \oplus V_1.$$

- b. Explain the difference between the vector spaces  $(V_1 \oplus V_2) \oplus V_3$  and  $V_1 \oplus (V_2 \oplus V_3)$ .
- c. Show that the operation of direct sum is “associative”: that is, there is a natural isomorphism

$$(V_1 \oplus V_2) \oplus V_3 \approx V_1 \oplus (V_2 \oplus V_3).$$

- d. Give a definition of the direct sum of  $k > 2$  vector spaces over  $F$  using  $k$ -tuples. Give an inductive definition assuming the case  $k = 2$  is given. Verify that the two definitions give isomorphic vector spaces.

**SumProd 8.** Let  $V$  be the internal direct sum of the subspaces  $V_1$  and  $V_2$ . Show that every element  $v$  of  $V$  can be written uniquely as a sum  $v = v_1 + v_2$  for some  $v_i \in V_i$ . Verify the assertion of Lemma 3.

**SumProd 9.** Let  $V$  be the internal direct sum of the subspaces  $V_i$ ,  $1 \leq i \leq k$ ,  $k > 2$ . State and prove the appropriate generalization of the preceding exercise. Give examples to show that pairwise intersection being 0 is not sufficient for  $k > 2$ . (See Remark 4.)

**SumProd 10.** Prove or disprove: Given a subspace  $W \subseteq V$ , there exists a unique subspace  $W'$  such that  $W + W' = V$  and  $W \cap W' = 0$ .

**SumProd 11.** Verify that Definition 8 indeed defines a vector space.

**SumProd 12.** Describe the direct sum and direct product of an “empty” collection of spaces (i.e., the index set  $I$  is empty).

**SumProd 13.** Let  $I = I_1 \sqcup I_2$  be a partition of the index set  $I$  into two disjoint subsets. Define two natural isomorphisms from  $\prod_{i \in I} V_i$  to  $(\prod_{i \in I_1} V_i) \oplus (\prod_{i \in I_2} V_i)$  and from  $\bigoplus_{i \in I} V_i$  to  $(\bigoplus_{i \in I_1} V_i) \oplus (\bigoplus_{i \in I_2} V_i)$ .

**SumProd 14.** Let  $I = \bigsqcup_{j \in J} I_j$  be a partition of the index set  $I$ . Are there any natural isomorphisms between either of  $\prod_{i \in I} V_i$  and  $\bigoplus_{i \in I} V_i$  with any of the vector spaces:

$$\begin{aligned} a) & \prod_{j \in J} \left( \prod_{i \in I_j} V_i \right) \\ b) & \bigoplus_{j \in J} \left( \prod_{i \in I_j} V_i \right) \\ c) & \prod_{j \in J} \left( \bigoplus_{i \in I_j} V_i \right) \\ d) & \bigoplus_{j \in J} \left( \bigoplus_{i \in I_j} V_i \right) \end{aligned}$$

Determine all of the natural inclusion maps between these four vector spaces (in the general case; that is, make no assumption on how large the sets  $J$  and  $I_j$  are).

**SumProd 15.** Show that if the index set  $I$  is infinite and all vector spaces  $V_i$  are non-trivial (i.e., they have elements different from 0), then the direct sum  $\bigoplus_{i \in I} V_i$  is a proper subspace of the direct product  $\prod_{i \in I} V_i$ . (Proving this requires using the Axiom of Choice.)

**SumProd 16.** Let  $V$  be a vector space over the field  $F$  and let  $S$  be a non-empty set. Interpret the examples  $F^S$ ,  $F^{(S)}$ ,  $V^S$ , and  $V^{(S)}$  from the section “Examples of Vector Spaces” in terms of the definitions given in this section.

**SumProd 17.** a. Show that  $F^n \oplus F^m \approx F^{n+m}$ .

b. Let the index set  $I = \{1, 2, \dots, n\}$  and each vector space  $V_i = F$ , the field  $F$ . Show that

$$\prod_{i=1}^n V_i \approx \bigoplus_{i=1}^n V_i \approx F^n.$$

c. Let the index set  $I$  be the set of natural numbers  $\mathbb{N}$  and each vector space  $V_i = F$ , the field  $F$ . Show that

$$\prod_{i \in I} V_i \approx F[[x]]$$

and

$$\bigoplus_{i \in I} V_i \approx F[x],$$

here  $F[x]$  is the vector space of formal polynomials over  $F$  and  $F[[x]]$  is the vector space of formal power series with coefficients in  $F$ . The isomorphisms above are as vector spaces over  $F$ .

d. Let the index set  $I$  be the set of real numbers  $\mathbb{R}$  and each vector space  $V_i = \mathbb{R}$ . What is  $\prod_{i \in I} V_i$ ? (That is, give a description (give an isomorphism) using some other notation we’ve previously discussed.)

**SumProd 18.** Let  $F = \mathbb{F}_p$  be the field of  $p$  elements with  $p$  a prime, and let  $V \subseteq F^n$  be a subspace. How many subspaces  $W$  such that  $V \cap W = 0$  and  $V + W = F^n$  exist?



## Math 4330 Homework Set 3

Due Monday, September 21, 2015

Keith Dennis Malott 524 255-4027 math4330@rkd.math.cornell.edu

TA: Gautam Gopal Krishnan 120 Malott Hall gk379@cornell.edu

**Read:** Notes on “Fields”, “Some Useful Definitions”, “Subobjects”, “Direct Sums and Products”, and “Equivalence Relations”.

Problems marked by  $\boxed{\text{box}}$  or  $\boxed{\star}$  are more challenging and may be turned in anytime during the semester. There will be several such problems assigned during the term. Please turn in *separately* from routine assignments – if incorrect or incomplete, they will be returned to you to complete correctly. Final deadline is Monday, Nov. 30, no exceptions.

Do the following problems from the handouts:

SumProd 3

SumProd 4

SumProd 5

SumProd 6

SumProd 16

and also

**Hw03 1.** a. Let  $W_1$  and  $W_2$  be subspaces of a vector space  $V$  such that their set-theoretic union is also a subspace. Prove that one of the spaces  $W_i$  is contained in the other.

$\boxed{\text{b.}}$  Prove this generalization of the first part. Assume that  $F$  is an infinite field. Let  $V_1, \dots, V_n$  be subspaces of a vector space  $V$  over  $F$ . Prove that  $V_1 \cup \dots \cup V_n$  is a subspace if and only if some  $V_i$  contains all the others. What happens if  $F$  is a finite field? [Give an exact statement relating  $n$  and the size of  $F$  in case that is necessary.]

**Hw03 2.** Let  $A$  be an abelian group (that is, addition  $+$  is defined and it satisfies exactly the same 4 properties that are satisfied by the addition of fields or vector spaces). For  $a \in A$  and  $n > 0$  an integer, we define  $n \cdot a = a + \cdots + a$  ( $n$  terms in sum) exactly as we did for fields. The *exponent* of  $A$  is the smallest positive integer  $n$ , if it exists, such that  $n \cdot a = 0$  for all  $a \in A$ . Otherwise we say the exponent of  $A$  is infinite. We write  $\exp(A) = n$  or  $\exp(A) = \infty$ . Note the similarity to (and difference from!) the definition of characteristic for a field. This exercise determines which abelian groups can be made into a vector space over the prime fields  $\mathbb{F}_p$  or  $\mathbb{Q}$ .

- a. Let  $F$  be a field and let  $V$  be a non-zero vector space over  $F$ . Are  $\exp(V)$  and  $\text{char}(F)$  related? If so, how? Prove your statement.
- b. Let  $\mathbb{F}_p$  be the finite field of integers modulo  $p$  for  $p$  a prime. Let  $A$  be an abelian group. Determine precisely when  $A$  can be made into a vector space over  $\mathbb{F}_p$ , define the scalar multiplication in that case, and verify that indeed  $A$  is a vector space over  $\mathbb{F}_p$  with the definition you have given.
- c. Let  $A$  be an abelian group. Give necessary and sufficient conditions on  $A$  such that it can be made into a vector space over  $\mathbb{Q}$ . [Hint: For  $n$  a non-zero integer and  $a \in A$ , what can one say about  $\frac{1}{n} \cdot a$ ?]