

## REAL ANALYSIS GENERAL EXAM FALL 2022

Solve as many problems as you can. Full solutions on a smaller number of problems will be worth more than partial solutions on several problems.

### Problem 1.

Let  $(X, \mu)$  be a  $\sigma$ -finite measure space and  $p \in [1, +\infty)$ . Let  $(f_n)_{n=1}^\infty$  be a sequence in  $L^p(X, \mu)$  and suppose that  $\|f_n\|_p \leq 1$ , and that  $f_n$  converges a.e. to a measurable function  $f$ . Show that  $\|f\|_p \leq 1$ .

Fatou's lemma

### Problem 2.

Let  $\mu$  be a Borel probability measure on  $\mathbb{R}$  without atoms. Suppose that  $E \subseteq \mathbb{R}$  is a Borel set with  $\mu(E) > 0$ . Show that there is a  $t \in \mathbb{R}$  with  $\mu(E \cap (-\infty, t)) = \frac{1}{2}\mu(E)$ .

### Problem 3.

Let  $X$  be a set equipped with a  $\sigma$ -algebra of sets  $\Sigma$ . Suppose that  $\mu, \nu: \Sigma \rightarrow [0, +\infty)$  are finite measures. Set  $\lambda = \mu + \nu$ . Let  $f: X \rightarrow \mathbb{R}$  be any  $\Sigma$ -measurable function so that

$$\nu(E) = \int_E f d\lambda$$

for all  $E \in \Sigma$ .

- (i) Show that  $0 \leq f \leq 1$   $\lambda$ -a.e.
- (ii) If  $F = \{x : f(x) = 1\}$ , show that  $\mu(F) = 0$ .
- (iii) If  $A \subseteq \{x : 0 \leq f(x) < 1\}$  and  $\mu(A) = 0$ , show that  $\nu(A) = 0$ .

### Problem 4.

Fix  $p \in [1, +\infty)$ . Let  $W^p([0, 1])$  consist of all absolutely continuous functions  $f: [0, 1] \rightarrow \mathbb{C}$  so that  $f' \in L^p([0, 1])$ . For  $f \in W^p([0, 1])$  define

$$\|f\| = |f(0)| + \|f'\|_p.$$

Show that  $\|\cdot\|$  is a norm which makes  $W^p([0, 1])$  into a Banach space. (You are allowed to use that  $L^p([0, 1])$  is a Banach space).

### Problem 5.

Let  $m$  be Lebesgue measure on  $\mathbb{R}$ . Let  $\Omega = \{1_E : E \subseteq \mathbb{R} \text{ is Borel and } m(E) < +\infty\}$  regarded as a subset of  $L^1(\mathbb{R})$  (recall that we identify two elements of  $L^1$  if they agree almost everywhere). Throughout this problem regard  $\Omega$  as a metric space equipped with the  $L^1$ -distance.

- (i) If  $a < b$  are real numbers, show that the function  $\Omega \rightarrow \mathbb{R}$  given by

$$1_E \mapsto m(E \cap [a, b])$$

is a continuous function.

- (ii) If  $a < b$  are real numbers, let  $U_{a,b}$  be the subset of  $\Omega$  consisting of all  $1_E$  where  $E \subseteq \mathbb{R}$  is Borel and

$$0 < m(E \cap [a, b]) < b - a.$$

Show that  $U_{a,b}$  is open and dense in  $\Omega$ .

(iii) Let  $D$  be the set of all  $1_E$  where  $E \subseteq \mathbb{R}$  is Borel and so that for every interval  $I$  of positive measure we have  $0 < m(E \cap I) < m(I)$ . Use (ii)

$$0 < m(E \cap I) < m(I).$$

Show that there is a countable collection  $\{U_j\}_{j \in J}$  of open and dense subsets of  $\Omega$  with

$$D \supseteq \bigcap_{j \in J} U_j.$$

## REAL ANALYSIS GENERAL EXAM FALL 2022

Solve as many problems as you can. Full solutions on a smaller number of problems will be worth more than partial solutions on several problems.

### Problem 1.

Compute

$$\lim_{n \rightarrow \infty} \int_0^\infty \frac{n \sin(x/n)}{x(1+x^2)} dx.$$

### Problem 2.

Fix  $a < b$  in  $\mathbb{R}$ . Recall that  $h: [a, b] \rightarrow \mathbb{C}$  is *absolutely continuous* if for every  $\varepsilon > 0$  there is a  $\delta > 0$  so that if  $((a_j, b_j))_{j=1}^k$  are disjoint intervals in  $[a, b]$  with  $\sum_{j=1}^k (b_j - a_j) < \delta$ , then  $\sum_{j=1}^k (f(b_j) - f(a_j)) < \varepsilon$ . For a Lipschitz function  $g: [a, b] \rightarrow \mathbb{C}$  we set

$$\|g\|_{Lip} = \sup_{x \neq y, x, y \in [a, b]} \frac{|g(x) - g(y)|}{|x - y|}.$$

- (a) Show that  $f: [a, b] \rightarrow \mathbb{C}$  is Lipschitz if and only if  $f$  is absolutely continuous and  $f' \in L^\infty([a, b])$ .
- (b) If  $f: [a, b] \rightarrow \mathbb{C}$  is Lipschitz, show that  $\|f\|_{Lip} = \|f'\|_\infty$ .

### Problem 3.

Let  $(X, \mu)$  be a  $\sigma$ -finite measure space. Show that if  $f, g \in L^1(X, \mu)$  with  $0 \leq f, g$  a.e., then

$$\|f - g\|_1 = \int_0^\infty \mu(\{x : f(x) > t\} \Delta \{x : g(x) > t\}) dt.$$

Here  $E \Delta F = E \setminus F \cup F \setminus E$  for sets  $E, F \subseteq X$ . Suggestion: it might be helpful to first show that for  $a, b \in [0, \infty)$  we have

$$|a - b| = \int_0^\infty |1_{(t, \infty)}(a) - 1_{(t, \infty)}(b)| dt.$$

**Note:** for this problem you may take for granted that the function  $X \times (0, \infty) \rightarrow \{0, 1\}$  given by  $(y, t) \mapsto 1_{\{x: f(x) > t\}}(y)$  and that the function  $t \mapsto \mu(\{x : f(x) > t\} \Delta \{x : g(x) > t\})$  are measurable functions.

### Problem 4.

Let  $(X, \Sigma)$  be a measurable space. Recall that if  $\eta$  is a signed measure on  $\Sigma$ , then  $|\eta| = \eta_1 + \eta_2$  where  $\eta_1, \eta_2$  are the unique nonnegative measures with  $\eta = \eta_1 - \eta_2$  and  $\eta_1 \perp \eta_2$ . Further,  $\|\eta\|_{TV} = |\eta|(X)$ . Suppose that  $\mu, \nu$  are signed measures on  $\Sigma$ , that  $\|\mu\|_{TV}, \|\nu\|_{TV} < +\infty$  and that  $|\mu|, |\nu|$  are mutually singular.

- (a) If  $\mu = \mu_1 - \mu_2$ ,  $\nu = \nu_1 - \nu_2$  with  $\mu_i, \nu_j$  nonnegative measures and  $\mu_1 \perp \mu_2$ ,  $\nu_1 \perp \nu_2$ , show that  $\mu_i \perp \nu_j$  for all  $i, j \in \{1, 2\}$ .
- (b) Show that

$$\|\mu + \nu\|_{TV} = \|\mu\|_{TV} + \|\nu\|_{TV}.$$

**Problem 5.**

- (a) For  $f \in L^1([0, 1])$ , set  $L_f$  be the set of  $x \in [0, 1]$  so that

$$\lim_{r \rightarrow 0} \frac{1}{2r} \int_{(x-r, x+r)} |f(y) - f(x)| dy = 0.$$

State the conclusion of the Lebesgue's differentiation theorem for  $L_f$ .

- (b) For  $n \in \mathbb{N}$ , and  $0 \leq j \leq 2^n - 1$ , set  $I_{n,j} = [j2^{-n}, (j+1)2^{-n})$ . For  $f \in L^1([0, 1])$ , define

$$E_n f = \sum_{j=0}^{2^n-1} \left( \frac{1}{m(I_{n,j})} \int_{I_{n,j}} f(t) dt \right) 1_{I_{n,j}}.$$

Show that

$$\lim_{n \rightarrow \infty} (E_n f)(x) = f(x) \text{ for almost every } x \in [0, 1].$$

**Instructions.** 2 hours. Closed book examination. Be neat in your presentation. When invoking a theorem from previous courses, name the theorem and thoroughly check its hypotheses. You must solve a significant portion of each of the three problems in order to pass the exam.

1. Let  $(X, \mathcal{M}, \mu)$  be a measure space. Let  $(f_n)$  be a sequence of nonnegative functions in  $L^1(X, \mathcal{M}, \mu)$  and let  $f$  be a nonnegative function in  $L^1(X, \mathcal{M}, \mu)$ . Suppose that

$$\int_X f_n \, d\mu \longrightarrow \int_X f \, d\mu$$

and that  $f_n \rightarrow f$  pointwise. Prove that  $f_n$  converges to  $f$  in  $L^1(X, \mathcal{M}, \mu)$ . Hint: consider  $g_n = \min(f, f_n)$ .

2. Let  $(X, \mathcal{M}, \mu)$  be a measure space. Let  $p \in [1, \infty)$ .

(a) Let  $(f_n)$  be a sequence of functions in  $L^p(X, \mathcal{M}, \mu)$  and let  $f$  be a function in  $L^p(X, \mathcal{M}, \mu)$ . Suppose that  $f_n$  converges to  $f$  in  $L^p(X, \mathcal{M}, \mu)$ . Prove that there exists a subsequence  $(f_{n_k})$  such that for  $\mu$ -almost all  $x$ ,  $\lim_{k \rightarrow \infty} f_{n_k}(x) = f(x)$ . Hint: remember the proof of completeness of  $L^p$ .

(b) Let  $h$  be a measurable function on  $X$ . Let

$$D = \{f \in L^p(X, \mathcal{M}, \mu) \mid hf \in L^p(X, \mathcal{M}, \mu)\}.$$

Let  $(f_n)$  be a sequence of elements of  $D$ , and let  $f, g \in L^p(X, \mathcal{M}, \mu)$  be such that  $f_n$  converges to  $f$  in  $L^p$ , and  $hf_n$  converges to  $g$  in  $L^p$ . Show that  $f \in D$  and  $g = hf$ .

3. For  $\mu$  a Borel probability measure on  $\mathbb{R}$ , we will denote by  $\hat{\mu}$  the function  $\mathbb{R} \rightarrow \mathbb{C}$  given by

$$\hat{\mu}(t) = \int_{\mathbb{R}} e^{itx} d\mu(x).$$

We will also adopt the notational convention  $\text{sinc}(x) = \frac{\sin x}{x}$  if  $x \neq 0$  and  $\text{sinc}(0) = 1$ .

(a) Show that  $\hat{\mu}$  is a bounded continuous function.

(b) Let  $\delta > 0$ . Show that

$$\frac{1}{2\delta} \int_{-\delta}^{\delta} (1 - \text{Re}(\hat{\mu}(t))) \, dt = \int_{\mathbb{R}} (1 - \text{sinc}(\delta x)) \, d\mu(x).$$

(c) Show that for all  $u \in \mathbb{R}$ ,

$$1 - \text{sinc}(u) \geq \frac{1}{2} \chi_{(-\infty, -2) \cup (2, \infty)}(u),$$

and deduce that

$$\mu(\{x \in \mathbb{R} \mid |x| > 2\delta^{-1}\}) \leq \frac{1}{\delta} \int_{-\delta}^{\delta} (1 - \operatorname{Re}(\widehat{\mu}(t))) \, dt .$$

(d) Let  $\mu_n$  be a sequence of Borel probability measures on  $\mathbb{R}$ . Suppose that for all  $t$ , the limit  $\Phi(t) = \lim_{n \rightarrow \infty} \widehat{\mu_n}(t)$  exists and that the resulting function  $\Phi(t)$  is continuous at  $t = 0$ . Prove that for all  $\epsilon > 0$ , there exists a compact set  $K$  inside  $\mathbb{R}$  such that, for all  $n$ ,  $\mu_n(K) \geq 1 - \epsilon$ .

# Real analysis Qualifying exam, January 2021

## DO NOT WRITE YOUR NAME ON YOUR WORK

My cellphone in case of zoom disconnection: \*\*\*

In order to receive the full credit for a problem, a detailed argument (rather than a sketch of the proof) is needed. Whenever applying one of the standard theorems, please indicate that clearly. Full solutions on a smaller number of problems will be worth more than partial solutions on more problems.

- 1.** Let  $f_n$ ,  $n \geq 1$ , and  $f$  be measurable functions on a space  $(\Omega, \mathcal{F}, \mu)$ , such that  $f_n \rightarrow f$  in measure. Does this imply that there exists a measurable set  $A \subseteq \Omega$  with  $\mu(\Omega \setminus A) = 0$  such that  $f_n(x) \rightarrow f(x)$  for all  $x \in A$ ?  
If yes, prove this. If no, give a counterexample.

- 2.** Let  $B$  be a measurable subset of the two-dimensional plane such that the intersection of  $B$  with every vertical line is finite or countable. Find  $\mu(B)$ , where  $\mu$  is the two-dimensional Lebesgue measure. Justify your answer.

- 3.** Let  $(\Omega, \mathcal{F})$  be a measurable space, and  $\mu, \nu, \rho$  be three finite positive measures on  $(\Omega, \mathcal{F})$  such that  $\mu \ll \nu$  (i.e.,  $\mu$  is absolutely continuous with respect to  $\nu$ ). Show that there exists a measurable function  $f$  on  $\Omega$  such that for all  $E \in \mathcal{F}$  we have

$$\mu(E) = \int_E f \, d\nu + \int_E (f - 1) \, d\rho.$$

(Hint: use Radon-Nikodym's Theorem)

- 4.** Let  $f, g$  be nonnegative measurable functions on  $[0, 1]$ , and  $a, b, c, d \geq 0$  be arbitrary nonnegative numbers. Show that then

$$\left( ac + bd + \int_0^1 f(x)g(x) \, dx \right)^3 \leq \left( a^3 + b^3 + \int_0^1 (f(x))^3 \, dx \right) \left( c^{3/2} + d^{3/2} + \int_0^1 (g(x))^{3/2} \, dx \right)^2.$$

Partial credit is given for proving the inequality in the particular case  $a = b = c = d = 0$ .

- 5.** Let  $f(x)$  be a continuous function on  $[0, 1]$ . Show that for every  $\varepsilon > 0$  there exists  $n \in \mathbb{Z}_{\geq 0}$  and constants  $a_0, a_1, \dots, a_n \in \mathbb{R}$  such that for the differential operator

$$D := \sum_{k=0}^n a_k \left( \frac{d}{dx} \right)^k = a_0 + a_1 \frac{d}{dx} + a_2 \left( \frac{d}{dx} \right)^2 + \dots + a_n \left( \frac{d}{dx} \right)^n$$

we have  $\left| f(x) - e^{x^2} (D e^{-x^2}) \right| < \varepsilon$  for all  $x \in [0, 1]$ . Here  $e^{x^2} (D e^{-x^2})$  is the function obtained by applying  $D$  to  $e^{-x^2}$  and after that multiplying the result by  $e^{x^2}$ .

# Real analysis Qualifying exam, August 2020

Make sure that you have signed the Honor Pledge on Collab.

In order to receive the full credit for a problem, a detailed argument (rather than a sketch of the proof) is needed. Whenever applying one of the standard theorems, please indicate that clearly.

**Cantor function increasing and continuous such that derivative is 0 a.e.**

- 1.** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be continuous, almost everywhere differentiable, and such that  $f'(x) = 1$  almost everywhere. (Both “almost everywhere” properties are assumed with respect to the Lebesgue measure on  $\mathbb{R}$ .) Does this imply that  $f(2) - f(1) = 1$ ?

If yes, prove this. If no, give a counterexample.

- 2.** Is every open set in  $\mathbb{R}^2$  a countable union of closed sets?

If yes, prove this. If no, give a counterexample.

- 3.** Let  $\mathcal{H}$  be a separable complex Hilbert space with basis (complete orthonormal system)  $f_1, f_2, f_3, \dots$ . Define a linear operator  $P$  in  $\mathcal{H}$  by setting

$$P(f_n) = f_{n+1}, \quad n = 1, 2, \dots$$

(a) Find the adjoint  $P^*$  to  $P$ .

(b) Find the operators  $PP^*$  and  $P^*P$ .

- 4.** Let  $(X, \mathcal{F}, \mu)$  be a measure space with  $\mu(X) = 1$ . Let  $f_n: X \rightarrow \mathbb{R}$  be measurable functions such that for all  $t \in \mathbb{R}$ ,

$$\lim_{n \rightarrow +\infty} \mu(x: f_n(x) \leq t) = \begin{cases} 0, & t < 0; \\ 1, & t \geq 0. \end{cases}$$

Show that  $f_n \rightarrow 0$  in measure.

- 5.** Show that the operator

$$(Tf)(x) := \int_0^\infty \frac{f(y)}{x+y} dy$$

is bounded in the space  $L^p(\mathbb{R}_{\geq 0})$  for all  $1 < p < +\infty$ .



## Real analysis general exam, January 2020

1. Let  $\mu$  be the Lebesgue measure on  $\mathbb{R}$ . For a Lebesgue measurable set  $A \subset [0, 1]$ , is it true that

- (a)  $\mu(A) = \sup_{U \subset A, U \text{ open}} \mu(U)$  ? If true, prove this. If false, give a counterexample.
- (b)  $\mu(A) = \inf_{U \supset A, U \text{ open}} \mu(U)$  ? If true, prove this. If false, give a counterexample.

2. Find a polynomial  $P(x)$  of degree at most 3 such that  $\int_{-1}^1 |x^4 - P(x)|^2 dx$  is minimal.

3. Let  $X$  be a compact metric space, and  $C(X)$  be the space of all real-valued continuous functions on  $X$  with the supremum norm. Assume that the subset  $\mathcal{A} \subset C(X)$  satisfies the following properties:

- (algebra) For all  $f, g \in \mathcal{A}$  and  $\alpha, \beta \in \mathbb{R}$  we have  $\alpha f + \beta g \in \mathcal{A}$  and  $fg \in \mathcal{A}$ .
- (separates points) For any  $x \neq y$  from  $X$  there exists a function  $f \in \mathcal{A}$  such that  $f(x) \neq f(y)$ .

This question has two parts:

- (a) Show by example that  $\mathcal{A}$  need not be dense in  $C(X)$ , explicitly checking all the properties of your example  $\mathcal{A}$ .
- (b) In order to conclude that  $\mathcal{A}$  is dense by Stone-Weierstrass Theorem, what additional condition(s) should be added?

4. Let  $\mu$  be a measure on  $(\mathbb{R}, \mathcal{B})$ , where  $\mathcal{B}$  is the Borel  $\sigma$ -algebra. Let  $\mu(\mathbb{R}) = 1$ . Next, let  $\mathcal{F} \subset \mathcal{B}$  be the sub- $\sigma$ -algebra of symmetric Borel sets, that is,  $\mathcal{F}$  generated by all intervals of the form  $(-a, a)$  with  $a > 0$ .

Let  $f \in L^1(\mathbb{R}, \mathcal{B}, \mu)$ . Find a function  $g$  such that:

- (a)  $g \in L^1(\mathbb{R}, \mathcal{F}, \mu)$  (in particular,  $g$  is  $\mathcal{F}$ -measurable).
- (b) For all  $E \in \mathcal{F}$  we have  $\int_E g d\mu = \int_E f d\mu$ .

5. Let  $\mu$  be a finite measure on some measurable space  $(X, \mathcal{F})$ .

Show that a sequence of  $\mathcal{F}$ -measurable functions  $f_n$  converges to a function  $f$  in measure if and only if  $\int_X \min\{1, |f_n - f|\} \mu(dx) \rightarrow 0$  as  $n \rightarrow +\infty$ .

## Real analysis Qualifying exam, August 2019

1. Let  $\mathcal{C}$  be the Cantor set on  $[0, 1]$ . Recall that it is obtained by iteratively deleting the open middle third:  $(\frac{1}{3}, \frac{2}{3})$ , then  $(\frac{1}{9}, \frac{2}{9}) \cup (\frac{7}{9}, \frac{8}{9})$ , and so on.

(a) Show that  $\mathcal{C} + \mathcal{C} := \{a + b : a, b \in \mathcal{C}\}$  is the full segment  $[0, 2]$ .

(b) Find two sets  $A, B \subset \mathbb{R}$ , each of which is closed and has Lebesgue measure zero, such that  $A + B = \{a + b : a \in A, b \in B\}$  is the full line  $\mathbb{R}$ .

2. Does there exist a measure space  $(X, \mathcal{F}, \mu)$  with a finite measure  $\mu$ , and a sequence of  $\mu$ -measurable functions  $\{f_n\}_{n=1,2,\dots}$  on  $X$  such that:

- $f_n(x) \geq 0$  for all  $n, x$ ;
- $f_n(x) \rightarrow 0$  as  $n \rightarrow +\infty$  for all  $x$ ;
- $\int f_n(x) \mu(dx) \rightarrow 0$  as  $n \rightarrow +\infty$ ;
- $\Phi(x) := \sup_n f_n(x)$  has infinite integral?

If yes, give an example of such a sequence  $\{f_n\}$ . If no, give a proof of nonexistence.

3. Let  $\mu$  be a signed Borel measure on  $\mathbb{R}^n$  which is bounded on bounded sets. Suppose that  $\int f d\mu = 0$  for all continuous functions  $f$  with bounded support. Show that then  $\mu = 0$ .

4. Let  $L^1(\mathbb{R})$  be the space of Lebesgue integrable functions on  $\mathbb{R}$ . For a positive function  $f \in L^1(\mathbb{R})$  show that the function  $\frac{1}{f(x)}$  does not belong to  $L^1(\mathbb{R})$ .

(Hint: look at the function  $1 = f^{1/2} f^{-1/2}$ .)

5. Applying the Gram-Schmidt orthogonalization to  $1, x, x^2, \dots$  in the Hilbert space  $L^2([-1, 1])$  (with Lebesgue measure), one gets the Legendre polynomials  $L_n(x)$ ,  $n = 0, 1, 2, \dots$

(a) Show that the Legendre polynomials form a basis (= complete orthogonal system) in the Hilbert space  $L^2([-1, 1])$

(b) Show that the Legendre polynomials are given by the formula  $L_n(x) = c_n \frac{d^n}{dx^n} (x^2 - 1)^n$  (you do not need to specify  $c_n$ ).

(Hint: employ integration by parts.)