

Class 8: PCA mini project

Sion Kang (PID: A17052234)

2025-01-30

Today we will do a complete analysis of some breast cancer biopsy data but first let's revisit the main PCA function in R `prcomp()` and see what `'scale=TRUE'`

```
head(mtcars)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Mazda RX4	21.0	6	160	110	3.90	2.620	16.46	0	1	4	4
Mazda RX4 Wag	21.0	6	160	110	3.90	2.875	17.02	0	1	4	4
Datsun 710	22.8	4	108	93	3.85	2.320	18.61	1	1	4	1
Hornet 4 Drive	21.4	6	258	110	3.08	3.215	19.44	1	0	3	1
Hornet Sportabout	18.7	8	360	175	3.15	3.440	17.02	0	0	3	2
Valiant	18.1	6	225	105	2.76	3.460	20.22	1	0	3	1

Find the mean value per column of this dataset

```
apply(mtcars, 2, mean)
```

mpg	cyl	disp	hp	drat	wt	qsec
20.090625	6.187500	230.721875	146.687500	3.596563	3.217250	17.848750
vs	am	gear	carb			
0.437500	0.406250	3.687500	2.812500			


```
apply(mtcars, 2, sd)
```

mpg	cyl	disp	hp	drat	wt
6.0269481	1.7859216	123.9386938	68.5628685	0.5346787	0.9784574
qsec	vs	am	gear	carb	
1.7869432	0.5040161	0.4989909	0.7378041	1.6152000	

It is clear “disp” and “hp” have the highest mean values and the highest standard deviation. They will likely dominate any analysis I do on this dataset. Let’s see

```
pc.noscale <- prcomp(mtcars, scale=FALSE)
pc.scale <- prcomp(mtcars, scale=TRUE)
```

```
biplot(pc.noscale)
```



```
pc.noscale$rotation[,1]
```


mpg	cyl	disp	hp	drat	wt
-0.038118199	0.012035150	0.899568146	0.434784387	-0.002660077	0.006239405
qsec	vs	am	gear	carb	
-0.006671270	-0.002729474	-0.001962644	-0.002604768	0.005766010	

Plot the loadings

```
library(ggplot2)


r1 <- as.data.frame(pc.noscale$rotation)
r1$names <- rownames(pc.noscale$rotation)
```

```
ggplot(r1) +
  aes(PC1, names)+
  geom_col()
```




```
r2 <- as.data.frame(pc.scale$rotation)
r2$names <- rownames(pc.scale$rotation)

ggplot(r2) +
  aes(PC1, names)+
  geom_col()
```



```
biplot(pc.scale)
```



Take-home: Generally we always want to set `scale=TRUE` when we do this type

of analysis to avoid our analysis being dominated by individual variables with the largest variance just due to their unit of measurement.

FNA breast cancer data

Load the data into R.

```
wisc.df <- read.csv("WisconsinCancer.csv", row.names = 1)
head(wisc.df)
```

	diagnosis	radius_mean	texture_mean	perimeter_mean	area_mean
842302	M	17.99	10.38	122.80	1001.0
842517	M	20.57	17.77	132.90	1326.0
84300903	M	19.69	21.25	130.00	1203.0
84348301	M	11.42	20.38	77.58	386.1
84358402	M	20.29	14.34	135.10	1297.0
843786	M	12.45	15.70	82.57	477.1

	smoothness_mean	compactness_mean	concavity_mean	concave.points_mean
842302	0.11840	0.27760	0.3001	0.14710
842517	0.08474	0.07864	0.0869	0.07017
84300903	0.10960	0.15990	0.1974	0.12790
84348301	0.14250	0.28390	0.2414	0.10520
84358402	0.10030	0.13280	0.1980	0.10430
843786	0.12780	0.17000	0.1578	0.08089

	symmetry_mean	fractal_dimension_mean	radius_se	texture_se	perimeter_se
842302	0.2419	0.07871	1.0950	0.9053	8.589
842517	0.1812	0.05667	0.5435	0.7339	3.398
84300903	0.2069	0.05999	0.7456	0.7869	4.585
84348301	0.2597	0.09744	0.4956	1.1560	3.445
84358402	0.1809	0.05883	0.7572	0.7813	5.438
843786	0.2087	0.07613	0.3345	0.8902	2.217

	area_se	smoothness_se	compactness_se	concavity_se	concave.points_se
842302	153.40	0.006399	0.04904	0.05373	0.01587
842517	74.08	0.005225	0.01308	0.01860	0.01340
84300903	94.03	0.006150	0.04006	0.03832	0.02058
84348301	27.23	0.009110	0.07458	0.05661	0.01867
84358402	94.44	0.011490	0.02461	0.05688	0.01885
843786	27.19	0.007510	0.03345	0.03672	0.01137

	symmetry_se	fractal_dimension_se	radius_worst	texture_worst
842302	0.03003	0.006193	25.38	17.33
842517	0.01389	0.003532	24.99	23.41

84300903	0.02250	0.004571	23.57	25.53
84348301	0.05963	0.009208	14.91	26.50
84358402	0.01756	0.005115	22.54	16.67
843786	0.02165	0.005082	15.47	23.75
	perimeter_worst	area_worst	smoothness_worst	compactness_worst
842302	184.60	2019.0	0.1622	0.6656
842517	158.80	1956.0	0.1238	0.1866
84300903	152.50	1709.0	0.1444	0.4245
84348301	98.87	567.7	0.2098	0.8663
84358402	152.20	1575.0	0.1374	0.2050
843786	103.40	741.6	0.1791	0.5249
	concavity_worst	concave.points_worst	symmetry_worst	
842302	0.7119	0.2654	0.4601	
842517	0.2416	0.1860	0.2750	
84300903	0.4504	0.2430	0.3613	
84348301	0.6869	0.2575	0.6638	
84358402	0.4000	0.1625	0.2364	
843786	0.5355	0.1741	0.3985	
	fractal_dimension_worst			
842302	0.11890			
842517	0.08902			
84300903	0.08758			
84348301	0.17300			
84358402	0.07678			
843786	0.12440			

Q1. How many observations are in this dataset?

```
nrow(wisc.df)
```

```
[1] 569
```

Q2. How many of the observations have a malignant diagnosis?

```
sum(wisc.df$diagnosis == "M")
```

```
[1] 212
```

The `table()` function is super useful here

```
table(wisc.df$diagnosis)
```

```
  B    M  
357 212
```

Q3. How many variables/features in the data are suffixed with `_mean`?

```
ncol(wisc.df)
```

```
[1] 31
```

```
colnames(wisc.df)
```

```
[1] "diagnosis"           "radius_mean"  
[3] "texture_mean"        "perimeter_mean"  
[5] "area_mean"           "smoothness_mean"  
[7] "compactness_mean"    "concavity_mean"  
[9] "concave.points_mean" "symmetry_mean"  
[11] "fractal_dimension_mean" "radius_se"  
[13] "texture_se"          "perimeter_se"  
[15] "area_se"             "smoothness_se"  
[17] "compactness_se"      "concavity_se"  
[19] "concave.points_se"   "symmetry_se"  
[21] "fractal_dimension_se" "radius_worst"  
[23] "texture_worst"       "perimeter_worst"  
[25] "area_worst"          "smoothness_worst"  
[27] "compactness_worst"   "concavity_worst"  
[29] "concave.points_worst" "symmetry_worst"  
[31] "fractal_dimension_worst"
```

A useful function for this is `grep()`

```
length( grep("_mean", colnames(wisc.df)) )
```

```
[1] 10
```

Before we go any further we need to exclude the diagnosis column from any future analysis - this tells us whether a sample is cancer or non-cancer.


```
diagnosis <- as.factor(wisc.df$diagnosis)
head(diagnosis)
```

```
[1] M M M M M M
Levels: B M
```

```
wisc.data <- wisc.df[, -1]
```

Let's see if we can cluster the wisc.data to find some structure in the dataset.

```
hc <- hclust( dist(wisc.data) )
plot(hc)
```



Principal Component Analysis (PCA)

```
wisc.pr <- prcomp( wisc.data, scale=T )
summary(wisc.pr)
```


Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

Q4. From your results, what proportion of the original variance is captured by the first principal components (PC1)?

44.27% of the original variance is capture by PC1.

Q5. How many principal components (PCs) are required to describe at least 70% of the original variance in the data?

PC1, PC2, PC3 (3 PCs) are required to describe at least 70% of the original variance.

Q6. How many principal components (PCs) are required to describe at least 90% of the original variance in the data?


PC1-7 are required to describe at least 90% of the original variance.

```
biplot(wisc.pr)
```


84300903	-5.728855	1.074229	-0.5512625	0.9112808	0.1769302	0.54097615
84348301	-7.116691	-10.266556	-3.2299475	0.1524129	2.9582754	3.05073750
84358402	-3.931842	1.946359	1.3885450	2.9380542	-0.5462667	-1.22541641
843786	-2.378155	-3.946456	-2.9322967	0.9402096	1.0551135	-0.45064213
	PC7	PC8	PC9	PC10	PC11	PC12
842302	2.15747152	0.39805698	-0.15698023	-0.8766305	-0.2627243	-0.8582593
842517	0.01334635	-0.24077660	-0.71127897	1.1060218	-0.8124048	0.1577838
84300903	-0.66757908	-0.09728813	0.02404449	0.4538760	0.6050715	0.1242777
84348301	1.42865363	-1.05863376	-1.40420412	-1.1159933	1.1505012	1.0104267
84358402	-0.93538950	-0.63581661	-0.26357355	0.3773724	-0.6507870	-0.1104183
843786	0.49001396	0.16529843	-0.13335576	-0.5299649	-0.1096698	0.0813699
	PC13	PC14	PC15	PC16	PC17	
842302	0.10329677	-0.690196797	0.601264078	0.74446075	-0.26523740	
842517	-0.94269981	-0.652900844	-0.008966977	-0.64823831	-0.01719707	
84300903	-0.41026561	0.016665095	-0.482994760	0.32482472	0.19075064	
84348301	-0.93245070	-0.486988399	0.168699395	0.05132509	0.48220960	
84358402	0.38760691	-0.538706543	-0.310046684	-0.15247165	0.13302526	
843786	-0.02625135	0.003133944	-0.178447576	-0.01270566	0.19671335	
	PC18	PC19	PC20	PC21	PC22	
842302	-0.54907956	0.1336499	0.34526111	0.096430045	-0.06878939	
842517	0.31801756	-0.2473470	-0.11403274	-0.077259494	0.09449530	
84300903	-0.08789759	-0.3922812	-0.20435242	0.310793246	0.06025601	
84348301	-0.03584323	-0.0267241	-0.46432511	0.433811661	0.20308706	
84358402	-0.01869779	0.4610302	0.06543782	-0.116442469	0.01763433	
843786	-0.29727706	-0.1297265	-0.07117453	-0.002400178	0.10108043	
	PC23	PC24	PC25	PC26	PC27	
842302	0.08444429	0.175102213	0.150887294	-0.201326305	-0.25236294	
842517	-0.21752666	-0.011280193	0.170360355	-0.041092627	0.18111081	
84300903	-0.07422581	-0.102671419	-0.171007656	0.004731249	0.04952586	
84348301	-0.12399554	-0.153294780	-0.077427574	-0.274982822	0.18330078	
84358402	0.13933105	0.005327110	-0.003059371	0.039219780	0.03213957	
843786	0.03344819	-0.002837749	-0.122282765	-0.030272333	-0.08438081	
	PC28	PC29	PC30			
842302	-0.0338846387	0.045607590	0.0471277407			
842517	0.0325955021	-0.005682424	0.0018662342			
84300903	0.0469844833	0.003143131	-0.0007498749			
84348301	0.0424469831	-0.069233868	0.0199198881			
84358402	-0.0347556386	0.005033481	-0.0211951203			
843786	0.0007296587	-0.019703996	-0.0034564331			


Plot of PC1 vs PC2 the first two columns

```
plot(wisc.pr$x[,1], wisc.pr$x[,2],col=diagnosis, xlab="PC1", ylab="PC2")
```



Q8. Generate a similar plot for principal components 1 and 3. What do you notice about these plots?

```
plot(wisc.pr$x[,1], wisc.pr$x[,3],col=diagnosis, xlab="PC1", ylab="PC3")
```




There is more overlapping occurring between the malignant and benign diagnosis groups. This is because PC3 explains less variance than PC2.

Make a ggplot version of this score plot

```
pc <- as.data.frame(wisc.pr$x)

ggplot(pc) +
  aes(PC1, PC2, col=diagnosis) +
  geom_point()
```




Variance explained

```
# Calculate variance of each component
pr.var <- wisc.pr$sdev^2
head(pr.var)
```


```
[1] 13.281608  5.691355  2.817949  1.980640  1.648731  1.207357
```

```
# Variance explained by each principal component: pve
pve <- pr.var / sum(pr.var)

# Plot variance explained for each principal component
plot(pve, xlab = "Principal Component",
     ylab = "Proportion of Variance Explained",
     ylim = c(0, 1), type = "o")
```



```
# Alternative scree plot of the same data, note data driven y-axis
barplot(pve, ylab = "Precent of Variance Explained",
        names.arg=paste0("PC",1:length(pve)), las=2, axes = FALSE)
axis(2, at=pve, labels=round(pve,2)*100 )
```



Communicating PCA results

Q9. For the first principal component, what is the component of the loading vector (i.e. `wisc.pr$rotation[,1]`) for the feature `concave.points_mean`?

```
wisc.pr$rotation["concave.points_mean",1]
```

```
[1] -0.2608538
```

Q10. What is the minimum number of principal components required to explain 80% of the variance of the data?

```
summary(wisc.pr)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624

Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

PC1-5 are needed to explain 80% of the variance of the data.

Hierarchical clustering

```
# Scale the wisc.data data using the "scale()" function
data.scaled <- scale(wisc.data)
```

```
data.dist <- dist(data.scaled)

wisc.hclust <- hclust(data.dist, method="complete")
wisc.hclust
```

Call:

```
hclust(d = data.dist, method = "complete")
```


```
Cluster method   : complete
Distance         : euclidean
Number of objects: 569
```

Results of hierarchical clustering

Q11. Using the `plot()` and `abline()` functions, what is the height at which the clustering model has 4 clusters?

At $h=19$

```
plot(wisc.hclust)
abline(h=19, col="red", lty=2)
```



```
data.dist
hclust (*, "complete")
```

Selecting number of clusters

Use `cutree()` to cut the tree so that it has 4 clusters.

```
wisc.hclust.clusters <- cutree(wisc.hclust, k=4)
table(wisc.hclust.clusters, diagnosis)
```

	diagnosis	
wisc.hclust.clusters	B	M
1	12	165
2	2	5
3	343	40
4	0	2

Q12. Can you find a better cluster vs diagnoses match by cutting into a different number of clusters between 2 and 10?

```
wisc.hclust.clusters.fxn <- function(x) {
  table(cutree(wisc.hclust, k=x), diagnosis)
}

wisc.hclust.clusters.fxn(2)
```

```
diagnosis
  B   M
1 357 210
2   0   2
```

```
wisc.hclust.clusters.fxn(3)
```

```
diagnosis
  B   M
1 355 205
2   2   5
3   0   2
```

```
wisc.hclust.clusters.fxn(5)
```

```
diagnosis
  B   M
1  12 165
2   0   5
3 343  40
4   2   0
5   0   2
```

No, regardless of the number of clusters made, there are no better cluster vs diagnoses matches. With 2 or 3 clusters, nearly all the malignant and benign diagnoses are in one cluster together while with a greater number of clusters, one diagnosis is separated in multiple clusters.

Using different methods

Q13. Which method gives your favorite results for the same data.dist dataset?
Explain your reasoning.


Complete, as it uses the largest of all pair-wise similarities and would lead to tight, well-separated groups.

Combining methods

Clustering in PC space

Class example

```
hc <- hclust( dist( wisc.pr$x[,1:2]), method="ward.D2")  
  
plot(hc)  
abline(h=70, col="red")
```



```
dist(wisc.pr$x[, 1:2])  
hclust (*, "ward.D2")
```

```
grps <- cutree(hc, h=70)  
table(grps)
```

```
grps  
  1  2  
195 374
```

```
table(diagnosis)
```

```
diagnosis
  B   M
357 212
```

Cross-table to see how my clustering groups correspond to the expert diagnosis vector of M and B values

```
table(grps, diagnosis)
```

```
      diagnosis
grps  B   M
1    18 177
2   339  35
```

Positive => cancer M Negative => non-cancerous B

True = cluster/grp 1 False= grp 2


True Positive 177 False Positive 18 True Negative 339 False Negative 35

sensitivity 177/212

Clustering on PCA results

```
wisc.pr.hclust <- hclust( dist( wisc.pr$x[,1:7]), method="ward.D2")
plot(wisc.pr.hclust)
```

Cluster Dendrogram




```
dist(wisc.pr$x[, 1:7])
hclust (*, "ward.D2")
```

```
grps <- cutree(wisc.pr.hclust, k=2)
table(grps, diagnosis)
```

```
      diagnosis
grps   B    M
  1   28 188
  2  329  24
```

```
plot(wisc.pr$x[,1:2], col=grps)
```



Cut the hierarchical clustering model into 2 clusters

```
wisc.pr.hclust.clusters <- cutree(wisc.pr.hclust, k=2)
```

Q15. How well does the newly created model with four clusters separate out the two diagnoses?

```
table(wisc.pr.hclust.clusters, diagnosis)
```

	diagnosis	
wisc.pr.hclust.clusters	B	M
1	28	188
2	329	24

Positive => cancer M Negative => non-cancerous B

True Positive 188 False Positive 28 True Negative 329 False Negative 24

`wisc.pr.hclust.clusters` with two clusters was able to separate clusters based on diagnosis with pretty high accuracy, with cluster 1 largely corresponding to malignant and cluster 2 to benign. Precision is also seen when comparing the true positive/negative values with the false positive/negative values.

Q16. How well do the k-means and hierarchical clustering models you created in previous sections (i.e. before PCA) do in terms of separating the diagnoses? Again, use the `table()` function to compare the output of each model (`wisc.km$cluster` and `wisc.hclust.clusters`) with the vector containing the actual diagnoses.

```
table(wisc.hclust.clusters, diagnosis)
```

	diagnosis	
wisc.hclust.clusters	B	M
1	12	165
2	2	5
3	343	40
4	0	2

Cluster 1 largely corresponds to malignancy and cluster 3 to benign cells. But the clustering on PCA results provided a more accurate and precise clustering compared to hierarchical.

Sensitivity

Q17. Which of your analysis procedures resulted in a clustering model with the best specificity? How about sensitivity?

Hierarchical clustering: Positive => cancer M Negative => non-cancerous B

True Positive 165 False Positive 14 True Negative 343 False Negative 47

Sensitivity: $TP/(TP+FN) = 165/212 = 0.778$ Specificity: $TN/(TN+FP) = 343/390 = 0.879$

PC Clustering: True Positive 188 False Positive 28 True Negative 329 False Negative 24

Sensitivity: $TP/(TP+FN) = 188/212 = 0.887$ Specificity: $TN/(TN+FP) = 329/353 = 0.932$

The clustering on PCA results had better specificity and sensitivity.


Prediction

We can use our PCA results (`wisc.pr`) to make predictions on new unseen data.

```
#url <- "new_samples.csv"
url <- "https://tinyurl.com/new-samples-CSV"
new <- read.csv(url)
npc <- predict(wisc.pr, newdata=new)
```



```
plot(wisc.pr$x[,1:2], col=grps)
points(npc[,1], npc[,2], col="blue", pch=16, cex=3)
text(npc[,1], npc[,2], c(1,2), col="white")
```



Q18. Which of these new patients should we prioritize for follow up based on your results?

Patient 2 as they fall within the malignant cluster.