

Class 18: Pertussis Mini Project

Sion Kang (PID: A17052234)

2025-03-06

Pertussis (a.k.a.) Whooping Cough is a deadly lung infection caused by the bacteria B. Pertussis.

The CDC tracks Pertussis cases around the US. At <https://tinyurl.com/pertussiscdc>

We can “scrape” this data using the R **datapasta** package.


```
head(cdc)
```

```
  year  cases
1 1922 107473
2 1923 164191
3 1924 165418
4 1925 152003
5 1926 202210
6 1927 181411
```

Q1, Q2.

```
library(ggplot2)

ggplot(cdc) +
  aes(year, cases) +
  geom_line() +
  geom_vline(xintercept = 1946, col="blue") +
  geom_vline(xintercept = 1996, col="red") +
  geom_vline(xintercept = 2020, col="gray") +
  geom_vline(xintercept = 2003, col="darkgreen")
```



Q3. Describe what happened after the introduction of the aP vaccine? Do you have a possible explanation for the observed trend?

There were high case numbers before the first wP (whole-cell) vaccine roll out in 1946 then a rapid decline in case numbers until 2004 when we have our first large-scale outbreak of pertussis again, not long after the switch to aP vaccine. This could be due to people being more skeptical of vaccination, but it is unclear what the difference is between people that have gotten the wP vs. aP vaccines. There is a notable COVID related dip and recent rapid rise.

Q. What is different about the immune response to infection if you had an older wP vaccine vs the newer aP vaccine?

Computational Models of Immunity - Pertussis Boost (CMI-PB)

The CMI-PB project aims to address this key question: what is different between aP and wP individuals?

We can get all the data from this ongoing project via JSON API calls For this we will use the **jsonlite** package. We can install with: `install.packages("jsonlite")`

```
library(jsonlite)
subject <- read_json("https://www.cmi-pb.org/api/v5_1/subject", simplifyVector = TRUE)
head(subject)
```

	subject_id	infancy_vac	biological_sex	ethnicity	race
1	1	wP	Female	Not Hispanic or Latino	White
2	2	wP	Female	Not Hispanic or Latino	White
3	3	wP	Female	Unknown	White
4	4	wP	Male	Not Hispanic or Latino	Asian
5	5	wP	Male	Not Hispanic or Latino	Asian
6	6	wP	Female	Not Hispanic or Latino	White

	year_of_birth	date_of_boost	dataset
1	1986-01-01	2016-09-12	2020_dataset
2	1968-01-01	2019-01-28	2020_dataset
3	1983-01-01	2016-10-10	2020_dataset
4	1988-01-01	2016-08-29	2020_dataset
5	1991-01-01	2016-08-29	2020_dataset
6	1988-01-01	2016-10-10	2020_dataset

Q. How many individuals “subjects” are in this data set?

```
nrow(subject)
```

```
[1] 172
```

Q4. How many wP and aP primed individuals are in this dataset?

```
table( subject$infancy_vac )
```

```
aP wP
87 85
```

Q5. How many male/female are there?

```
table ( subject$biological_sex )
```

```
Female    Male
  112      60
```

Q6. What is the breakdown of race and biological sex?

```
table(subject$race, subject$biological_sex)
```

	Female	Male
American Indian/Alaska Native	0	1
Asian	32	12
Black or African American	2	3
More Than One Race	15	4
Native Hawaiian or Other Pacific Islander	1	1
Unknown or Not Reported	14	7
White	48	32

This is not representative of the US population but it is the biggest dataset of its type so let's see what we can learn.

Obtain more data from CMI-PB:

```
specimen <- read_json("http://cmi-pb.org/api/v5_1/specimen", simplifyVector = TRUE)
ab_data <- read_json("http://cmi-pb.org/api/v5_1/plasma_ab_titer", simplifyVector = TRUE)
```

```
head(specimen)
```

	specimen_id	subject_id	actual_day_relative_to_boost	
1	1	1	-3	
2	2	1	1	
3	3	1	3	
4	4	1	7	
5	5	1	11	
6	6	1	32	

	planned_day_relative_to_boost	specimen_type	visit
1	0	Blood	1
2	1	Blood	2
3	3	Blood	3
4	7	Blood	4
5	14	Blood	5
6	30	Blood	6

```
head(ab_data)
```

	specimen_id	isotype	is_antigen_specific	antigen	MFI	MFI_normalised
1	1	IgE	FALSE	Total	1110.21154	2.493425
2	1	IgE	FALSE	Total	2708.91616	2.493425
3	1	IgG	TRUE	PT	68.56614	3.736992
4	1	IgG	TRUE	PRN	332.12718	2.602350
5	1	IgG	TRUE	FHA	1887.12263	34.050956
6	1	IgE	TRUE	ACT	0.10000	1.000000

	unit	lower_limit_of_detection
1	UG/ML	2.096133
2	IU/ML	29.170000
3	IU/ML	0.530000
4	IU/ML	6.205949
5	IU/ML	4.679535
6	IU/ML	2.816431

I now have 3 tables of data from CMI-PB: `subject`, `specimen`, `ab_data`. I need to “join” these tables so I will have all the info I need to work with.

For this we will use the `inner_join()` function from **dplyr** package.

Q9. Complete the code to join specimen and subject tables to make a new merged data frame containing all specimen records along with their associated subject details:

```
library(dplyr)
```

Attaching package: 'dplyr'

The following objects are masked from 'package:stats':

`filter`, `lag`

The following objects are masked from 'package:base':

`intersect`, `setdiff`, `setequal`, `union`

```
meta <- inner_join(subject, specimen)
```

Joining with `by = join_by(subject_id)`

```
head(meta)
```

	subject_id	infancy_vac	biological_sex	ethnicity	race
1	1	wP	Female Not Hispanic or Latino	White	
2	1	wP	Female Not Hispanic or Latino	White	
3	1	wP	Female Not Hispanic or Latino	White	
4	1	wP	Female Not Hispanic or Latino	White	
5	1	wP	Female Not Hispanic or Latino	White	
6	1	wP	Female Not Hispanic or Latino	White	

	year_of_birth	date_of_boost	dataset	specimen_id
1	1986-01-01	2016-09-12	2020_dataset	1
2	1986-01-01	2016-09-12	2020_dataset	2
3	1986-01-01	2016-09-12	2020_dataset	3
4	1986-01-01	2016-09-12	2020_dataset	4
5	1986-01-01	2016-09-12	2020_dataset	5
6	1986-01-01	2016-09-12	2020_dataset	6

	actual_day_relative_to_boost	planned_day_relative_to_boost	specimen_type
1	-3	0	Blood
2	1	1	Blood
3	3	3	Blood
4	7	7	Blood
5	11	14	Blood
6	32	30	Blood

	visit
1	1
2	2
3	3
4	4
5	5
6	6

```
dim(subject)
```

```
[1] 172  8
```

```
dim(specimen)
```

```
[1] 1503  6
```

```
dim(meta)
```

```
[1] 1503  13
```

Q10. Now using the same procedure join meta with titer data so we can further analyze this data in terms of time of visit aP/wP, male/female etc.

Now we can join our `ab_data` table to `meta` so we have all the info we need about antibody levels.

```
abdata <- inner_join(meta, ab_data)
```

Joining with ``by = join_by(specimen_id)``

```
head(abdata)
```

	subject_id	infancy_vac	biological_sex	ethnicity	race
1	1	wP	Female Not Hispanic or Latino	White	
2	1	wP	Female Not Hispanic or Latino	White	
3	1	wP	Female Not Hispanic or Latino	White	
4	1	wP	Female Not Hispanic or Latino	White	
5	1	wP	Female Not Hispanic or Latino	White	
6	1	wP	Female Not Hispanic or Latino	White	

	year_of_birth	date_of_boost	dataset	specimen_id
1	1986-01-01	2016-09-12	2020_dataset	1
2	1986-01-01	2016-09-12	2020_dataset	1
3	1986-01-01	2016-09-12	2020_dataset	1
4	1986-01-01	2016-09-12	2020_dataset	1
5	1986-01-01	2016-09-12	2020_dataset	1
6	1986-01-01	2016-09-12	2020_dataset	1

	actual_day_relative_to_boost	planned_day_relative_to_boost	specimen_type
1	-3	0	Blood
2	-3	0	Blood
3	-3	0	Blood
4	-3	0	Blood
5	-3	0	Blood
6	-3	0	Blood

	visit	isotype	is_antigen_specific	antigen	MFI	MFI_normalised	unit
1	1	IgE	FALSE	Total	1110.21154	2.493425	UG/ML
2	1	IgE	FALSE	Total	2708.91616	2.493425	IU/ML
3	1	IgG	TRUE	PT	68.56614	3.736992	IU/ML

4	1	IgG	TRUE	PRN	332.12718	2.602350	IU/ML
5	1	IgG	TRUE	FHA	1887.12263	34.050956	IU/ML
6	1	IgE	TRUE	ACT	0.10000	1.000000	IU/ML

	lower_limit_of_detection
1	2.096133
2	29.170000
3	0.530000
4	6.205949
5	4.679535
6	2.816431

Q11. How many different antibody isotypes are there in this dataset? How many specimens (i.e. entries in abdata) do we have for each isotype?

```
table(abdata$isotype)
```

IgE	IgG	IgG1	IgG2	IgG3	IgG4
6698	7265	11993	12000	12000	12000

Q12. What are the different \$dataset values in abdata and what do you notice about the number of rows for the most “recent” dataset?

```
table(abdata$dataset)
```

2020_dataset	2021_dataset	2022_dataset	2023_dataset
31520	8085	7301	15050

The most “recent” dataset (2023) has double the number of rows of the previous year.

```
table(abdata$antigen)
```


ACT	BETV1	DT	FELD1	FHA	FIM2/3	LOLP1	LOS	Measles	OVA
1970	1970	6318	1970	6712	6318	1970	1970	1970	6318
PD1	PRN	PT	PTM	Total	TT				
1970	6712	6712	1970	788	6318				

Q13. Complete the following code to make a summary boxplot of Ab titer levels (MFI) for all antigens:


I want a plot of antigen levels across the whole dataset.

```
ggplot(abdata) +  
  aes(MFI, antigen) +  
  geom_boxplot()
```

Warning: Removed 1 row containing non-finite outside the scale range
(`stat_boxplot()`).




```
ggplot(abdata) +  
  aes(MFI_normalised, antigen) +  
  geom_boxplot()
```



Antigens like FIM2/3, PT, FELD1 have quite a large range of values. Others like Measles don't show much activity.

```
ggplot(abdata) +
  aes(MFI_normalised, antigen) +
  geom_boxplot() +
  xlim(0,75) +
  facet_wrap(vars(visit), nrow=2)
```

Warning: Removed 983 rows containing non-finite outside the scale range (`stat_boxplot()`).




Q14. What antigens show differences in the level of IgG antibody titers recognizing them over time? Why these and not others?


IgG antibodies against antigens like PT, PRN, FIM2/23, and FHA seem to decrease over time. This is because these are the antigens that are introduced in the aP and wP vaccines and the presence of IgG that detect them decrease over time.

Q. Are there differences at the whole-dataset level between aP and wP?

```
ggplot(abdata) +
  aes(MFI_normalised, antigen, col=infancy_vac) +
  geom_boxplot()
```



```
ggplot(abdata) +
  aes(MFI_normalised, antigen, col=infancy_vac) +
  geom_boxplot() +
  facet_wrap(~infancy_vac)
```



Examine IgG Ab titer levels

For this I need to select out just isotype IgG.

```
igg <- abdata %>%
  filter(isotype == "IgG")
head(igg)
```

	subject_id	infancy_vac	biological_sex	ethnicity	race
1	1	wP	Female	Not Hispanic or Latino	White
2	1	wP	Female	Not Hispanic or Latino	White
3	1	wP	Female	Not Hispanic or Latino	White
4	1	wP	Female	Not Hispanic or Latino	White
5	1	wP	Female	Not Hispanic or Latino	White
6	1	wP	Female	Not Hispanic or Latino	White

	year_of_birth	date_of_boost	dataset	specimen_id
1	1986-01-01	2016-09-12	2020_dataset	1
2	1986-01-01	2016-09-12	2020_dataset	1
3	1986-01-01	2016-09-12	2020_dataset	1
4	1986-01-01	2016-09-12	2020_dataset	2
5	1986-01-01	2016-09-12	2020_dataset	2
6	1986-01-01	2016-09-12	2020_dataset	2


	actual_day_relative_to_boost	planned_day_relative_to_boost	specimen_type
1	-3	0	Blood
2	-3	0	Blood
3	-3	0	Blood
4	1	1	Blood
5	1	1	Blood
6	1	1	Blood

	visit	isotype	is_antigen_specific	antigen	MFI	MFI_normalised	unit
1	1	IgG	TRUE	PT	68.56614	3.736992	IU/ML
2	1	IgG	TRUE	PRN	332.12718	2.602350	IU/ML
3	1	IgG	TRUE	FHA	1887.12263	34.050956	IU/ML
4	2	IgG	TRUE	PT	41.38442	2.255534	IU/ML
5	2	IgG	TRUE	PRN	174.89761	1.370393	IU/ML
6	2	IgG	TRUE	FHA	246.00957	4.438960	IU/ML

	lower_limit_of_detection
1	0.530000
2	6.205949
3	4.679535
4	0.530000
5	6.205949
6	4.679535

An overview boxplot:

```
ggplot(igg) +
  aes(MFI_normalised, antigen, col=infancy_vac) +
  geom_boxplot()
```



Digging in further to look at the time course of IgG isotype PT antigen levels across aP and wP individuals:

```
abdata.21 <- abdata %>% filter(dataset == "2021_dataset") ## Filter to include 2021 data only

abdata.21 %>% ## Filter to look at IgG PT data only
  filter(isotype == "IgG", antigen == "PT") %>%
  ggplot() + ## Plot
    aes(x=planned_day_relative_to_boost,
         y=MFI_normalised,
         col=infancy_vac,
         group=subject_id) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept=0, linetype="dashed") +
  geom_vline(xintercept=14, linetype="dashed") +
  labs(title="2021 dataset IgG PT",
        subtitle = "Dashed lines indicate day 0 (pre-boost) and 14 (apparent peak levels)")
```

2021 dataset IgG PT

Dashed lines indicate day 0 (pre-boost) and 14 (apparent peak levels)

