

Universität Bonn

Physikalisches Institut

Measurement of a single top quark production in association with a Z boson in ATLAS

Kanhaiya Gupta

The total cross-section of a single top quark production in association with a Z boson is measured in proton-proton collision data at a center-of-mass energy of 13 TeV recorded by the ATLAS experiment at LHC. The full Run II dataset (2015-2018) which corresponds to an integrated luminosity of 139 fb^{-1} is analysed. In this production mechanism, the top quark is produced via the t-channel and the Z boson is either radiated off from one of the participating quarks or produced via W boson fusion, leading to a signature of a single top quark, a Z boson and an additional quark. Events are selected based on the presence of three leptons (electrons or muons), two or three jets (one identified as coming from a b-quark), and substantial missing transverse energy. The three leptons come from leptonic Z boson decay and semi-leptonic top quark decay. Major backgrounds come from diboson, $t\bar{t}V$, Z+jets and $t\bar{t}$. The non-prompt leptons background processes which are difficult to simulate are studied by implementing different isolation criteria. A python based (tensorflow) neural network algorithm is trained in order to separate the signal and background distributions. A simultaneous binned maximum-likelihood fit of the signal regions and control regions is performed and from the fit cross-section is extracted. The extracted cross-section of tllq is $\sigma(pp \rightarrow tZq \rightarrow Wbl^+l^-q) = 132 \pm 12 \text{ (stat)} \pm 17 \text{ (syst)} \text{ fb}$ with the SM prediction of 102 fb under the assumption of a top-quark mass of $m_t = 172.5 \text{ GeV}$.

Physikalisches Institut der
Universität Bonn
Nussallee 12
D-53115 Bonn

BONN-IR-2022-01
January 2022

Contents

1	Introduction	1
2	Theoretical concepts	3
2.1	The standard model of particle physics	3
2.2	Physics at hadron colliders	3
2.3	Top quark physics	3
3	The LHC and the ATLAS experiment	5
3.1	Large hadron collider	5
3.2	A toroidal LHC apparatus (ATLAS)	5
3.2.1	Inner detector	5
3.2.2	Calorimetry	5
3.2.3	Muon system	5
3.2.4	Trigger system	5
3.3	Physics objects reconstruction in ATLAS	5
3.3.1	Electrons	5
3.3.2	Muons	5
3.3.3	Jets	5
3.3.4	Missing transverse energy	5
4	Data and Monte Carlo simulated events	7
4.1	Data sample	7
4.2	Monte Carlo simulation	7
4.2.1	Signal sample	7
4.2.2	Background sample	7
4.2.3	Re-weighting of Monte Carlo simulated events.	7
5	Events selection	9
5.1	Full event reconstruction	9
5.2	Signal region	10
5.2.1	Lepton isolation working point	11
5.2.2	Optimization of cuts	13
5.2.3	Signal regions (SRs) plots and yields	15
5.3	Control regions (CRs)	17
5.3.1	Diboson CRs plots and yields	18
5.3.2	$t\bar{t}Z$ CRs plots and yield	19

5.3.3	$t\bar{t}$ CRs plots and yields	21
6	Analysis	23
6.1	Cross-section measurement analysis strategy	23
6.2	Signal and background separation	23
6.3	Multivariate analysis technique: Artificial neural network (NN)	24
6.3.1	Input variables	25
6.3.2	Data and MC comparison	25
6.3.3	NN training in the SRs and $t\bar{t}Z$ CRs	29
6.4	Signal extraction - Profile likelihood fit	33
6.4.1	Fitted regions	33
6.4.2	Binning optimization	33
6.5	Systematics uncertainties	33
6.5.1	Sources of systematics uncertainties	34
6.5.2	Symmetrizing, smoothing and pruning	34
7	Results	35
7.1	Expected fit results	35
7.2	Observed fit results	35
7.3	Discussion of the results	35
8	Conclusion	37
	Bibliography	39