



CSCI 2824: Discrete Structures

Lecture 28: Count-ing. Permutations. Combinations



Reminders

Submissions:

- Homework 9: **Fri 11/01 at noon** - Moodle

Readings:

- Ch. 6 – Counting

Midterm II

- Tuesday November 5th



$$E=mc^2$$




Last time

Recap:

- **Generalized Pigeonhole Principle (PHP)** -- If you want to sort n objects into k bins, at least one of the bins will contain $[n/k]$ objects.

Today:

- Permutations and Combinations.



$$E=mc^2$$




Permutations and combinations

Example: How many 4-digit PIN combinations are there



$$(10 \text{ choices for first digit}) \times (10 \text{ choices for second digit}) \times (10 \dots) \times (10 \dots) = 10,000$$



$$E=mc^2$$



Permutations and combinations

Example: How many 4-digit PIN combinations are there?



$$(10 \text{ choices for first digit}) \times (10 \text{ choices for second digit}) \times (10 \dots) \times (10 \dots) = 10,000$$

Digits 0-9 like 10 distinct items in a bag

- Pick first PIN digit: pull a number out of the bag
- Then put digit back into bag. Do this 4 times (digits)



$$E=mc^2$$



Permutations and combinations

Example: How many 4-digit PIN combinations are there?



$$(10 \text{ choices for first digit}) \times (10 \text{ choices for second digit}) \times (10 \dots) \times (10 \dots) = 10,000$$



Digits 0-9 like 10 distinct items in a bag

- Pick first PIN digit: pull a number out of the bag
- Then put digit back into bag. Do this 4 times (digits)



What if we picked from the bag without replacement and (still) care about their order?


What if we picked from the bag without replacement and don't care about order?

Permutations and combinations

Example: How many 4-digit PIN combinations are there?



$$(10 \text{ choices for first digit}) \times (10 \text{ choices for second digit}) \times (10 \dots) \times (10 \dots) = 10,000$$



Digits 0-9 like 10 distinct items in a bag

- Pick first PIN digit: pull a number out of the bag
- Then put digit back into bag. Do this 4 times (digits)



What if we picked from the bag without replacement and (still) care about their order?

⇒ *permutations*



What if we picked from the bag without replacement and don't care about order?


⇒ *combinations*



Permutations

When selecting r distinct items without replacing them, we call the possible selection an r -permutation

Example: Find all 2-permutations of the set $S = \{a, b, c\}$



$$E = mc^2$$



Permutations



BOE

$E=mc^2$



When selecting r distinct items without replacing them, we call the possible selection an **r -permutation**

Example: Find all 2-permutations of the set $S = \{a, b, c\}$

ab, ac, ba, bc, ca, cb

Note that there are $3 \times 2 = 6$ such 2-permutations, and the list contains both ab and ba (for example) because **order matters**

Permutations


Example: How many three-character strings can we make if each character is a distinct lowercase letter?



$$E = mc^2$$



Permutations



BOH

$E = mc^2$



11

Example: How many three-character strings can we make if each character is a distinct lowercase letter?

Solution: 26 choices for the first letter, 25 choices for second, 24 choices for third
 $\Rightarrow 26 \times 25 \times 24 = 15,600$ strings

Theorem: If n is a positive integer and r is an integer such that $0 \leq r \leq n$, then the number of r -permutations from a set of size n is

$$P(n, r) = n(n - 1)(n - 2) \dots (n - r + 1)$$

Special cases: $P(n, 0) = 1$ and $P(n, n) = n!$


$$\frac{n!}{(n-r)!}$$

Permutations

Example: How many different 1st-2nd-3rd place permutations can occur in a race with 5 runners?



Example: How many permutations of the letters *ABCDEFGHI* contain the string *ABC*?



$$BOE$$

$$E=mc^2$$



Permutations

Example: How many different 1st-2nd-3rd place permutations can occur in a race with 5 runners?

Solution: Let's use the factorial formula for $P(5,3)...$


$$P(5, 3) = 5! / (5-3)! = 5! / 2! = 120 / 2 = 60$$




Example: How many permutations of the letters ABCDEFGH contain the string ABC?

Think of ABC as a block, a single character. In addition to ABC, there are 5 other characters, for a total of 6 discrete items.

⇒ Total number of permutations is $P(6, 6) = 6! = 720$



Combinations



When we make an *unordered* selection of distinct items, it's called a **combination**

Example: Find all 2-letter combinations of the set $S = \{a, b, c\}$

The 2-permutations were: ab, ac, ba, bc, ca, cb

But now we throw out all the repeats, now that we don't care about order;

The 2-combinations are: ab, ac, bc

Repeats: ba, ca, cb

⇒ there are **3** 2-combos of the letters in S

Combinations

We can come up with a formula for the number of r -combinations ($C(n,r)$) by starting with the formula for the number of r -permutations ($P(n,r)$), and the **product rule**:

In English:

[# r -permutations of n elements] = [# pick r elements from n] \times [# ways to arrange them]

Combinations



$$E=mc^2$$



We can come up with a formula for the number of r -combinations ($C(n,r)$) by starting with the formula for the number of r -permutations ($P(n,r)$), and the **product rule**:

In English:

[# r -permutations of n elements] = [# pick r elements from n] \times [# ways to arrange them]

$$P(n, r) = C(n, r) \times P(r, r)$$

Which means:

$$C(n, r) = \frac{P(n, r)}{P(r, r)} = \frac{n!}{(n - r)! r!} = \binom{n}{r}$$

Alternative notation: Since we are *choosing* r elements from n elements, sometimes we say “ n choose r ” and denote it using the **binomial coefficient** (on ¹⁹the right)

Combinations

Example: How many 5-card poker hands can be drawn from a 52-card deck?

Example: There are about 650 computer science majors at CU. How many ways are there to choose 6 of them to serve as on an undergraduate CS committee?



$$E=mc^2$$



Combinations

Example: How many 5-card poker hands can be drawn from a 52-card deck?

We don't care about the order in which the cards are received so we want **5-combinations**

$$C(52, 5) = \frac{52!}{(52 - 5)! 5!} = \frac{52!}{47! 5!} = \frac{52 \times 51 \times 50 \times 49 \times 48}{5 \times 4 \times 3 \times 2 \times 1}$$

= ... = **2,598,960 distinct 5-card poker hands**

Example: There are about 650 computer science majors at CU. How many ways are there to choose 6 of them to serve as on an undergraduate CS committee?


Solution: We want $C(650, 6) = \dots$ = (a very large number)

Combinations

Example: How many bit-strings of length 10:

- a) Contain exactly four 1's?
- b) Contain at most four 1's?

Combinations



BOE

E = mc²



Example: How many bit-strings of length 10:

- a) Contain exactly four 1's?
- b) Contain at most four 1's?

Solution:

- a) Choose where the four 1s go:

$$C(10, 4) = \frac{10!}{6! 4!} = \frac{10 \cdot 9 \cdot 8 \cdot 7}{4 \cdot 3 \cdot 2 \cdot 1} = 210$$

- b) Count up the # ways to have zero 1s, one 1, two 1s, three 1s, or four 1s:

$$C(10, 0) + C(10, 1) + C(10, 2) + C(10, 3) + C(10, 4) = \dots = 386$$

Permutations and combinations

It can be tricky to know when to use permutations, or combinations, or both.

Example: The English alphabet contains 21 consonants and 5 vowels. How many strings of six letters contain exactly 2 vowels?

Permutations and combinations



$$E=mc^2$$



It can be tricky to know when to use permutations, or combinations, or both.

Example: The English alphabet contains 21 consonants and 5 vowels. How many strings of six letters contain exactly 2 vowels?

Solution: by product rule, we have 3 tasks:

1. Pick where the vowels go $\Rightarrow C(6, 2)$


2. Pick what the vowels are $\Rightarrow 5 \times 5$

3. Pick what the consonants are $\Rightarrow 21 \times 21 \times 21 \times 21$

\Rightarrow There are $C(6, 2) \times 5^2 \times 21^4 = 15 \times 25 \times 21^4 = \dots = 72,930,375$ such strings

Permutations and combinations

Example: How many ways are there for eight regular people and the five dysfunctional members of the Scooby Doo gang to stand in a line so that no two members of the Scooby Doo gang stand next to each other?



Permutations and combinations




$$E=mc^2$$



Example: How many ways are there for eight regular people and the five dysfunctional members of the Scooby Doo gang to stand in a line so that no two members of the Scooby Doo gang stand next to each other?

Solution:



Product rule, 2 tasks:

- 9 places SDG can stand, need to pick 5, and order matters $\Rightarrow P(9, 5)$
 - Arrange the 8 regular people $\Rightarrow P(8, 8)$
- \Rightarrow Have $P(8, 8) \times P(9, 5) = (8!) \times (9! / 4!) = 609,638,400$ possible arrangements

Permutations and combinations

FYOG: A club has 25 members. How many ways are there to choose a president, a vice president, a secretary and a treasurer? (Assume each member can hold at most one position)

FYOG: How many strings of six letters contain *at least* one vowel?

FYOG: How many strings of six letters contain *at least* two vowels?

FYOG: S'pose that a department contains 10 men and 15 women. How many ways are there to form a committee with six members if it must have the same number of men as women?

Permutations and combinations

Previously: calculate # of permutations and combinations of r objects from a set of n total objects **without repetition**

→ Care about order? **permutations:**

$$P(n, r) = n(n - 1)(n - 2) \cdots (n - r + 1) = \frac{n!}{(n - r)!}$$

→ Order doesn't matter? **combinations:**

$$C(n, r) = \frac{n!}{(n - r)! r!}$$

Now: we want to count permutations and/or combinations **with repetition**

Permutations with repetition

... return of the product rule!

Example: How many strings of length 4 can be formed using the uppercase letters of the English alphabet?

Permutations with repetition

... return of the product rule!

Example: How many strings of length 4 can be formed using the uppercase letters of the English alphabet?

Solution: Since order matters (AMCE is different from MACE, e.g.), we want permutations

⇒ We can bring back the analogy of filling slots:

[26 choices for 1st char.] x [26 choices for 2nd char.] x [26 choices for 3rd char.] x [26 choices for 4th char.]

$$= 26^4$$

Theorem: The number of r -permutations of n objects with repetition is n^r



$$E = mc^2$$



Combinations with repetition

Example: S'pose that a bakery in the Engineering Center Lobby has 4 different kinds of donuts. How many different ways can I select 6 delicious, *delicious* donuts?




$$E=mc^2$$



Combinations with repetition

Example: S'pose that a bakery in the Engineering Center Lobby has 4 different kinds of donuts. How many different ways can I select 6 delicious, ***delicious*** donuts?



Solution: Since order does *not* matter (I am for sure going to eat them all anyway), we want **combinations**

Theorem: There are $C(n+r-1, r) = C(n+r-1, n-1)$ r -combinations from a set with n elements where repetition of elements is allowed.

$$\text{So we have: } C(n + r - 1, r) = C(9, 6) = \frac{9!}{(9 - 6)! \cdot 6!} = \frac{9 \cdot 8 \cdot 7}{3 \cdot 2 \cdot 1} = 84$$

Okay... but why does this work?

Combinations with repetition

Theorem: There are $C(n+r-1, r) = C(n+r-1, n-1)$ r -combinations from a set with n elements where repetition of elements is allowed.

This problem is asking us to distribute our 6 donuts choices among the 4 donut types. We could represent each of our donut choices with a ★, like so:



$$E=mc^2$$



Combinations with repetition

Theorem: There are $C(n+r-1, r) = C(n+r-1, n-1)$ r -combinations from a set with n elements where repetition of elements is allowed.

This problem is asking us to distribute our 6 donuts choices among the 4 donut types. We could represent each of our donut choices with a ★, like so:

★★



★



★★★



|



$$E=mc^2$$



Combinations with repetition

Theorem: There are $C(n+r-1, r) = C(n+r-1, n-1)$ r -combinations from a set with n elements where repetition of elements is allowed.

This problem is asking us to distribute our 6 donuts choices among the 4 donut types. We could represent each of our donut choices with a ★, like so:

And we only really need to represent where the *divisions* between donut categories are. Let's use a bar (|) for this:

★★



|

★



|

★★★



|



This gives 2 glazed, 1 chocolate-frosted, 3 pink sprinkles, and 0 jelly-filled



DOE

$$E=mc^2$$



Combinations with repetition

Theorem: There are $C(n+r-1, r) = C(n+r-1, n-1)$ r -combinations from a set with n elements where repetition of elements is allowed.



DOE

$$E=mc^2$$



This problem is asking us to distribute our 6 donut choices among the 4 donut types. We could represent each of our donut choices with a \star , like so:

And we only really need to represent where the *divisions* between donut categories are. Let's use a bar (|) for this:



Upshot: the problem is reduced to asking how many different ways can we arrange r stars and $n-1$ bars.

Combinations with repetition

Upshot: the problem is reduced to asking how many different ways can we arrange r stars and $n-1$ bars.


Each unique arrangement of r stars (the things we're selecting) and $n-1$ bars (the divisions between the bins) gives a different distribution of our selections.

We have $n+r-1$ total objects (stars and bars) and need to pick r of them to be stars:

$$C(n+r-1, r)$$

Similarly, we could also pick $n-1$ of our objects to be bars:

$$C(n+r-1, n-1)$$



$$E=mc^2$$



Combinations with repetition

And these two quantities are equal:

$$C(n + r - 1, r) = \frac{(n + r - 1)!}{(n + r - 1 - (r))! \ r!} = \frac{(n + r - 1)!}{(n - 1)! \ r!}$$

... and...

$$C(n + r - 1, n - 1) = \frac{(n + r - 1)!}{(n + r - 1 - (n - 1))! \ (n - 1)!} = \frac{(n + r - 1)!}{r! \ (n - 1)!}$$



BOE

E = mc²



Combinations with repetition

Example: How many solutions does the equation

$$x_1 + x_2 + x_3 = 11$$

have, where x_1, x_2 and x_3 are all nonnegative integers?

Combinations with repetition



$$E=mc^2$$



Example: How many solutions does the equation

$$x_1 + x_2 + x_3 = 11$$

have, where x_1, x_2 and x_3 are all nonnegative integers?

Solution: Think of each of x_1, x_2 and x_3 as the bins into which we must sort 11 items. Then this is stars ‘n’ bars with 11 stars and 2 bars. For example, one possibility is

★★★ | ★★ | ★★★★★★

which represents $x_1 = 3, x_2 = 2, x_3 = 6$.

So we have $C(n + r - 1, r) = C(13, 11) = \frac{13!}{(13 - 11)! 11!} = \frac{13 \cdot 12}{2 \cdot 1} = 78$

Combinations with repetition

Example: How many solutions does the equation

$$x_1 + x_2 + x_3 = 11$$

have, if x_1, x_2 and x_3 are all nonnegative integers that satisfy $x_1 \geq 1, x_2 \geq 2, x_3 \geq 3$?

Combinations with repetition



$$E = mc^2$$



Example: How many solutions does the equation

$$x_1 + x_2 + x_3 = 11$$

have, if x_1, x_2 and x_3 are all nonnegative integers that satisfy $x_1 \geq 1, x_2 \geq 2, x_3 \geq 3$?

Solution: This is the same as the previous problem, except we first need to allocate 1 star to the left-most bin, 2 stars to the center bin, and 3 stars to the right-most bin.

★ | ★★ | ★★★


So we're left with 5 stars to distribute among the 3 bins (separated by 2 bars) as we wish.

So we have

43

$$C(n + r - 1, r) = C(7, 5) = \frac{7!}{(7 - 5)! 5!} = \frac{7 \cdot 6}{2 \cdot 1} = 21$$

Permutations/combinations with repetition



$$E=mc^2$$



Useful Recap:
(Rosen, p. 427)

TABLE 1 Combinations and Permutations With and Without Repetition.


Type	Repetition Allowed?	Formula
r -permutations	No	$\frac{n!}{(n - r)!}$
r -combinations	No	$\frac{n!}{r! (n - r)!}$
r -permutations	Yes	n^r
r -combinations	Yes	$\frac{(n + r - 1)!}{r! (n - 1)!}$

Next time:

- More permutations and combinations, and the Binomial Theorem
- More **counting**, and moving toward **probability!**

Pigeonhole principle

Example: Show that for any number n there is a multiple of n that has only 0s and 1s in its decimal representation.



$$E = mc^2$$



Pigeonhole principle



$$E=mc^2$$



Example: Show that for any number n there is a multiple of n that has only 0s and 1s in its decimal representation.

Solution: Consider the $n+1$ numbers $1, 11, 111, \dots, 111\cdots11$, where the last number has $n+1$ ones in it.

Note that if we divide a number by n , there are only n possible remainders:

$$0, 1, 2, \dots, n-1$$

Since there are $n+1$ numbers above but only n possible remainders, by PHP at least 2 of them must have the same remainder when divided by n .

⇒ Let those numbers be a and b , where $b > a$

⇒ $n \mid (b-a)$, which means that $b-a$ is a multiple of n

⇒ $b-a$ is a number made up of only 0s and 1s (e.g., $b=111$ and $a=11$ give $b-a = 100$)

■

Pigeonhole principle



III

Example: Show that for any number n there is a multiple of n that has only 0s and 1s in its decimal representation.



An example of the example: Suppose we let $n = 7$



We check: 1 11 111 1,111 11,111 111,111 1,111,111 11,111,111


Then (it turns out) $1,111,111 \equiv 1 \pmod{7}$ so 1 and 1,111,111 have the same remainder when divided by 7



$\Rightarrow 1,111,111 - 1 = 1,111,110$ is divisible by 7

$(1,111,110 / 7 = 158,730)$

```
In [5]: n=7  
numbers = [1]  
for i in range(1,n):  
    numbers.append(10**i + numbers[i-1])
```



```
remainders = []  
for i in range(0,n):  
    remainders.append(numbers[i] % n)
```



```
In [6]: print(remainders)  
[1, 4, 6, 5, 2, 0, 1]
```



47