Requirements for the drug dispensing tool

i.Patients are identified by SSN, and their names, addresses, and also ages.

ii. Doctors are identified by an SSN, for each doctor, the name, specialty and years of experience must be recorded.

iii. Each pharmaceutical company is identified by name and has a phone number.

iv. For each drug, the trade name and formula must be reordered. Each drug is sold by a given pharmaceutical company, and the trade name identifies a drug uniquely from among the products of that company. If a pharmaceutical company is deleted, you need not keep track of its products any longer.

v. Each pharmacy has a name, address, and phone number.

vi. Every patient has a primary physician. Every doctor has at least one patient.

vii. Each pharmacy sells several drugs and has a price for each. A drug could be sold at several pharmacies, and the price could vary from one pharmacy to another.

viii. Doctors prescribe drugs for patients. A doctor could prescribe one or more drugs for several patients, and a patient could obtain prescriptions from several doctors. Each prescription has a date and a quantity associated with it. You can assume that if a doctor prescribes the same drug for the same patient more than once, only

the last such prescription needs to be stored.

ix. Pharmaceutical companies have long-term contracts with pharmacies. A
pharmaceutical company can contract with several pharmaceutical companies.
For each contract, you have to store a start date, and end date, and the text of the
contract.

x. Pharmacies appoint a supervisor for each contract. There must always a supervisor
for each contract

Added requirements

- The patient should be able to make multiple orders.
- The user should be required to log in with a unique username and password.
- The patient should be invoiced after making an order and the invoice details recorded.
- The payment details of the order made should be recorded.
- The interface should be user-friendly.
- The program should be able to rectify errors for example, where the wrong drug has been prescribed, it should be rectified before the patient uses the drug.