

# **A Lecture on Topological Defect**

Kantaro Ohmori

2023-09-26

# Table of contents

<b>1</b>	<b>What is this</b>	<b>3</b>
1.1	Prerequisite . . . . .	3
1.2	What is contained and what is not . . . . .	3
1.3	Other Lectures/Reviews . . . . .	3
<b>2</b>	<b>Introduction</b>	<b>4</b>
2.1	Symmetry . . . . .	4
2.2	Locality . . . . .	4
<b>3</b>	<b>Summary</b>	<b>5</b>
	<b>References</b>	<b>6</b>

# 1 What is this

This is a lecture note prepared for two sets of “intensive lectures”:<sup>1</sup>

- at Tohoku University, Oct. 11-13, 2023, and
- at Yukawa Insititute for Theoretical Physics, Kyoto University, Nov. 29-1, 2023.

In this lecture I will try to explain the constructions of topological defects corresponding to generalized symmetries. Due to lack of time and (more significantly) my understanding, the lecture will focus on bosonic systems, and the generalization to fermionic systems is left for the readers/audiences.

## 1.1 Prerequisite

- Basic knowledge about scalar field theory and (abelian) gauge theory in path-integral formalism, and
- Knowledge about renormalization group (RG) flows to understand motivations.

## 1.2 What is contained and what is not

## 1.3 Other Lectures/Reviews

Recently there has been a surge of lecture notes/ review articles on generalized symmetries. The ones I have noticed are [1–6]. Because this lecture will focus on the fundamental aspects of the topic and will not connect very well with the existent literature (so sorry about that), readers/audiences are strongly encouraged to refer to at least one of them, or something similar.

Also, about conventional symmetries and their anomalies, there are nice old lectures. The one I would particularly recommend is [7].

---

<sup>1</sup>In Japan, an “intensive lecture” is a format of a lecture course where a lecturer (usually from another university) gives lectures in consecutive days filling 7-9 slots in usually 3 days.

## 2 Introduction

### 2.1 Symmetry

**Symmetry** plays a fundamental role in theoretical physics. In this lecture we consider them in *quantum field theories* (QFTs). The fundamental fact about symmetry in QFTs is that it is preserved along the renormalization group flow. More precisely, when an ultraviolet (UV) theory  $\mathcal{T}_{\text{UV}}$  flows into an infrared theory  $\mathcal{T}_{\text{IR}}$ , there is a canonical homomorphism  $f_{\text{RG}}$  from the UV symmetry group  $G_{\text{UV}}$  to the IR symmetry group  $G_{\text{IR}}$ :<sup>1</sup>

$$f_{\text{RG}} : G_{\text{UV}} \rightarrow G_{\text{IR}}. \quad (2.1)$$

Given this relation, there are two ways of applying symmetry in QFT:

- UV to IR: Given a microscopic model (e.g. a model of elementary particles or electrons in a matter), constrain/guess what happens in the macroscopic scale.
- IR to UV: Given some macroscopic phenomena, constrain/guess what could be the microscopic origin of it (e.g. guessing QCD lagrangian from hadron spectrum).

#### Warning

In this lecture, whenever something is called “symmetry”, it means a *global* symmetry. Here *global* means that the symmetry operation acts on the entire space.

### 2.2 Locality

The second important keyword in this lecture is **locality**.

---

<sup>1</sup>If the RG flow is from a finite energy to a lower nonzero energy, and if one retains all the (even very massive) degrees of freedom in the description of  $\mathcal{T}_{\text{IR}}$ ,  $f_{\text{RG}}$  is an isomorphism. However, typically one integrates out massive dofs in the description of  $\mathcal{T}_{\text{RG}}$ , in which case some symmetry can decouple and thus  $f_{\text{RG}}$  can be non-surjective (if  $G_{\text{IR}}$  is taken to be the *faithful* symmetry). Also, if one also drops some higher-order interaction terms, or runs the flow to the zero energy, there can be an *emergent* symmetry, in which case  $f_{\text{RG}}$  can be non-injective.

## 3 Summary

In summary, this book has no content whatsoever.

# References

- [1] J. McGreevy, “Generalized Symmetries in Condensed Matter”, [10.1146/annurev-conmatphys-040721-021029](#) (2022), [arXiv:2204.03045 \[cond-mat.str-el\]](#).
- [2] S. Schafer-Nameki, “ICTP Lectures on (Non-)Invertible Generalized Symmetries”, (2023), [arXiv:2305.18296 \[hep-th\]](#).
- [3] P. R. S. Gomes, “An introduction to higher-form symmetries”, [SciPost Phys. Lect. Notes](#) **74**, 1 (2023), [arXiv:2303.01817 \[hep-th\]](#).
- [4] L. Bhardwaj, L. E. Bottini, L. Fraser-Taliente, L. Gladden, D. S. W. Gould, A. Platschorre, and H. Tillim, “Lectures on Generalized Symmetries”, (2023), [arXiv:2307.07547 \[hep-th\]](#).
- [5] R. Luo, Q.-R. Wang, and Y.-N. Wang, “Lecture Notes on Generalized Symmetries and Applications”, in (July 2023), [arXiv:2307.09215 \[hep-th\]](#).
- [6] S.-H. Shao, “What’s Done Cannot Be Undone: TASI Lectures on Non-Invertible Symmetry”, (2023), [arXiv:2308.00747 \[hep-th\]](#).
- [7] T. Yuji, “Lecture on anomalies and topological phases”, (2019), <https://member.ipmu.jp/yuji.tachikawa/lectures/2019-top-anom/>.