

SIG Proceedings Paper in LaTeX Format*

Extended Abstract[†]

Lars Thørväld[‡]
The Thørväld Group
Hekla, Iceland
larst@affiliation.org

ABSTRACT

This paper provides a sample of a \LaTeX document which conforms, somewhat loosely, to the formatting guidelines for ACM SIG Proceedings.¹

CCS CONCEPTS

• **Computer systems organization** → **Embedded systems**; *Redundancy*; Robotics; • **Networks** → Network reliability;

KEYWORDS

ACM proceedings, \LaTeX , text tagging

ACM Reference Format:

Lars Thørväld. 1997. SIG Proceedings Paper in LaTeX Format: Extended Abstract. In *Proceedings of ACM Woodstock conference (WOODSTOCK'97)*, Jennifer B. Sartor, Theo D'Hondt, and Wolfgang De Meuter (Eds.). ACM, New York, NY, USA, Article 4, 12 pages. https://doi.org/10.475/123_4

1 PATHS AND ALIASING

Model dynamics using judgement of the form:

$$\boxed{V, H, R, F \vdash_{P, \Sigma} e \Downarrow v, H', F'}$$

Where $V : \text{Var} \rightarrow \text{Val} \times \text{State}$, $H : \text{Loc} \rightarrow \text{Val}$, $R \subseteq \text{Loc}$, $F \subseteq \text{Loc}$, and $\Sigma : \text{Var} \rightarrow \text{FTypes}$. This can be read as: under stack V , heap H , roots R , freelist F , and program P with signature Σ , the expression e evaluates to v , and engenders a new heap H' and freelist F' .

A *program* is then a Σ indexed map P from Var to pairs $(y_f, e_f)_{f \in \Sigma}$, where $\Sigma(y_f) = A \rightarrow B$, and $\Sigma; y_f : A \vdash e_f : B$ (typing rules are discussed in 6). We write $P : \Sigma$ to mean P is a program with signature Σ . Because the signature Σ for the mapping of function names to first order functions does not change during evaluation, we drop the subscript Σ from \vdash_Σ when the context of evaluation is clear. It is convenient to think of the evaluation judgement \vdash as being indexed by a family of signatures Σ 's,

*Produces the permission block, and copyright information

[†]The full version of the author's guide is available as `acmart.pdf` document

[‡]This author is the one who did all the really hard work.

¹This is an abstract footnote

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).

WOODSTOCK'97, July 1997, El Paso, Texas USA

© 2016 Copyright held by the owner/author(s).

ACM ISBN 123-4567-24-567/08/06...\$15.00

https://doi.org/10.475/123_4

BTypes	$\tau ::=$		
	<code>nat</code>	<code>nat</code>	natura
	<code>unit</code>	<code>unit</code>	unit
	<code>bool</code>	<code>bool</code>	boolea
	<code>prod($\tau_1; \tau_2$)</code>	$\tau_1 \times \tau_2$	produc
	<code>list(τ)</code>	$L(\tau)$	list
FTypes	$\rho ::=$		
	<code>arr($\tau_1; \tau_2$)</code>	$\tau_1 \rightarrow \tau_2$	first o
Exp	$e ::=$		
	<code>x</code>	<code>x</code>	variab
	<code>nat[n]</code>	\bar{n}	numbe
	<code>unit</code>	<code>()</code>	unit
	<code>T</code>	<code>T</code>	true
	<code>F</code>	<code>F</code>	false
	<code>if($x; e_1; e_2$)</code>	<code>if x then e_1 else e_2</code>	if
	<code>lam($x : \tau. e$)</code>	$\lambda x : \tau. e$	abstra
	<code>ap($f; x$)</code>	$f(x)$	applic
	<code>tpl($x_1; x_2$)</code>	$\langle x_1, x_2 \rangle$	pair
	<code>case($x_1, x_2. e_1$)</code>	<code>case $p \{ (x_1; x_2) \hookrightarrow e_1 \}$</code>	match
	<code>nil</code>	<code>[]</code>	nil
	<code>cons($x_1; x_2$)</code>	$x_1 :: x_2$	cons
	<code>case{l}($e_1; x. xs. e_2$)</code>	<code>case $l \{ \text{nil} \hookrightarrow e_1 \mid \text{cons}(x; xs) \hookrightarrow e_2 \}$</code>	match
	<code>let($e_1; x : \tau. e_2$)</code>	<code>let $x = e_1$ in e_2</code>	let
	<code>share($x; x_1, x_2. e$)</code>	<code>share x as x_1, x_2 in e</code>	share
Val	$v ::=$		
	<code>val(n)</code>	n	numen
	<code>val(T)</code>	<code>T</code>	true v
	<code>val(F)</code>	<code>F</code>	false v
	<code>val(Null)</code>	<code>Null</code>	null v
	<code>val(cl($V; x. e$))</code>	$(V, x. e)$	functi
	<code>val(l)</code>	l	loc val
	<code>val(pair($v_1; v_2$))</code>	$\langle v_1, v_2 \rangle$	pair v
State	$s ::=$		
	<code>alive</code>	<code>alive</code>	live v
	<code>dead</code>	<code>dead</code>	dead v
Loc	$l ::=$		
	<code>loc(l)</code>	l	locatio
Var	$l ::=$		
	<code>var(x)</code>	x	variab

each of which is a set of “top-level” first-order declarations to be used during evaluation.

For a partial map $f : A \rightarrow B$, we write dom for the defined values of f . Sometimes we shorten $x \in dom(f)$ to $x \in f$. We write $f[x \mapsto y]$ for the extension of f where x is mapped to y , with the constraint that $x \notin dom(f)$.

Roots represents the set of locations required to compute the continuation *excluding* the current expression. We can think of roots as the heap allocations necessary to compute the context with a hole that will be filled by the current expression.

In order prove soundness of the type system, we need some auxiliary judgements to defining properties of a heap. Below we define $reach : Val \rightarrow \{\{Loc\}\}$ that maps stack values its the root *multiset*, the multiset of locations that’s already on the stack.

Next we define reachability of values:

$$\begin{aligned} reach_H(\langle v_1, v_2 \rangle) &= reach_H(v_1) \uplus reach_H(v_2) \\ reach_H(l) &= \{l\} \uplus reach_H(H(l)) \\ reach_H(_) &= \emptyset \end{aligned}$$

For a multiset S , we write $\mu_S : S \rightarrow \mathbb{N}$ for the multiplicity function of S , which maps each element to the count of its occurrence. If $\mu_S(x) \geq 1$ for a multiset S , then we write $x \in S$ as in the usual set membership relation. If for all $s \in S$, $\mu(s) = 1$, then S is a property set, and we denote it by $set(S)$. Additionally, $A \uplus B$ denotes counting union of sets where $\mu_{A \uplus B}(s) = \mu_A(s) + \mu_B(s)$, and $A \cup B$ denotes the usual union where $\mu_{A \cup B}(s) = \max(\mu_A(s), \mu_B(s))$. For the disjoint union of sets A and B , we write $A \sqcup B$.

Next, we define the predicates `no_alias`, `stable`, and `disjoint`:

$$\begin{aligned} \text{no_alias}(V, H): \quad & \forall x, y \in V, x \neq y. \text{ Let } r_x = reach_H(V(x)), \\ & r_y = reach_H(V(y)). \text{ Then:} \\ (1) \quad & set(r_x), set(r_y) \\ (2) \quad & r_x \cap r_y = \emptyset \\ \text{stable}(R, H, H'): \quad & \forall l \in R. H(l) = H'(l). \\ \text{safe}(V, H, F): \quad & \forall x \in V. reach_H(V(x)) \cap F = \emptyset \\ \text{disjoint}(C): \quad & \forall X, Y \in C. X \cap Y = \emptyset \end{aligned}$$

For a stack V and a heap H , whenever `no_alias`(V, H) holds, visually, one can think of the situation as the following: the induced graph of heap H with variables on the stack as additional leaf nodes is a forest: a disjoint union of arborescences (directed trees); consequently, there is at most one path from a live variable on the stack V to a location in H by following the pointers.

First, we define $FV^*(e)$, the multiset of free variables of e . As the usual FV , it is defined inductively over the structure of e ; the only unusual thing is that multiple occurrences of a free variable x in e will be reflected in the multiplicity of $FV^*(e)$.

Next, we define $locs_{V,H}$ using the previous notion of reachability.

$$locs_{V,H}(e) = \bigcup_{x \in FV(e)} reach_H(V(x))$$

size calculates the *literal size* of a value, e.g. the size to store its address.

$$\begin{aligned} size(\langle v_1, v_2 \rangle) &= size(v_1) + size(v_2) \\ size(_) &= 1 \end{aligned}$$

Let $card(S)$ denote the number of unique elements, e.g. the cardinality of a multiset S . We write $\|v\|_H$ for $card(reach_H(v))$. As usual, we extend it to stacks V : $\|V\|_H = \sum_{V(x)=v} \|v\|_H$

$copy(H, L, v)$ takes a heap H , a set of locations L , and a value v , and returns a new heap H' and a location l such that l maps to v in H' .

$$\begin{aligned} copy(H, L, \langle v_1, v_2 \rangle) &= \\ \text{let } L_1 \sqcup L_2 &\subseteq L \\ \text{where } |L_1| &= \|v_1\|_H, |L_2| = \|v_2\|_H \\ \text{let } H_1, v'_1 &= copy(H, L_1, v_1) \\ \text{let } H_2, v'_2 &= copy(H_1, L_2, v_2) \text{ in} \\ H_2, \langle v'_1, v'_2 \rangle & \\ copy(H, L, l) &= \\ \text{let } l' \in L &\text{ in} \\ \text{let } H', v &= copy(H, L \setminus \{l'\}, H(l)) \text{ in} \\ H' \{l' \mapsto v\}, l' & \\ copy(H, L, v) &= \\ H, v & \end{aligned}$$

2 GARBAGE COLLECTION SEMANTICS

$$\frac{V(x) = v}{V, H, R, F \vdash x \Downarrow v, H, F} (S_1)$$

$$\frac{}{V, H, R, F \vdash \bar{n} \Downarrow \text{val}(n), H, F} (S_2)$$

$$\frac{}{V, H, R, F \vdash \top \Downarrow \text{val}(\top), H, F} (S_3)$$

$$\frac{}{V, H, R, F \vdash F \Downarrow \text{val}(F), H, F} (S_4)$$

$$\frac{}{V, H, R, F \vdash () \Downarrow \text{val}(\text{Null}), H, F} (S_5)$$

$$\frac{V = V'[x \mapsto \top] \quad g = \{l \in H \mid l \notin F \cup R \cup \text{locs}_{V, H}(e_1)\} \quad V', H, R, F \cup g \vdash e_1 \Downarrow v, H', F'}{V, H, R, F \vdash \text{if}(x; e_1; e_2) \Downarrow v, H', F'} (S_6)$$

$$\frac{V = V'[x \mapsto F] \quad g = \{l \in H \mid l \notin F \cup R \cup \text{locs}_{V, H}(e_2)\} \quad V', H, R, F \cup g \vdash e_2 \Downarrow v, H', F'}{V, H, R, F \vdash \text{if}(x; e_1; e_2) \Downarrow v, H', F'} (S_7)$$

$$\frac{V(x) = v' \quad P(f) = (y_f, e_f) \quad g = \{l \in H \mid l \notin F \cup R \cup \text{locs}_{V, H}(e_f)\} \quad [y_f \mapsto v'], H, R, F \cup g \vdash e_f \Downarrow v, H', F'}{V, H, R, F \vdash f(x) \Downarrow v, H', F'} (S_8)$$

$$\frac{V(x_1) = v_1 \quad V(x_2) = v_2}{V, H, R, F \vdash \langle x_1, x_2 \rangle \Downarrow \langle v_1, v_2 \rangle, H, F} (S_9)$$

$$\frac{V = V'[x \mapsto \langle v_1, v_2 \rangle] \quad g = \{l \in H \mid l \notin F \cup R \cup \text{locs}_{V, H}(e)\} \quad V'' = V'[x_1 \mapsto v_1, x_2 \mapsto v_2] \quad V'', H, R, F \cup g \vdash e \Downarrow v, H', F'}{V, H, R, F \vdash \text{case } x \{ (x_1; x_2) \hookrightarrow e \} \Downarrow v, H', F'} (S_{10})$$

$$\frac{}{V, H, R, F \vdash \text{nil} \Downarrow \text{val}(\text{Null}), H, F} (S_{11})$$

$$\frac{v = \langle V(x_1), V(x_2) \rangle \quad H' = H\{l \mapsto v\}}{V, H, R, F \vdash \text{cons}(x_1; x_2) \Downarrow l, H', F'} (S_{12})$$

$$\frac{V = V'[x \mapsto v'] \quad g = \text{reach}_H(v') \quad V', H, R, F \cup g \vdash e \Downarrow v, H', F'}{V, H, R, F \vdash \text{drop}(x; e) \Downarrow v, H', F'} (S_{13})$$

$$\frac{V = V'[x \mapsto v'] \quad L \subseteq F \quad |L| = \|v'\|_H \quad H', v'' = \text{copy}(H, L, v') \quad V'[x_1 \mapsto v', x_2 \mapsto v''], H', R, F \setminus L \vdash e \Downarrow v, H'', F'}{V, H, R, F \vdash \text{shareCopy } x \text{ as } x_1, x_2 \text{ in } e \Downarrow v, H'', F'} (S_{14})$$

$$\frac{V(x) = \text{Null} \quad V' \subseteq V \quad \text{dom}(V') = FV(e_1) \quad g = \{l \in H \mid l \notin F \cup R \cup \text{locs}_{V, H}(e_1)\} \quad V', H, R, F \cup g \vdash e_1 \Downarrow v, H', F'}{V, H, R, F \vdash \text{case } x \{ \text{nil} \hookrightarrow e_1 \mid \text{cons}(x_h; x_t) \hookrightarrow e_2 \} \Downarrow v, H', F'} (S_{15})$$

$$\frac{V(x) = l \quad H(l) = \langle v_h, v_t \rangle \quad V' \subseteq V \quad \text{dom}(V') = FV(e_2) \setminus \{x_h, x_t\} \quad V'' = V'[x_h \mapsto v_h, x_t \mapsto v_t] \quad g = \{l \in H \mid l \notin F \cup R \cup \text{locs}_{V'', H}(e_2)\} \quad V'', H, R, F \cup g \vdash e_2 \Downarrow v, H', F'}{V, H, R, F \vdash \text{case } x \{ \text{nil} \hookrightarrow e_1 \mid \text{cons}(x_h; x_t) \hookrightarrow e_2 \} \Downarrow v, H', F'} (S_{16})$$

3 OPERATIONAL SEMANTICS

In order to prove the soundness of the type system, we also define a simplified operational semantics that does not account for garbage collection.

$$\boxed{V, H \vdash e \Downarrow v, H'}$$

This can be read as: under stack V , heap H the expression e evaluates to v , and engenders a new heap H' . We write the representative rules.

$$\frac{v = \langle V(x_1), V(x_2) \rangle \quad (L \sqcup \{l\}) \cap \text{dom}(H) = \emptyset \quad H', l = \text{copy}(H, L, v)}{V, H \vdash \text{cons}(x_1; x_2) \Downarrow l, H'} \quad (\text{S}_{18})$$

$$\frac{V(x) = l \quad H(l) = \langle v_h, v_t \rangle \quad V' \subseteq V \quad \text{dom}(V') = FV(e_2) \setminus \{x_h, x_t\} \quad V'' = V'[x_h \mapsto v_h, x_t \mapsto v_t] \quad V'', H \vdash e_2 \Downarrow v, H'}{V, H \vdash \text{case } x \{ \text{nil} \hookrightarrow e_1 \mid \text{cons}(x_h; x_t) \hookrightarrow e_2 \} \Downarrow v, H'} \quad (\text{S}_{19})$$

$$\frac{V = V_1 \sqcup V_2 \quad \text{dom}(V_1) = FV(e_1) \quad \text{dom}(V_2) = FV(\text{lam}(x : \tau. e_2)) \quad V_1, H \vdash e_1 \Downarrow v_1, H_1 \quad V'_2 = V_2[x \mapsto v_1] \quad V'_2, H_1 \vdash e_2 \Downarrow v_2, H_2}{V, H \vdash \text{let}(e_1; x : \tau. e_2) \Downarrow v_2, H_2} \quad (\text{S}_{20})$$

$$\frac{V = V'[x \mapsto v'] \quad L \text{ fresh} \quad |L| = \|v'\|_H \quad H', v'' = \text{copy}(H, L, v') \quad V'[x_1 \mapsto v', x_2 \mapsto v''], H' \vdash e \Downarrow v, H''}{V, H \vdash \text{shareCopy } x \text{ as } x_1, x_2 \text{ in } e \Downarrow v, H''} \quad (\text{S}_{21})$$

4 WELL DEFINED ENVIRONMENTS

In order to define the potential for first-order types, we need a notion of well-define environments, one that relates heap values to semantic values of a type. We first give a denotational semantics for the first-order types:

$$\begin{aligned} () &\in \llbracket \text{unit} \rrbracket \\ \perp &\in \llbracket \text{bool} \rrbracket \\ \top &\in \llbracket \text{bool} \rrbracket \\ 0 &\in \llbracket \text{nat} \rrbracket \\ n + 1 &\in \llbracket \text{nat} \rrbracket \text{ if } n \in \llbracket \text{nat} \rrbracket \\ [] &\in \llbracket L(A) \rrbracket \\ \pi(a, l) &\in \llbracket L(A) \rrbracket \text{ if } a \in \llbracket A \rrbracket \text{ and } l \in \llbracket L(A) \rrbracket \end{aligned}$$

Where semantic set for each type is the least set such that the above holds. Note $\pi(x, y)$ is the usual set-theoretic pairing function, and write $[a_1, \dots, a_n]$ for $\pi(a_1, \dots, \pi(a_n, []))$.

Now we give the judgements relating heap values to semantic values, in the form $\boxed{H \models v \mapsto a : A}$, which can be read as: under heap H , heap value v defines the semantic value $a \in \llbracket A \rrbracket$.

$$\frac{n \in \mathbb{Z}}{H \models n \mapsto n : \text{nat}} \quad (\text{V:ConstI})$$

$$\frac{}{H \models \text{Null} \mapsto n : \text{unit}} \quad (\text{V:ConstI})$$

$$\frac{A \in \text{BType}}{H \models \text{Null} \mapsto n : L(A)} \quad (\text{V:Nil})$$

$$\frac{}{H \models \top \mapsto \top : \text{bool}} \quad (\text{V:True}) \quad \frac{}{H \models \text{F} \mapsto \perp : \text{bool}} \quad (\text{V:False})$$

$$\frac{l \in \text{Loc} \quad H(l) = \langle v_h, v_t \rangle \quad H \models v_h \mapsto a_1 : A \quad H \models v_t \mapsto [a_2, \dots, a_n] : L(A)}{H \models l \mapsto [a_1, \dots, a_n] : L(A)} \quad (\text{V:Cons})$$

5 STACK VS HEAP ALLOCATED TYPES

In order to share variables, we need to distinguish between types that are allocated on the stack and the heap. We write $\boxed{\text{stack}(A)}$ to denote that values of type A will be allocated *entirely* on the stack at run time (no references into the heap).

$$\frac{A \in \{\text{unit}, \text{bool}, \text{nat}\}}{\text{stack}(A)} \quad (\text{S:Const})$$

$$\frac{\text{stack}(A_1) \quad \text{stack}(A_2)}{\text{stack}(A_1 \times A_2)} \quad (\text{S:Product})$$

6 LINEAR GARBAGE COLLECTION TYPE RULES

The linear version of the type system takes into account of garbage collected cells by returning potential locally in a match construct. Since we are interested in the number of heap cells, all constants are assumed to be nonnegative. The second let rule expresses the fact that since stack types don't reference heap cells, any heap cells used in the evaluation of e_1 can be deallocated, as there are no longer references to them in v_1 .

$$\frac{n \in \mathbb{Z}}{\Sigma; \emptyset \mid \frac{q}{q} n : \text{nat}} \text{ (L:ConstI)} \quad \frac{}{\Sigma; \emptyset \mid \frac{q}{q} () : \text{unit}} \text{ (L:ConstU)}$$

$$\frac{}{\Sigma; \emptyset \mid \frac{q}{q} \top : \text{bool}} \text{ (L:ConstT)} \quad \frac{}{\Sigma; \emptyset \mid \frac{q}{q} \text{F} : \text{bool}} \text{ (L:ConstF)}$$

$$\frac{}{\Sigma; x : B \mid \frac{q}{q} x : B} \text{ (L:Var)} \quad \frac{\Sigma(f) = A \xrightarrow{q/q'} B}{\Sigma; x : A \mid \frac{q}{q'} f(x) : B}$$

$$\frac{\Sigma; \Gamma \mid \frac{q}{q'} e_t : B \quad \Sigma; \Gamma \mid \frac{q}{q'} e_f : B}{\Sigma; \Gamma, x : \text{bool} \mid \frac{q}{q'} \text{if } x \text{ then } e_t \text{ else } e_f : B} \text{ (L:Cond)}$$

$$\frac{}{\Sigma; x_1 : A_1, x_2 : A_2 \mid \frac{q}{q} \langle x_1, x_2 \rangle : A_1 \times A_2} \text{ (L:Pair)}$$

$$\frac{\Sigma; \Gamma, x_1 : A_1, x_2 : A_2 \mid \frac{q}{q'} e : B}{\Sigma; \Gamma, x : (A_1, A_2) \mid \frac{q}{q'} \text{case } x \{ (x_1, x_2) \hookrightarrow e \} : B} \text{ (L:MatP)}$$

$$\frac{}{\Sigma; \emptyset \mid \frac{q}{q} \text{nil} : L^P(A)} \text{ (L:Nil)}$$

$$\frac{}{\Sigma; x_h : A, x_t : L^P(A) \mid \frac{q+p+1}{q} \text{cons}(x_h; x_t) : L^P(A)} \text{ (L:Cons)}$$

$$\frac{\Sigma; \Gamma \mid \frac{q}{q'} e_1 : B \quad \Sigma; \Gamma, x_h : A, x_t : L^P(A) \mid \frac{q+p+1}{q'} e_2 : B}{\Sigma; \Gamma, x : L^P(A) \mid \frac{q}{q'} \text{case } x \{ \text{nil} \hookrightarrow e_1 \mid \text{cons}(x_h; x_t) \hookrightarrow e_2 \} : B} \text{ (L:MatL)}$$

$$\frac{\Sigma; \Gamma_1 \mid \frac{q}{p} e_1 : A \quad \Sigma; \Gamma_2, x : A \mid \frac{p}{q'} e_2 : B}{\Sigma; \Gamma_1, \Gamma_2 \mid \frac{q}{q'} \text{let}(e_1; x : \tau.e_2) : B} \text{ (L:Let)}$$

$$\frac{\Sigma; \Gamma_1 \mid \frac{q}{p} e_1 : A \quad \text{stack}(A) \quad \Sigma; \Gamma_2, x : A \mid \frac{\max(p, q)}{q'} e_2 : B}{\Sigma; \Gamma_1, \Gamma_2 \mid \frac{q}{q'} \text{let}(e_1; x : \tau.e_2) : B} \text{ (L:LetS)}$$

$$\frac{\Sigma; \Gamma \mid \frac{q}{q'} e : B}{\Sigma; \Gamma, x : A \mid \frac{q}{q'} \text{drop}(x; e) : B} \text{ (L:Drop)}$$

$$\frac{\text{stack}(A) \quad \Sigma; \Gamma, x_1 : A, x_2 : A \mid \frac{q}{q'} e : B}{\Sigma; \Gamma, x : A \mid \frac{q}{q'} \text{share } x \text{ as } x_1, x_2 \text{ in } e : B} \text{ (L:Share)}$$

Now if we take $\dagger : L^P(A) \mapsto L(A)$ as the map that erases resource annotations, we obtain a simpler typing judgement

$$\boxed{\Sigma^\dagger; \Gamma^\dagger \vdash e : B^\dagger}.$$

7 TYPE RULES FOR SHARING

$$\begin{aligned} L^P(A) - n &= L^{\max(p-n, 0)}(A - n) \\ A_1 \times A_2 - n &= A_1 - n \times A_2 - n \\ A - n &= A \end{aligned}$$

$$\frac{A \curlyvee A_1, A_2, 1 \quad \Sigma; \Gamma, x_1 : A_1, x_2 : A_2 \mid \frac{q}{q'} e : B}{\Sigma; \Gamma, x : A \mid \frac{q}{q'} \text{share } x \text{ as } x_1, x_2 \text{ in } e : B} \text{ (M:Share)}$$

$$\frac{\Sigma; \Gamma_1 \mid \frac{q}{p} e_1 : A \quad \Sigma; \Gamma_2, x : (A' - 1) \mid \frac{p}{q'} e_2 : B}{\Sigma; \Gamma_1, \Gamma_2 \mid \frac{q}{q'} \text{let}(e_1; x : \tau.e_2) : B} \text{ (M:Let)}$$

Where $A \curlyvee A_1, A_2, n$ is the sharing relation defined as:

$$\begin{aligned} L^P(A) \curlyvee L^q(A_1), L^r(A_2), n & \quad \text{if } p = q + r + n \text{ and } A \curlyvee A_1, A_2, n \\ A \curlyvee A_1, A_2 & \quad \text{if } \text{stack}(A), \text{stack}(A_1), \text{stack}(A_2) \text{ and } A \equiv A_1 \equiv A_2 \end{aligned}$$

8 SOUNDNESS FOR LINEAR GC

We simplify the soundness proof of the type system for the general metric to one with monotonic resource. (No function types for now)

Definition 8.1 (Well-formed computation). When considering the input mode arguments of a evaluation judgment $V, H, R, F \vdash e \Downarrow v, H', F'$, we say the 5-tuple (V, H, R, F, e) is a *well-formed computation* given the following:

- (1) $\text{dom}(V) = FV(e)$
- (2) $\text{no_alias}(V, H)$, and
- (3) $\text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$

And we write $\text{wfc}(V, H, R, F, e)$ to denote this fact.

LEMMA 1.1. *If $\Sigma; \Gamma \mid \frac{q}{q'} e : B$, then $\Sigma^\dagger; \Gamma^\dagger \vdash e : B^\dagger$.*

LEMMA 1.2. *If $\Sigma; \Gamma \mid \frac{q}{q'} e : B$, then $\text{set}(FV^\star(e))$ and $\text{dom}(\Gamma) = FV(e)$.*

PROOF. Induction on the typing judgement. \square

LEMMA 1.3. *Let $H \models v \mapsto a : A$. For all sets of locations R , if $\text{reach}_H(v) \subseteq R$ and $\text{stable}(R, H, H')$, then $H' \models v \mapsto a : A$ and $\text{reach}_{H'}(v) = \text{reach}_H(v)$.*

PROOF. Induction on the structure of v . \square

COROLLARY 1.3.1. *Let $H \models V : \Gamma$. For all sets of locations R , if $\bigcup_{x \in V} \text{reach}_H(V(x)) \subseteq R$ and $\text{stable}(R, H, H')$, then $H' \models V : \Gamma$.*

PROOF. Follows from Lemma 1.3. \square

LEMMA 1.4. *Let $H \models v \mapsto a : A$. If $\text{stack}(A)$, then $\Phi_H(v : A) = 0$.*

PROOF. Induction on $H \models v \mapsto a : A$. \square

LEMMA 1.5 (HEAP CONSERVATION). *Let $\text{wfc}(V, H, R, F, e), V, H, R, F \vdash e \Downarrow v, H', F'$, and $g = \text{gc}(H', R, F')$. Then $\|V\|_H + |F| \leq \|v\|_{H'} + |F' \cup g|$.*

PROOF. Induction on evaluation.

Case 1: E:Var

$$\begin{aligned} V &= [x \mapsto v] && (\text{since } \text{dom}(V) = FV(e) = \{x\}) \\ \|V\|_H &= \|v\|_H && (\text{def of } \|\cdot\|_H) \\ \|V\|_H + |F| &\leq \|v\|_{H'} + |F \cup g| \end{aligned}$$

Case 2: E:Const* Due to similarity, we show only for E:ConstI

$$\begin{aligned} V &= \emptyset && (\text{since } \text{dom}(V) = FV(e) = \emptyset) \\ \|V\|_H &= \|v\|_H && (\text{def of } \|\cdot\|_H) \\ \|V\|_H + |F| &\leq \|v\|_{H'} + |F \cup g| \end{aligned}$$

Case 4: E:App

Case 5: E:CondT Similar to E:MatNil

Case 6: E:CondF Similar to E:CondT

Case 7: E:Let

$$\|V_1\|_H + |F| \leq \|v_1\|_{H_1} + |F_1 \cup g| \quad (\text{IH on first premise})$$

Let $g' = \text{gc}(H_2, R, F_2)$

$$\|V_2'\|_{H_1} + |F_1 \cup g| \leq \|v_2\|_{H_2} + |F \cup g'| \quad (\text{IH on second premise})$$

$$\begin{aligned} \|V_2'\|_{H_1} &= \|V_2\|_{H_1} + \|v_1\|_{H_1} && (\text{definition of semantic size}) \\ &= \|V_2\|_H + \|v_1\|_{H_1} && (\text{main lemma}) \end{aligned}$$

$$\|V_2\|_H + \|v_1\|_{H_1} + |F_1 \cup g| \leq \|v_2\|_{H_2} + |F \cup g'|$$

$$\|V_1\|_H + \|V_2\|_H + \|v_1\|_{H_1} + |F| + |F_1 \cup g| \leq \|v_1\|_{H_1} + \|v_2\|_{H_2} + |F_1 \cup g| + |F|$$

$$\|V\|_H + |F| \leq \|v_2\|_{H_2} + |F \cup g'|$$

Case 8: E:Pair Similar to E:Var

Case 9: E:MatP Similar to E:MatCons

Case 10: E:Nil Similar to E:Const*

Case 11: E:Cons

$V = [x_1 \mapsto v_1, x_2 \mapsto v_2]$ (since $\text{dom}(V) = FV(e) = \{x_1, x_2\}$)

$$\|V\|_H = \|v_1\|_H + \|v_2\|_H \quad (\text{def of } \|\cdot\|_H)$$

$$\|l\|_{H'} = 1 + \|H'(l)\|_{H'} = 1 + \|v\|_{H''} = 1 + \|v_1\|_{H''} + \|v_2\|_{H''} \quad (\text{def of semantic size})$$

$$= 1 + \|v_1\|_H + \|v_1\|_H$$

$$= 1 + \|V\|_H$$

$$L \sqcup \{l\} \subseteq g \quad (R \cap F = \emptyset \text{ and } L \sqcup \{l\} \subseteq H'')$$

$$|g| \geq |L \sqcup \{l\}| = \text{size}(v) + 1$$

$$|F' \cup g| \geq |F|$$

$$\|V\|_H + |F| \leq \|v\|_{H'} + |F \cup g|$$

Case 12: E:MatNil

Case 13: E:MatCons

Let $g' = \text{gc}(H', R, F')$

$$\|V''\|_H + |F \cup g| \leq |F' \cup g'| \quad (\text{IH (wfc from main lemma)})$$

$$\|V''\|_H = \|V'[x_h \mapsto v_h, x_t \mapsto v_t]\|_H$$

$$= \|V'\|_H + \|v_h\|_H + \|v_t\|_H$$

$$= \|V'\|_H + \|l\|_H - 1$$

$$= \|V\|_H - 1$$

$$\|V\|_H - 1 + |F \cup g| \leq |F' \cup g'|$$

$$\|v\|_H - 1 + |F| + |g| \leq |F' \cup g'| \quad (F \cap g = \emptyset)$$

$$\|v\|_H + |F| \leq |F' \cup g'| \quad (|g| \geq 1 \text{ from main lemma})$$

Case 13: E:Drop

Let $g' = \text{gc}(H', R, F')$

$$\|V'\|_H + |F \cup g| \leq \|v\|_{H'} + |F' \cup g'| \quad (\text{IH})$$

$$HV = \|v\|_{V'} \parallel$$

$$v' \|V\|_H - \|v\|_{v'} |F \cup \text{reach}_H(v')| \leq \|v\|_{H'} + |F' \cup g'|$$

$$\|V\|_H - \|v\|_{v'} |F| + |\text{reach}_H(v')| \leq \|v\|_{H'} + |F' \cup g'|$$

$$\|V\|_H + |F| \leq \|v\|_{H'} + |F' \cup g'|$$

Case 13: E:ShareCopy

$e = \text{shareCopy } x \text{ as } x_1, x_2 \text{ in } e$ (case)

Let $g' = \text{gc}(H', R, F')$

$$\|V'[x_1 \mapsto v', x_2 \mapsto v'']\|_{H'} + |F \setminus L| \leq \|v\|_{H''} + |F' \cup g'|$$

(IH, well-formedness from main lemma)

$$\|V'[x_1 \mapsto v', x_2 \mapsto v'']\|_{H'} + |F \setminus L| = \|V\|_H + \|v''\|_{H'} + |F| - |L|$$

(stability lemma for copy)

$$= \|V\|_H + |L| + |F| - |L|$$

(lemma about copy)

$$= \|V\|_H + |F|$$

□

LEMMA 1.6. Let $\Sigma; \Gamma \vdash \frac{q}{q'} e : B$ and $V, H, R, F \vdash e \Downarrow v, H', F'$. Then $\|V\|_H - \|v\|_{H'} + q \geq q'$.

LEMMA 1.7 (MAIN LEMMA). For all stacks V and heaps H , let $V, H, R, F \vdash e \Downarrow v, H', F'$ and $\Sigma; \Gamma \vdash e : B$. Then given the following:

- (1) $\text{dom}(V) = FV(e)$
- (2) $\text{no_alias}(V, H)$, and
- (3) $\text{disjoint}(\{R, F, \text{locsv}_{V,H}(e)\})$

We have the following:

- (1) $\text{set}(\text{reach}_{H'}(v))$
- (2) $\text{disjoint}(\{R, F', \text{reach}_{H'}(v)\})$, and
- (3) $\text{stable}(R, H, H')$

PROOF. Nested induction on the evaluation judgement and the typing judgement.

Case 1: E:Var

Suppose $H \models V : \Gamma, \text{dom}(V) = FV(e), \text{no_alias}(V, H), \text{disjoint}(\{R, F, \text{locsv}_{V,H}(e)\})$
 $\text{set}(\text{reach}_H(v))$ (no_alias(V, H))
 $\text{disjoint}(\{R, F, \text{reach}_H(v)\})$ (disjoint({R, F, locsv_{V,H}(e)}))
 $\text{no_alias}(V, H)$ (Sp.)
 $\text{stable}(R, H, H')$ ($H = H'$)

Case 2: E:Const* Due to similarity, we show only for E:ConstI

Suppose $H \models V : \Gamma, \text{dom}(V) = FV(e), \text{no_alias}(V, H), \text{disjoint}(\{R, F, \text{locsv}_{V,H}(e)\})$
 $\text{set}(\text{reach}_H(v))$ ($\text{reach}_H(v) = \emptyset$)
 $\text{disjoint}(\{R, F, \emptyset\})$ (disjoint(R, F))
 $\text{no_alias}(V, H)$ (Sp.)
 $\text{stable}(R, H, H')$ ($H = H'$)

Case 4: E:App

Case 5: E:CondT Similar to E:MatNil

Case 6: E:CondF Similar to E:CondT

Case 7: E:Let

$V, H, R, F \vdash \text{let}(e_1; x : \tau.e_2) \Downarrow v_2, H_2, F_2$ (case)

$V, H, R', F \vdash e_1 \Downarrow v_1, H_1, F_1$ (ad.)

$\Sigma; \Gamma_1, \Gamma_2 \vdash \text{let}(e_1; x : \tau.e_2) : B$ (case)

$\Sigma; \Gamma_1 \vdash e_1 : A$ (ad.)

Suppose $H \models V : \Gamma, \text{dom}(V) = FV(e), \text{no_alias}(V, H), \text{disjoint}(\{R, F, \text{locsv}_{V,H}(e)\})$

$H \models V_1 : \Gamma_1$ (def of W.D.E and Lemma 1.2)

By IH, we have invariant on J_1

NTS (1) - (3) to instantiate invariant on J_1

(1) $\text{dom}(V_1) = FV(e_1)$ (def of V_1)

(2) $\text{no_alias}(V_1, H)$ (no_alias(V, H) and $V_1 \subseteq V$)

(3) $\text{disjoint}(R', F, \text{locsv}_{V,H}(e_1))$

$$F \cap R' = \emptyset$$

$$(F \cap \text{locsv}_{V,H}(e) = \emptyset \text{ and } \text{locsv}_{V_2,H}(\text{lam}(x : \tau.e_2)) \subseteq \text{locsv}_{V,H}(e))$$

$$FV(e_1) \cap FV(\text{lam}(x : \tau.e_2)) = \emptyset$$
 (Lemma 1.2)

$$\text{locsv}_{V,H}(e_1) \cap \text{locsv}_{V_2,H}(\text{lam}(x : \tau.e_2)) = \emptyset$$
 (no_alias(V, H))

$$R' \cap \text{locsv}_{V,H}(e_1) = \emptyset$$
 (disjoint({R, locsv_{V,H}(e)}))

$$F \cap \text{locsv}_{V,H}(e_1) = \emptyset$$
 (Sp.)

Thus we have $\text{disjoint}(R', F, \text{locsv}_{V,H}(e_1))$

By IH,

(1) $\text{set}(\text{reach}_{H_1}(v_1))$

(2) $\text{disjoint}(\{R', F_1, \text{reach}_{H_1}(v_1)\})$

(3) $\text{stable}(R', H, H_1)$

$$V'_2, H_1, R, F_1 \cup g \vdash e_2 \Downarrow v_2, H_2, F_2$$
 (ad.)

$$\Sigma; \Gamma_2, x : A \vdash e_2 : B$$
 (ad.)

$$H_1 \models V'_2 : (\Gamma_2, x : A)$$
 (???)

By IH, we have invariant on J_2

NTS (1) - (3) to instantiate invariant on J_2

(1) $\text{dom}(V'_2) = FV(e_2)$ (def of V'_2)

(2) $\text{no_alias}(V'_2, H_1)$

Let $x_1, x_2 \in V'_2, x_1 \neq x_2$ be arb.

case: $x_1 = x, x_2 \neq x$

$$\text{reach}_H(V'_2(x_1)) \subseteq R'$$

$$(\text{reach}_H(V'_2(x_1)) \subseteq \text{locsv}_{V'_2,H}(\text{lam}(x : \tau.e_2)))$$

$$\text{reach}_H(V'_2(x_2)) \subseteq R'$$

$$(\text{reach}_H(V'_2(x_2)) \subseteq \text{locsv}_{V'_2,H}(\text{lam}(x : \tau.e_2)))$$

$$\text{reach}_H(V'_2(x_1)) = \text{reach}_{H_1}(V'_2(x_1)), \text{reach}_H(V'_2(x_2)) = \text{reach}_{H_1}(V'_2(x_2))$$

$$(\text{stable}(R', H, H_1) \text{ and Lemma 1.3})$$

$$\text{reach}_{H_1}(V'_2(x_1)) = \text{reach}_H(V(x_1)), \text{reach}_{H_1}(V'_2(x_2)) = \text{reach}_H(V(x_2))$$

$$(\text{stable}(R', H, H_1) \text{ and Lemma 1.3})$$

$$\text{no_alias}(V'_2, H_1)$$
 (no_alias(V, H))

case: $x_1 = x, x_2 \neq x$

$$\text{reach}_{H_1}(V'_2(x_1)) = \text{reach}_{H_1}(v_1)$$
 (def of V'_2)

$$\text{reach}_{H_1}(V'_2(x_2)) \subseteq R'$$
 (same as above)

$$\text{set}(\text{reach}_{H_1}(v_1))$$
 (IH 1.1)

$$\text{reach}_{H_1}(V'_2(x_2)) = \text{reach}_H(V(x_2))$$
 (same as above)

$$\text{set}(\text{reach}_{H_1}(V'_2(x_2)))$$
 (no_alias(V, H))

$$\text{reach}_{H_1}(V'_2(x_1)) \cap \text{reach}_{H_1}(V'_2(x_2)) = \emptyset$$

$$(\text{disjoint}(\{R', \text{reach}_{H_1}(v_1)\}))$$

Thus we have $\text{no_alias}(V'_2, H_1)$
(3) $\text{disjoint}(\{R, F_1 \cup g, \text{locs}_{V'_2, H_1}(e_2)\})$
 $R \cap F_1 = \emptyset$ (disjoint($\{R', F_1\}$) from 1.2 and $R \subseteq R'$)
 $R \cap (F_1 \cup g) = \emptyset$ (def of g)
NTS $(F_1 \cup g) \cap \text{locs}_{V'_2, H_1}(e_2) = \emptyset$
Let $l \in \text{locs}_{V'_2, H_1}(e_2)$ be arb.
 $l \in \text{reach}_{H_1}(V'_2(x'))$ for some $x' \in V'_2$
case: $x' \neq x$
 $\text{reach}_H(V_2(x')) = \text{reach}_{H_1}(V'_2(x'))$ (same as above)
 $\text{reach}_{H_1}(V'_2(x')) \subseteq R'$ (def of R')
 $\text{reach}_{H_1}(V'_2(x')) \cap F_1 = \emptyset$ (disjoint($\{R', F_1\}$) from 1.2)
case: $x' = x$
 $\text{reach}_{H_1}(V'_2(x')) = \text{reach}_{H_1}(v_1)$ (def of V'_2)
 $\text{reach}_{H_1}(V'_2(x')) \cap F_1 = \emptyset$
(disjoint($\{F_1, \text{reach}_{H_1}(v_1)\}$) from 1.2)
 $\text{reach}_{H_1}(V'_2(x')) \subseteq \text{locs}_{V'_2, H_1}(e_2)$ (def of $\text{locs}_{V, H}$)
 $\text{reach}_{H_1}(V'_2(x')) \cap g = \emptyset$ (def of g)
Thus $\text{reach}_{H_1}(V'_2(x')) \cap (F_1 \cup g) = \emptyset$
NTS $R \cap \text{locs}_{V'_2, H_1}(e_2) = \emptyset$
Let $l \in \text{locs}_{V'_2, H_1}(e_2)$ be arb.
 $l \in \text{reach}_{H_1}(V'_2(x'))$ for some $x' \in V'_2$
case: $x' \neq x$
 $\text{reach}_H(V_2(x')) = \text{reach}_{H_1}(V'_2(x'))$ (same as above)
 $l \in \text{locs}_{V, H}(e)$ (def of $\text{locs}_{V, H}$)
 $l \notin R$ (disjoint($\{R, \text{locs}_{V, H}(e)\}$) from 0.3)
case: $x' = x$
 $\text{reach}_{H_1}(V'_2(x')) = \text{reach}_{H_1}(v_1)$ (def of V'_2)
 $\text{reach}_{H_1}(V'_2(x')) \cap R = \emptyset$
(disjoint($\{R', \text{reach}_{H_1}(v_1)\}$) from 1.2 and $R \subseteq R'$)
Thus $\text{reach}_{H_1}(V'_2(x')) \cap R = \emptyset$
Hence we have (3) $\text{disjoint}(R, F_1 \cup g, \text{locs}_{V'_2, H_1}(e_2))$
By instantiating the invariant on J_2 , we have
(1) $\text{set}(\text{reach}_{H_2}(v_2))$
(2) $\text{disjoint}(\{R, F_2, \text{reach}_{H_2}(v_2)\})$
(3) $\text{stable}(R, H_1, H_2)$
Lastly, showing (1) - (3) holds for the original case J_0 :
(1) $\text{set}(\text{reach}_{H_2}(v_2))$ (By 2.1)
(2) $\text{disjoint}(\{R, F_2, \text{reach}_{H_2}(v_2)\})$ (By 2.2)
(3) $\text{stable}(R, H_1, H_2)$
Let $l \in R$ be arb.
 $H(l) = H_1(l)$ (stable(R', H, H_1) from 1.3)
 $H_1(l) = H_2(l)$ (stable(R, H_1, H_2) from 2.3)
 $H(l) = H_2(l)$
Hence $\text{stable}(R, H, H_2)$

Case 8: E:Pair Similar to E:Var

Case 9: E:MatP Similar to E:MatCons

Case 10: E:Nil Similar to E:Const*

Case 11: E:Cons

$V, H, R, F \vdash e \Downarrow l, H'', F'$ (case)

Suppose $H \models V : \Gamma, \text{dom}(V) = FV(e), \text{no_alias}(V, H), \text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$

NTS (1) - (3) holds after evaluation

(1) $\text{set}(\text{reach}_{H''}(l))$

$\text{stable}(\{\text{locs}_{V, H}(e)\}, H, H'')$

(disjoint($\{F, \text{locs}_{V, H}(e)\}$) and *copy* only updates $l \in L \subseteq F$)

$\text{reach}_H(V(x_i)) = \text{reach}_{H''}(V(x_i))$

($\text{reach}_H(V(x_i)) \subseteq \text{locs}_{V, H}(e)$ and 1.3 for $i = 1, 2$)

$\text{reach}_{H''}(l) = \{l\} \cup \text{reach}_{H''}(V(x_1)) \cup \text{reach}_{H''}(V(x_2))$

(def of reach_H)

$\text{set}(\text{reach}_{H''}(l))$ ($l \notin \text{locs}_{V, H}(e)$ and $\text{no_alias}(V, H)$)

(2) $\text{disjoint}(\{R, F', \text{reach}_{H''}(l)\})$

$R \cap F' = \emptyset$ ($F' \subseteq F$ and $\text{disjoint}(\{R, F\})$)

$R \cap \text{reach}_{H''}(l) = \emptyset$ ($l \in F$ and $\text{disjoint}(\{R, \text{locs}_{V, H}(e)\})$)

$F' \cap \text{reach}_{H''}(l) = \emptyset$ ($F' \subseteq F$ and $\text{disjoint}(\{F, \text{locs}_{V, H}(e)\})$)

Thus we have (2) $\text{disjoint}(\{R, F', \text{reach}_{H''}(l)\})$

(3) $\text{stable}(R, H, H'')$

(since *copy* only updates $l \in L \subseteq F$ and $F \cap R = \emptyset$)

Case 12: E:MatNil

Suppose $H \models V : \Gamma, \text{dom}(V) = FV(e), \text{no_alias}(V, H), \text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$

$\Sigma; \Gamma' \vdash e_1 : B$ (ad.)

$V, H, R, F \cup g \vdash e_1 \Downarrow v, H', F'$ (ad.)

$H \models V' : \Gamma'$ (def of W.D.E)

By IH, we have invariant on J_1

NTS (1) - (3) to instantiate invariant on J_1

(1) $\text{dom}(V') = FV(e_1)$ (def of V')

(2) $\text{no_alias}(V', H)$ ($\text{no_alias}(V, H)$ and $V' \subseteq V$)

(3) $\text{disjoint}(\{R, F, \text{locs}_{V', H}(e_1)\})$

(disjoint($\{R, F, \text{locs}_{V, H}(e)\}$) and $\text{locs}_{V', H}(e_1) \subseteq \text{locs}_{V, H}(e)$)

Instantiating invariant on J_1 ,

(1) $\text{set}(\text{reach}_{H'}(v))$

(2) $\text{disjoint}(\{R, F_1, \text{reach}_{H'}(v)\})$

(3) $\text{stable}(R, H, H')$

Case 13: E:MatCons

$V(x) = l$ (ad.)

$H(l) = \langle v_h, v_t \rangle$ (ad.)

$\Gamma = \Gamma', x : L(A)$ (ad.)

$\Sigma; \Gamma', x_h : A, x_t : L(A) \vdash e_2 : B$ (ad.)

$V'', H, R, F \cup g \vdash e_2 \Downarrow v_2, H_2, F'$ (ad.)

Suppose $H \models V : \Gamma, \text{dom}(V) = FV(e), \text{no_alias}(V, H), \text{disjoint}(\{F, R, \text{locs}_{V, H}(e)\})$

$H \models V(x) : L(A)$ (def of W.D.E)

$$H'' \models v_h : A, H'' \models v_t : L(A) \quad (\text{ad.})$$

$$H \models v_h : A, H \models v_t : L(A) \quad (???)$$

$$H \models V'' : \Gamma', x_h : A, x_t : L(A) \quad (\text{def of W.D.E})$$

By IH, we have invariant on J_1

NTS (1) - (3) to instantiate invariant on J_1

$$(1) \quad \text{dom}(V'') = FV(e_2) \quad (\text{def of } V'')$$

$$(2) \quad \text{no_alias}(V'', H)$$

Let $x_1, x_2 \in V'', x_1 \neq x_2, r_{x_1} = \text{reach}_H(V''(x_1)), r_{x_2} = \text{reach}_H(V''(x_2))$

case: $x_1 \notin \{x_h, x_t\}, x_2 \notin \{x_h, x_t\}$

$$(1), (2) \text{ from } \text{no_alias}(V, H)$$

case: $x_1 = x_h, x_2 \notin \{x_h, x_t\}$

$$\text{set}(r_{x_1}) \quad (\text{since } \text{set}(\text{reach}_H(V(x))) \text{ from } \text{no_alias}(V, H))$$

$$\text{set}(r_{x_2}) \quad (\text{since } \text{no_alias}(V, H))$$

$$x_2 \in FV(e) \quad (\text{def of } FV)$$

$$\text{reach}_H(V(x)) \cap r_{x_2} = \emptyset \quad (\text{def of } \text{reach} \text{ and } \text{no_alias}(V, H))$$

$$\text{hence } r_{x_1} \cap r_{x_2} = \emptyset$$

case: $x_1 = x_h, x_2 = x_t$

$$\text{set}(r_{x_1}) \text{ since } \text{set}(\text{reach}_H(V(x))) \text{ from } \text{no_alias}(V, H)$$

$$\text{set}(r_{x_2}) \text{ since } \text{set}(\text{reach}_H(V(x))) \text{ from } \text{no_alias}(V, H)$$

$$r_{x_1} \cap r_{x_2} = \emptyset \quad (\text{set}(\text{reach}_H(V(x))))$$

case: otherwise

similar to the above

Thus we have $\text{no_alias}(V'', H)$

$$(3) \quad \text{disjoint}(\{R, F \cup g, \text{locs}_{V'', H}(e_2)\})$$

$$(F \cup g) \cap R = \emptyset \quad (\text{since } F \cap R = \emptyset \text{ and by def of } g)$$

$$\text{NTS } R \cap \text{locs}_{V'', H}(e_2) = \emptyset$$

Let $l' \in \text{locs}_{V'', H}(e_2)$ be arb.

case: $l' \in \text{reach}_H(V''(x'))$ for some $x' \in FV(e_2)$ where $x' \notin \{x_h, x_t\}$

$$x' \in V \quad (\text{def of } V'')$$

$$l' \in \text{reach}_H(V(x'))$$

$$x' \in FV(e) \quad (\text{def of } FV)$$

$$l' \in \text{locs}_{V, H}(e) \quad (\text{def of } \text{locs}_{V, H})$$

$$l' \notin R \quad (\text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\}))$$

case: $l' \in \text{reach}_H(V''(x_h))$

tom $l' \in \text{reach}_H(v_h)$

$$l' \in \text{reach}_H(V(x)) \quad (\text{def of } \text{reach})$$

$$l' \in \text{locs}_{V, H}(e) \quad (\text{def of } \text{locs}_{V, H})$$

$$l' \notin R \quad (\text{since } \text{disjoint}(\{F, R, \text{locs}_{V, H}(e)\}))$$

case: $l' \in \text{reach}_H(V''(x_t))$

similar to above

$$\text{Hence } R \cap \text{locs}_{V'', H}(e_2) = \emptyset$$

$$F \cap \text{locs}_{V'', H}(e_2) = \emptyset \quad (\text{Similar to above})$$

$$g \cap \text{locs}_{V'', H}(e_2) = \emptyset \quad (\text{def. of } g)$$

$$(F \cup g) \cap \text{locs}_{V'', H}(e_2) = \emptyset$$

$$\text{Thus } \text{disjoint}(\{R, F \cup g, \text{locs}_{V'', H}(e_2)\})$$

Instantiating invariant on J_1 ,

$$(1) \quad \text{set}(\text{reach}_{H'}(v))$$

$$(2) \quad \text{disjoint}(\{R, F', \text{reach}_{H'}(v)\})$$

$$(3) \quad \text{stable}(R, H, H')$$

Case 13: E:Drop

$$e = \text{drop}(x; e') \quad (\text{case})$$

$$V', H, R, F \cup g \vdash e' \Downarrow v, H', F'(\mathcal{J}_1) \quad (\text{ad.})$$

$$\Gamma = \Gamma', x : A \quad (\text{case})$$

$$\Sigma; \Gamma' \vdash \frac{q}{q'} e' : B$$

Suppose $\text{dom}(V) = FV(e)$, $\text{no_alias}(V, H)$, $\text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$

By IH, we have invariant on \mathcal{J}_1

NTS (1) - (3) for \mathcal{J}_1

$$(1) \quad \text{dom}(V') = FV(e') \quad (\text{dom}(V) = FV(e) \text{ and def of } FV)$$

$$(2) \quad \text{no_alias}(V', H) \quad (\text{no_alias}(V, H) \text{ and } V' \subseteq V)$$

$$(3) \quad \text{disjoint}(\{R, F \cup g, \text{locs}_{V', H}(e')\})$$

$$g = \text{reach}_H(v') \quad (\text{case})$$

$$g \subseteq \text{locs}_{V, H}(e) \quad (\text{def of } \text{locs}_{V, H})$$

$$R \cap (F \cup g) = \emptyset \quad (\text{disjoint}(\{R, F\}) \text{ and } \text{disjoint}(\{R, \text{locs}_{V, H}(e)\}))$$

$$R \cap \text{locs}_{V', H}(e') = \emptyset \quad (\text{disjoint}(\{R, \text{locs}_{V, H}(e)\}) \text{ and } \text{locs}_{V', H}(e) \subseteq \text{locs}_{V, H}(e))$$

$$F \cap \text{locs}_{V', H}(e') = \emptyset \quad (\text{disjoint}(\{F, \text{locs}_{V, H}(e)\}) \text{ and } \text{locs}_{V', H}(e) \subseteq \text{locs}_{V, H}(e))$$

$$g \cap \text{locs}_{V', H}(e') = \emptyset \quad (\text{no_alias}(V, H))$$

Instantiating invariant on \mathcal{J}_1 ,

$$(1) \quad \text{set}(\text{reach}_{H'}(v))$$

$$(2) \quad \{R, F', \text{reach}_{H'}(v)\}$$

$$(3) \quad \text{stable}(R, H, H')$$

Case 13: E:Share

$$e = \text{shareCopy } x \text{ as } x_1, x_2 \text{ in } e' \quad (\text{case})$$

Suppose $H \models V : \Gamma, \text{dom}(V) = FV(e)$, $\text{no_alias}(V, H)$, $\text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$
(def. of wtf)

$$\text{Let } V_2 = V'[x_1 \text{mapstov}', x_2 \text{mapstov}']$$

We show the subsequent computation is also well-formed to invoke the IH:

$$(1) \quad \text{dom}(V_2) = FV(e') \quad (\text{dom}(V) = FV(e) \text{ and def of } FV)$$

$$(2) \quad \text{no_alias}(V'[x_1 \mapsto v', x_2 \mapsto v''], H) \quad (\text{no_alias}(V, H))$$

$$\text{Let } x' \mapsto v''' \in V'[x_1 \mapsto v']. \text{STS } \text{reach}_{H'}(v''') \cap \text{reach}_{H'}(v'') = \emptyset$$

$$\text{reach}_{H'}(v'') \subseteq L \subseteq F \quad (\text{lemma about copy})$$

$$\text{reach}_{H'}(v''') \subseteq \text{locs}_{V'[x_1 \mapsto v'], H'}(e') \subseteq \text{locs}_{V, H}(e) \quad (\text{stability lemma for copy})$$

$$(3) \text{ disjoint}(\{R, F \setminus L, \text{locs}_{V_2, H'}(e')\})$$

By IH:

$$(2) \{R, F', \text{reach}_{H''}(v)\}$$

$$(3) \text{ stable}(R, H', H'')$$

STS $\text{stable}(R, H, H')$, which follows from $L \cap R = \emptyset$ and stability for

□

TASK 1.8 (SOUNDNESS). *let $H \models V : \Gamma, \Sigma; \Gamma \mid \frac{q}{q'} e : B$, and $V, H \vdash e \Downarrow v, H'$. Then $\forall C \in \mathbb{Q}^+$ and $\forall F, R \subseteq \text{Loc}$, if we have the following (existence lemma):*

- (1) $\text{dom}(V) = FV(e)$
- (2) $\text{no_alias}(V, H)$
- (3) $\text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$, and
- (4) $|F| \geq \Phi_{V, H}(\Gamma) + q + C$

then there exists $F' \subseteq \text{Loc}$ s.t.

- (1) $V, H, R, F \vdash e \Downarrow v, H', F'$
- (2) $|F'| \geq \Phi_{H'}(v : B) + q' + C$

PROOF. Nested induction on the evaluation judgement and the typing judgement.

Case 1: E:Var

$$V, H, R, F \vdash x \Downarrow V(x), H, F \quad (\text{admissibility})$$

$$\Sigma; x : B \mid \frac{q}{q'} x : B \quad (\text{admissibility})$$

$$|F| - |F'| \quad (1)$$

$$= |F| - |F| \quad (\text{ad.})$$

$$= 0 \quad (2)$$

$$\Phi_{V, H}(\Gamma) + q - (\Phi_{H'}(v : B) + q') \quad (3)$$

$$= \Phi_{V, H}(x : B) + q - (\Phi_H(V(x) : B) + q) \quad (\text{ad.})$$

$$= \Phi_H(V(x) : B) + q - (\Phi_H(V(x) : B) + q) \quad (\text{def. of } \Phi_{V, H})$$

$$= 0 \quad (4)$$

$$|F| - |F'| \leq \Phi_{V, H}(\Gamma) + q - (\Phi_{H'}(v : B) + q') \quad ((3), (5))$$

Case 2: E:Const* Due to similarity, we show only for E:ConstI

$$|F| - |F'| = |F| - |F| \quad (\text{ad.})$$

$$= 0$$

$$\Phi_{V, H}(\Gamma) + q - (\Phi_{H'}(v : B) + q') = \Phi_{V, H}(\emptyset) + q - (\Phi_H(v : \text{int}) + q) \quad (\text{ad.})$$

$$= 0 \quad (\text{def of } \Phi_{V, H})$$

$$|F| - |F'| \leq \Phi_{V, H}(\Gamma) + q - (\Phi_{H'}(v : B) + q')$$

Case 4: E:App

Case 5: E:CondT

$$\Gamma = \Gamma', x : \text{bool} \quad (\text{ad.})$$

$$H \models V : \Gamma' \quad (\text{def of W.F.E})$$

$$\Sigma; \Gamma' \mid \frac{q}{q'} e_t : B \quad (\text{ad.})$$

$$V, H, R, F \cup g \vdash e \Downarrow v_2, H_2 \quad (\text{ad.})$$

$$|F \cup g| - |F'| \leq \Phi_{V, H}(\Gamma) + q - (\Phi_{H'}(v : B) + q') \quad (\text{IH})$$

$$|F| - |F'| \leq \Phi_{V, H}(\Gamma) + q - (\Phi_{H'}(v : B) + q')$$

$$(1) \text{ set}(\text{reach}_{H''}(v))$$

Case 6: E:CondF Similar to E:CondT

Case 7: E:Let

$$\text{copy} \vdash e \Downarrow v_2, H_2 \quad (\text{case})$$

$$V, H \vdash e_1 \Downarrow v_1, H_1 \quad (\text{ad.})$$

$$\Sigma; \Gamma_1 \mid \frac{q}{p} e_1 : A \quad (\text{ad.})$$

$$H \models V_1 : \Gamma_1 \quad (\text{def of W.D.E})$$

Let $C \in \mathbb{Q}^+, F, R \subseteq \text{Loc}$ be arb.

Suppose $\text{dom}(V) = FV(e)$, $\text{no_alias}(V, H)$, $\text{disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$, and $|F| \geq$

NTF F' s.t.

$$1. V, H, R, F \vdash e \Downarrow v_2, H_2, F' \text{ and}$$

$$2. |F'| \geq \Phi_{H_2}(v_2 : B) + q' + C$$

$$\text{Let } R' = R \cup \text{locs}_{V, H}(\text{lam}(x : \tau. e_2))$$

$$\text{disjoint}(\{R', F, \text{locs}_{V, H}(e_1)\})$$

(Similar to case in Lemma 1.7)

Instantiate IH with $C = C + \Phi_{V_2, H}(\Gamma_2)$, $F = F$, $R = R'$, we get existence lemma

NTS (1) - (4) to instantiate existence lemma on J_1

$$(1) \text{ dom}(V_1) = FV(e_1)$$

$$(2) \text{ no_alias}(V_1, H)$$

$$(3) \text{ disjoint}(\{R, F, \text{locs}_{V, H}(e)\})$$

((1) - (3) all verbatim as in Lemma 1.7)

$$(4) |F| \geq \Phi_{V_1, H}(\Gamma_1) + q + C + \Phi_{V, H}(\Gamma_2)$$

$$(|F| \geq \Phi_{V, H}(\Gamma) + q + C \text{ and } \Phi_{V, H}(\Gamma) \geq \Phi_{V_1, H}(\Gamma_1) + \Phi_{V, H}(\Gamma_2))$$

Instantiating existence lemma on J_1 , we get F'' s.t.

$$1. V, H, R', F \vdash e_1 \Downarrow v_1, H_1, F'' \text{ and}$$

$$2. |F''| \geq \Phi_{H_1}(v_1 : A) + p + C + \Phi_{V_2, H_1}(\Gamma_2)$$

For the second premise:

$$\Sigma; \Gamma_2, x : A \mid \frac{p}{q} e_2 : B \quad (\text{ad.})$$

$$H_1 \models v_1 : A \text{ and} \quad (\text{Theorem 3.3.4})$$

$$H_1 \models V : \Gamma_2 \quad (???)$$

$$H_1 \models V' : \Gamma_2, x : A \quad (\text{def of } \models)$$

$$V', H_1 \vdash e_2 \Downarrow v_2, H_2 \quad (\text{ad.})$$

$$\text{Let } g = \{l \in H_1 \mid l \notin F_1 \cup R \cup \text{locs}_{V', H_1}(e_2)\}$$

Instantiate IH with $C = C$, $F = F'' \cup g$, $R = R$, we get existence lemma on J_2 :

NTS (1) - (4) to instantiate existence lemma on J_1

$$(1) \text{ dom}(V'_2) = FV(e_2)$$

$$(2) \text{ no_alias}(V'_2, H_1)$$

$$(3) \text{ disjoint}(\{R, F'' \cup g, \text{locs}_{V'_2, H_1}(e_2)\})$$

((1) - (3) all verbatim as in Lemma 1.7)

$$(4) |F'' \cup g| \geq \Phi_{V'_2, H_1}(\Gamma_2, x : (A - 1)) + p + C$$

$$\text{STS } |F'' \cup g| \geq \Phi_{V_2, H_1}(\Gamma_2) + \Phi_{H_1}(v_1 : (A - 1)) + p + C$$

$$|F'' \cup g| \geq \|V_1\|_H + |F| - \|v_1\|_{H_1} \quad (\text{conservation lemma})$$

$\geq \Phi_{V,H}(\Gamma) + q + C + \ V_1\ _H - \ v_1\ _{H_1}$ ($ F \geq \Phi_H(V) + q + C$)	NTS (1) - (4) to instantiate existence lemma on J_1
STS $\Phi_{V_1,H}(\Gamma_1) + q + C \ V_1\ _H - \ v_1\ _{H_1} \geq \Phi_{H_1}(v_1 : (A - 1)) + p$	(1) $\text{dom}(V'_2) = FV(e_2)$
$\Phi_{V_1,H}(\Gamma_1) \geq \Phi_{H_1}(v_1 : (A - 1))$ (lemma about cf typing)	(2) $\text{no_alias}(V'_2, H_1)$
STS $\ V_1\ _H - \ v_1\ _{H_1} + q \geq p$ (done by aux lemma)	(3) $\text{disjoint}(\{R, F'' \cup g, \text{locs}_{V'_2, H_1}(e_2)\})$
Instantiating existence lemma on J_2 , we get $F^{(3)}$ s.t.	((1) - (3) all verbatim as in Lemma 1.7)
1. $V'_2, H_1, R, F'' \cup g \vdash e_2 \Downarrow v_2, H_2, F^{(3)}$	(4) $ F'' \cup g \geq \Phi_{V'_2, H_1}(\Gamma_2, x : A) + q + C$
2. $ F^{(3)} \geq \Phi_{H_2}(v_2 : B) + q' + C$	$ F'' \cup g \geq F'' $
Take $F' = F^{(3)}$	$\geq \Phi_{H_1}(v_1 : A) + p + C + \Phi_{V_2, H}(\Gamma_2)$ (IH)
$V, H, R, F \vdash e \Downarrow v_2, H_2, F'$ and	$= \Phi_{H_1}(v_1 : A) + p + C + \Phi_{V'_2, H_1}(\Gamma_2)$ (Lemma 4.3.3)
$ F' \geq \Phi_{H_2}(v_2 : B) + q' + C$ (from IH)	$= \Phi_{V'_2, H_1}(\Gamma_2, x : A) + p + C$ (def of Φ)
Case 14: E:Let1	Instantiating existence lemma on J_2 , we get $F^{(3)}$ s.t.
$V, H \vdash e \Downarrow v_2, H_2$ (case)	1. $V'_2, H_1, R, F'' \cup g \vdash e_2 \Downarrow v_2, H_2, F^{(3)}$
$V, H \vdash e_1 \Downarrow v_1, H_1$ (ad.)	2. $ F^{(3)} \geq \Phi_{H_2}(v_2 : B) + q' + C$
$\Sigma; \Gamma_1 \mid \frac{q}{p} e_1 : A$ (ad.)	Take $F' = F^{(3)}$
$H \models V_1 : \Gamma_1$ (def of W.D.E)	$V, H, R, F \vdash e \Downarrow v_2, H_2, F'$ and (E:Let)
Let $C \in \mathbb{Q}^+, F, R \subseteq \text{Loc}$ be arb.	$ F' \geq \Phi_{H_2}(v_2 : B) + q' + C$ (from IH)
Suppose $\text{dom}(V) = FV(e)$, $\text{no_alias}(V, H)$, $\text{disjoint}(\{R, F, \text{locs}_{V,H}(e)\})$, and $ F \geq \Phi_V(V)$	Case 8: E:Pair Similar to E:Const*
NTF F' s.t.	Case 9: E:MatP Similar to E:MatCons
1. $V, H, R, F \vdash e \Downarrow v_2, H_2, F'$ and	Case 10: E:Nil Similar to E:Const*
2. $ F' \geq \Phi_{H_2}(v_2 : B) + q' + C$	Case 11: E:Cons
Let $R' = R \cup \text{locs}_{V,H}(\text{lam}(x : \tau.e_2))$	$V, H \vdash \text{cons}(x_1; x_2) \Downarrow l, H'$ (case)
$\text{disjoint}(\{R', F, \text{locs}_{V,H}(e_1)\})$	Let $C \in \mathbb{Q}^+, F, R \subseteq \text{Loc}$ be arb.
(Similar to case in Lemma 1.7)	Suppose $\text{dom}(V) = FV(e)$, $\text{no_alias}(V, H)$, $\text{disjoint}(\{R, F, \text{locs}_{V,H}(e)\})$, $ F \geq \Phi_V(V)$
Instantiate IH with $C = C + \Phi_{V_2, H}(\Gamma_2)$, $F = F$, $R = R'$, we get existence lemma on J_1	NTF F' s.t.
NTS (1) - (4) to instantiate existence lemma on J_1	1. $V, H, R, F \vdash e \Downarrow v, H', F'$ and
(1) $\text{dom}(V_1) = FV(e_1)$	2. $ F' \geq \Phi_{H'}(v : B) + q' + C$
(2) $\text{no_alias}(V_1, H)$	Let $F' = F$
(3) $\text{disjoint}(\{R, F, \text{locs}_{V,H}(e)\})$	Case 12: E:MatNil Similar to E:Cond*
((1) - (3) all verbatim as in Lemma 1.7)	Case 13: E:MatCons
(4) $ F \geq \Phi_{V_1, H}(\Gamma_1) + q + C + \Phi_{V, H}(\Gamma_2)$	$V(x) = (l, \text{alive})$ (ad.)
$(F \geq \Phi_{V, H}(\Gamma) + q + C \text{ and } \Phi_{V, H}(\Gamma) \geq \Phi_{V_1, H}(\Gamma_1) + \Phi_{V, H}(\Gamma_2))$	$H(l) = \langle v_h, v_t \rangle$ (ad.)
Instantiating existence lemma on J_1 , we get F'' s.t.	$\Gamma = \Gamma', x : L^P(A)$ (ad.)
1. $V, H, R', F \vdash e_1 \Downarrow v_1, H_1, F''$ and	$\Sigma; \Gamma', x_h : A, x_t : L^P(A) \mid \frac{q+p+1}{q'} e_2 : B$ (ad.)
2. $ F'' \geq \Phi_{H_1}(v_1 : A) + p + C + \Phi_{V_2, H_1}(\Gamma_2)$	$V'', H \vdash e_2 \Downarrow v, H'$ (ad.)
For the second premise:	Let $C \in \mathbb{Q}^+, F, R \subseteq \text{Loc}$ be arb.
$\Sigma; \Gamma_2, x : A \mid \frac{\max(p, q)}{q'} e_2 : B$ (ad.)	$H \models V(x) : L^P(A)$ (def of W.D.E)
$H_1 \models v_1 : A$ and (Theorem 3.3.4)	$H'' \models v_h : A, H'' \models v_t : L^P(A)$ (ad.)
$H_1 \models V : \Gamma_2$ (???)	$H \models v_h : A, H \models v_t : L^P(A)$ (???)
$H_1 \models V' : \Gamma_2, x : A$ (def of \models)	$H \models V'' : \Gamma', x_h : A, x_t : L^P(A)$ (def of W.D.E)
$V', H_1 \vdash e_2 \Downarrow v_2, H_2$ (ad.)	Suppose $\text{no_alias}(V, H)$, $\text{disjoint}(\{R, F, \text{locs}_{V,H}(e)\})$, and $ F \geq \Phi_{V,H}(\Gamma) + q + C$
Let $g = \{l \in H_1 \mid l \notin F'' \cup R \cup \text{locs}_{V', H_1}(e_2)\}$	NTF F' s.t.
Instantiate IH with $C = C, F = F'' \cup g, R = R$, we get existence lemma on J_1	1. $V, H, R, F \vdash e \Downarrow v, H', F'$ and
	2. $ F' \geq \Phi_{H'}(v : B) + q' + C$

Let $g = \{l \in H \mid l \notin F \cup R \cup \text{locsv}''_{V,H}(e_2)\}$

We want to g nonempty, in particular, that $l \in g$

$$l \notin F \cup R \quad (\text{disjoint}(\{R, F, \text{locsv}_{V,H}(e)\}))$$

$$\text{AFSOC } l \in \text{locsv}''_{V,H}(e_2)$$

Then $l \in \text{reach}_H(\bar{V}''(x'))$ for some $x' \neq x$

$$x' \in \{x_h, x_t\}$$

(since $\text{reach}_H(\bar{V}(x')) \cap \text{reach}_H(\bar{V}(x)) = \emptyset$ from $\text{no_alias}(V, H)$)

WLOG let $x' = x_h$

But then $\mu_{\text{reach}_H(\bar{V}(x))}(l) \geq 2$ and $\text{set}(\text{reach}_H(\bar{V}(x)))$ doesn't hold

$$l \notin \text{locsv}''_{V,H}(e_2)$$

Hence $l \in g$

Next, we have $\text{no_alias}(V'', H)$ and $\text{disjoint}(\{R, F \cup g, \text{locsv}''_{V,H}(e_2)\})$
(similar to case in Lemma 1.2)

By IH with $C' = C, F'' = F \cup g$ and the above conditions, we have: $F^{(3)}$ s.t.

$$1. V'', H, R, F \cup g \vdash e_2 \Downarrow v, H', F^{(3)}$$

$$2. |F^{(3)}| \geq \Phi_{H'}(v : B) + q' + C$$

Where we also verify the precondition that

$$|F''| \geq \Phi_{V'',H}(\Gamma', x_h : A, x_t : L^P(A)) + q + p + 1 + C' :$$

$$\begin{aligned} |F''| &= |F \cup g| \\ &= |F| + |g| \quad (F \text{ and } g \text{ disjoint}) \\ &\geq \Phi_{V,H}(\Gamma) + q + C + |g| \quad (\text{Sp.}) \\ &= \Phi_{V,H}(\Gamma', x_h : A, x_t : L^P(A)) + p + q + C + |g| \\ &\quad (\text{Lemma 4.1.1}) \\ &= \Phi_{V,H}(\Gamma', x_h : A, x_t : L^P(A)) + p + q + C + 1 \\ &\quad (g \text{ nonempty}) \end{aligned}$$

Now take $F' = F^{(3)}$

$$V, H, R, F \vdash e \Downarrow v, H', F' \quad (\text{E:MatCons})$$

$$|F'| \geq \Phi_{H'}(v : B) + q' + C \quad (\text{From the IH})$$

Case 13: E:Share

$$V, H \vdash e \Downarrow v, H'' \quad (\text{case})$$

$$V'[x_1 \mapsto v', x_2 \mapsto v''], H' \vdash e' \Downarrow v, H'' \quad (\text{ad})$$

$$\Sigma; \Gamma, x : A \mid \frac{q}{q'} e : B \quad (\text{case})$$

$$A \curlywedge A_1, A_2, 1 \quad (\text{ad.})$$

$$\Sigma; \Gamma, x_1 : A_1, x_2 : A_2 \mid \frac{q}{q'} e : B \quad (\text{ad.})$$

Let $C \in \mathbb{Q}^+, F, R \subseteq \text{Loc}$ be arb.

Suppose $\text{no_alias}(V, H)$, $\text{disjoint}(\{R, F, \text{locsv}_{V,H}(e)\})$, and $|F| \geq \Phi_{V,H}(\Gamma, x : A) + q + C$

NTF F'' s.t.

$$1. V, H, R, F \vdash e \Downarrow v, H'', F'' \text{ and}$$

$$2. |F''| \geq \Phi_{H''}(v : B) + q' + C$$

We need to show the freelist is sufficient for the subsequent computation to invoke the IH:

Instantiate with $C, F \setminus L$, and R

$$\text{STS } |F \setminus L| \geq \Phi_{V_2,H'}(\Gamma, x_1 : A_1, x_2 : A_2) + q + C$$

$$\iff |F| - |L| \geq \Phi_{V_2,H'}(\Gamma) + \Phi_{V_2,H'}(x_1 : A_1) + \Phi_{V_2,H'}(x_2 : A_2) + q + C$$

$$\iff |F| \geq \Phi_{V_2,H'}(\Gamma) + \Phi_{V_2,H'}(x_1 : A_1) + \Phi_{V_2,H'}(x_2 : A_2) + \|v'\|_H + q + C$$

$$\iff |F| \geq \Phi_{V_2,H'}(\Gamma) + \Phi_{V,H}(x : A) + q + C \quad (\text{definition of sharing relation})$$

$$\iff |F| \geq \Phi_{V,H}(\Gamma, x : A) + q + C \quad (\text{stability of copying})$$

done from assumption

By IH, we get F'' fulfilling the previous two points for the case. \square

9 COPY-FREE GARBAGE COLLECTION SEMANTICS

Consider the GC semantics (from now on copy semantics) above, with the share rule replaced with the following:

$$\frac{V = V'[x \mapsto v'] \quad V'[x_1 \mapsto v', x_2 \mapsto v''], H', R, F \vdash e \Downarrow v, H'', F'}{V, H, R, F \vdash \text{share } x \text{ as } x_1, x_2 \text{ in } e \Downarrow v, H'', F'} \quad (\text{S}_{22})$$

Call this new semantics free semantics for copy-free. It is easy to see that any terminating computation in copy has a corresponding one in free that can be instantiated with an equal or smaller freelist. This is expressed as the following lemma:

LEMMA 1.9. *Let $V, H, R, F \vdash^{\text{copy}} v, H', F'$. Then there exists an F'' s.t. $V, H, R, F \vdash^{\text{free}} v, H', F''$ and $|F''| \geq |F'|$.*