

19/11/14 340 + 1000

$$= p^2 \sum_{y \in \mathbb{N}_0} (1-p)^y \sum_{x \in \{y+1, y+2, \dots\}} (1-p)^x$$

looks "close" to
geometric series

let $i := x - (y+1)$ then $i \in \{1, 2, \dots\} = \mathbb{N}_0$
 $\Rightarrow x = i + y + 1$

$$= p^2 \sum_{y \in \mathbb{N}_0} (1-p)^y \sum_{i \in \mathbb{N}_0} (1-p)^{i+y+1}$$

$$= p^2 \sum_{y \in \mathbb{N}_0} (1-p)^{2y+1} \sum_{i \in \mathbb{N}_0} (1-p)^i$$

$\frac{1}{1-(1-p)} = \frac{1}{p}$

$$= p(1-p) \sum_{y \in \mathbb{N}_0} ((1-p)^2)^y$$

$$\frac{1}{1-(1-p)^2} = \frac{1}{1-1+2p-p^2}$$

$$= \frac{p(1-p)}{p(2-p)} = \frac{1-p}{2-p}$$

$X, Y \sim \text{Geom}(p)$

1


$$P(X > Y) = \sum_{x \in \mathbb{R}} \sum_{y \in \mathbb{R}} P_{X,Y}(x,y) \mathbb{1}_{x>y}$$

$$= \sum_{x \in \mathbb{R}} \sum_{y \in \mathbb{R}} P_X(x) P_Y(y) \mathbb{1}_{x>y}$$

$$= \sum_{x \in \mathbb{R}} \sum_{y \in \mathbb{R}} P_X^{\text{d.f.}}(x) P_Y^{\text{d.f.}}(y) \mathbb{1}_{x>y}$$

$$= \sum_{x \in \mathbb{N}_0} \sum_{y \in \mathbb{N}_0} (1-p)^x p (1-p)^y p \mathbb{1}_{x>y}$$

as $p \rightarrow 0$, $P(X > Y) \rightarrow \frac{1}{2}$



CDF of Geometric (corrected)

B_1, B_2, \dots ^{i.i.d} Bern(p) # of zeroes

1 2 3 ... X X+1

$$S(x) = P(X > x) = P(X \geq x+1) \quad \begin{array}{l} x+1 \text{ zeroes or more} \\ \text{before the first one} \end{array}$$

$$= P(X \geq x+1, X=x+1) + P(X \geq x+1, X=x+2) + P(X \geq x+1, X=x+3) + \dots$$

$$P(B_1=0, B_2=0, \dots, B_{x+1}=0) = (1-p)^{x+1}$$

$$\Rightarrow F(x) = 1 - (1-p)^{x+1}$$


By the book:

$$\text{let } j := i - (x+1) \Rightarrow i = j + x + 1$$

$$\begin{aligned} S(x) &= P(X \geq x+1) = \sum_{i \in \{x+1, x+2, \dots\}} p(1-p)^x \\ &= p \sum_{j \in \mathbb{N}_0} (1-p)^{j+x+1} = p(1-p)^{x+1} \underbrace{\sum_{j \in \mathbb{N}_0} (1-p)^j}_{\frac{1}{1-(1-p)}} \\ &= (1-p)^{x+1} \end{aligned}$$

$$X \sim \text{Geom}(p)$$

Imagine X is time on a clock, maybe every "second". Imagine "seconds" are seconds.
What if instead, we had n ~~seconds~~ per second:



$$\Rightarrow F_{X_n}(x) = 1 - (1-p)^{nx+1}$$

Let X_n denote stopping time $X_n \sim P_{X_n}(x) = (1-p)^{nx} p \mathbb{1}_{x \in \{0, \frac{1}{n}, \frac{2}{n}, \dots, 1+\frac{1}{n}, 1+\frac{2}{n}, \dots\}}$
The process "stops" almost immediately if n is large and p is modest
But what if p was very small? $p \rightarrow 0, n \rightarrow \infty$ s.t. $\lambda = np > 0$

And we substitute...

$$P_{X_n}(x) = \left(1 - \frac{\lambda}{n}\right)^{nx} \frac{\lambda}{n} \mathbb{1}_{x \in \{0, \frac{1}{n}, \frac{2}{n}, \dots\}}$$

Now we let $n \rightarrow \infty$ effectively running infinite Bernoulli's in each second:

$$\lim_{n \rightarrow \infty} P_{X_n}(x) = \lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^{nx} \frac{\lambda}{n} \mathbb{1}_{x \in \{0, \frac{1}{n}, \frac{2}{n}, \dots\}}$$

Recall limit calculus... $\lim_{n \rightarrow \infty} f(n)g(n) = \lim_{n \rightarrow \infty} f(n) \lim_{n \rightarrow \infty} g(n)$

$$\Rightarrow \lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^{nx} \lim_{n \rightarrow \infty} \frac{\lambda}{n} \lim_{n \rightarrow \infty} \mathbb{1}_{x \in \{0, \frac{1}{n}, \frac{2}{n}, \dots\}}$$

$$\lim_{n \rightarrow \infty} f(n) + g(n) = \lim_{n \rightarrow \infty} f(n) + \lim_{n \rightarrow \infty} g(n)$$

$$= \left(\lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^n \right)^x \left(\lim_{n \rightarrow \infty} \frac{1}{n} \right) \lim_{n \rightarrow \infty} \mathbb{1}_{x \in \{1, \frac{2}{n}, \frac{3}{n}, \dots\}}$$

$$= (e^{-\lambda x})(0) \mathbb{1}_{x \in [0, \infty)} \quad \leftarrow \text{continuous support}$$

= 0 Not a pmf!!!

\Rightarrow We're in trouble!

$$|S_{X_{\infty}}| \neq |\mathbb{N}| \quad \text{ctbl } \infty$$

No longer a discrete rv

$$|S_{X_{\infty}}| = |\mathbb{R}|$$

ctbl ∞


Let's take a look at the CDF...

$$F_{X_{\infty}}(x) = \lim_{n \rightarrow \infty} F_{X_n}(x) = \lim_{n \rightarrow \infty} 1 - \left(1 - \frac{\lambda}{n}\right)^{nx+1} = 1 - \left(\lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n \right)^x$$

$\lim_{n \rightarrow \infty} (1 - \lambda/n)$

This term in red becomes 1

$$= 1 - e^{-\lambda x}$$



Monotonically incr.

$$\textcircled{3} \frac{d}{dx} F(x) \geq 0$$

$$\lambda e^{-\lambda x} > 0$$

for all $x \geq 0$

and = 0 for

all $x < 0$

Is this a valid CDF?

1) $F(x) \in [0, 1]$ since it's a prob

If $x < 0 \Rightarrow F(x) = 0$

If $x > 0 \Rightarrow F(x) \in [0, 1]$ since $1 - e^{-\lambda x} \in [0, 1]$

2) $\lim_{x \rightarrow -\infty} F(x) = 0$ since $x < 0 \Rightarrow F(x) = 0$

3) $\lim_{x \rightarrow \infty} F(x) = 1$ $\lim_{x \rightarrow \infty} 1 - e^{-\lambda x} = 1 - \lim_{x \rightarrow \infty} e^{-\lambda x} = 1 - 0 = 1$ ✓

\Rightarrow It's a valid CDF

Def: A cont. rv X has $|S_X| = |\mathbb{R}|$ and no PMF,
 the PMF is $p(x) = 0$ & $P(X=x) = 0 !!$

They have CDF's, The deriv. of the CDF
 (how fast the rv collects prob at any x) is useful:

$f_X(x) := \frac{d}{dx}(F_X(x))$ is called the rv's prob 'density' function (PDF)

Why density? It's the deriv of prob in any given region:

$$a < b, P(X \in (a, b)) = F(b) - F(a) \stackrel{\text{FTC}}{=} \int_a^b f_X(x) dx \Rightarrow \begin{aligned} P(X \in (-\infty, b]) &= F(b) - F(-\infty) \\ &= \int_{-\infty}^b f_X(x) dx \end{aligned}$$

Also, many times the PDF is avail. in closed form but
 the CDF is not! Proposition:

$$= P(X \in (-\infty, \infty)) = \int_{\mathbb{R}} f_X(x) dx \quad \text{the Hairy-Integration Lemma for cont. rv's,}$$


$f_X(x) \geq 0$ since $F_X(x)$ is monotonically Non-dec: f_X can be > 0

$$S_X = \{x: f(x) > 0\}$$

It's not a problem
 the PMF!!!

The Xoo we discussed is a famous brand name rv, the exponential

$$X \sim \text{Exp}(\lambda) := \underbrace{\lambda e^{-\lambda x}}_{\text{pdf}} \mathbb{1}_{x \geq 0} \quad \lambda \in (0, \infty) \text{ since } p \in (0, 1), n \in \mathbb{N}$$




Cont.

Another form of C.R.V. is the uniform r.v. or just "Uniform"

$$X \sim U(a, b) := \frac{1}{b-a} \mathbb{1}_{X \in [a, b]}$$

$$S_X = [a, b], \quad a \in \mathbb{R}, b \in \mathbb{R}, a < b$$



Valid density? $\int_{\mathbb{R}} \frac{1}{b-a} \mathbb{1}_{X \in [a, b]} = \int_a^b \frac{1}{b-a} dx = \frac{1}{b-a} (x)_a^b = \frac{b-a}{b-a} = 1 \checkmark$

$\forall x \ f(x) \geq 0$ \wedge $x \in (-\infty, a) \Rightarrow f(x) = 0$, $x \in [a, b] \Rightarrow f(x) = \frac{1}{b-a} > 0$
 $x \in (b, \infty) \Rightarrow f(x) = 0 \checkmark$

\wedge $a=0, b=1 \Rightarrow X \sim U(0, 1) = \mathbb{1}_{X \in [0, 1]}$ the standard uniform

Very important r.v. especially in comp. science!

if $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$ $f_{\vec{x}}(\vec{x}) = f_{x_1, \dots, x_n}(x_1, \dots, x_n)$ joint density function (JDF) (7)


if x_1, \dots, x_n ind $\Rightarrow f_{x_1}(x_1) f_{x_2}(x_2) \dots f_{x_n}(x_n)$

if $x_1, \dots, x_n \stackrel{iid}{\sim} = f(x_1) f(x_2) \dots f(x_n)$

$\int_{\mathbb{R}^n} f_{\vec{x}}(\vec{x}) d\vec{x} = 1$ 'Happy Dimples'

To get prob's from a JDF, you need to integrate for $n=2$

over a region $P(\vec{x} \in A) = \int_A f_{\vec{x}}(\vec{x}) d\vec{x}$



A calc. fact: Leibnitz's Rule

$$\frac{d}{dt} \int_{a(t)}^{b(t)} g(x, y) dy = g(x, b(t)) b'(t) + g(x, a(t)) a'(t)$$

If $b(t) = t, a(t) = -\infty$

$$\frac{d}{dt} \int_{-\infty}^t g(x, y) dy = g(x, t) \quad \text{since } a(t) \text{ is constant} \Rightarrow a'(t) = 0$$

let X_1, X_2 be two cont. rv's


$$T = X_1 + X_2 \sim f_T(t) = ?$$

Method 1

Note: if $F_T(t)$ is known
 $\Rightarrow f_T(t) = \frac{d}{dt} [F_T(t)]$ What if not known?

We want to derive a general formula like before
 as a function of
 the j.d.f or
 densities (if indep.)

Method 2
 p145 of the textbook



$$F_T(t)$$

$$P(T \leq t)$$

$$P(\vec{x} \in A_t) = \int_A f_{X_1, X_2}(x_1, x_2) dx_1 dx_2 = \int_{x_1 \in \mathbb{R}} \int_{x_2 = -\infty}^{x_2 = t - x_1} f_{X_1, X_2}(x_1, x_2) dx_2 dx_1$$

we need to make this look like Fubini's rule so
 To do so we need to reindex similar to what we
 did with the sums for the geometric problem with $P(X > Y)$
 let $X_2 = V - X_1 \Rightarrow V = X_2 + X_1 \Rightarrow dv = dx_2$

$$V = X_2 + X_1$$

↑↑

$$\text{let } x_2 = V - x_1$$

which is a random variable like we

did with the $P(X > v)$

geometric problem

$$\Rightarrow x_2 = -\infty \Rightarrow v = -\infty$$

$$\Rightarrow x_2 = t - x_1 \Rightarrow v = t$$

$$\frac{dx_2}{dv} = 1 \Rightarrow dv = dx_2$$

drop subscript on $x_1 \Rightarrow x$

$$= \int_{x \in \mathbb{R}} \int_{v=-\infty}^{v=t} f_{x_1, x_2}(x, v-x) dv dx$$

free variable of integration

$$F_T(t) = \int_{-\infty}^t \underbrace{\left(\int_{x \in \mathbb{R}} f_{x_1, x_2}(x, v-x) dx \right)}_{\text{density}} dv$$

Recall

$$P(X \in (-\infty, b]) = F(b) - F(-\infty)$$

$$= \int_{-\infty}^b f_X(x) dx$$

Any integral that looks like this means density is under!

$$F(x) = \int_{-\infty}^x f(v) dv$$

↑ this is the density of the integral

↑ free variable

general conv.
formula