Math 342W / 642 / RM742 Spring 2026
Midterm Examination One

Professor Adam Kapelner

March 18, 2026

Full Name _____

Code of Academic Integrity

Since the college is an academic community, its fundamental purpose is the pursuit of knowledge. Essential to the success of this educational mission is a commitment to the principles of academic integrity. Every member of the college community is responsible for upholding the highest standards of honesty at all times. Students, as members of the community, are also responsible for adhering to the principles and spirit of the following Code of Academic Integrity.

Activities that have the effect or intention of interfering with education, pursuit of knowledge, or fair evaluation of a student's performance are prohibited. Examples of such activities include but are not limited to the following definitions:

Cheating Using or attempting to use unauthorized assistance, material, or study aids in examinations or other academic work or preventing, or attempting to prevent, another from using authorized assistance, material, or study aids. Example: using an unauthorized cheat sheet in a quiz or exam, altering a graded exam and resubmitting it for a better grade, etc.

I acknowledge and agree to uphold this Code of Academic Integrity.

_____ signature

_____ date

Instructions

This exam is 110 minutes (variable time per question) and closed-book. You are allowed **two** pages (front and back) of a "cheat sheet", blank scrap paper (provided by the proctor) and a graphing calculator (which is not your smartphone). Please read the questions carefully. Within each problem, I recommend considering the questions that are easy first and then circling back to evaluate the harder ones. No food is allowed, only drinks.

Problem 1 George Box and Norman Draper in 1987 wrote "All models are wrong but some are useful". Below are some conceptual questions about this aphorism and modeling in general. Let $< \boldsymbol{X}, \boldsymbol{y} >$ be a historical dataset. Let $\boldsymbol{x}_*$ refer to a future object that is not a row in the historical dataset $\boldsymbol{X}$.

(a) [11 pt / 11 pts] For each true statement, circle the roman numeral on the left.

(i) In the quote, "models are wrong" since $\hat{y}_* := g(\boldsymbol{x}_*) \neq y_*$ for all $\boldsymbol{x}_*$.

(ii) In the quote, "models are wrong" since they are approximations to reality.

(iii) In the quote, "models are wrong" since you cannot get an honest assessment of how well they will predict on average for $\boldsymbol{x}_*$.

(iv) In the quote, "models are wrong" but in context they only were speaking about non-mathematical models.

(v) In the quote, "models are wrong" only in the case where $\mathcal{Y} \nsubseteq \mathbb{R}$.

(vi) In the quote, "some [models] are useful" since they are approximations to reality.

(vii) In the quote, "some [models] are useful" only because you can compute oos error metrics.

(viii) In the quote, "some [models] are useful" since g_{final} is more accurate than g if you do a training-test split

(ix) In the quote, "some [models] are useful" but in context they meant only those that are mathematical.

(x) In the quote, "some [models] are useful" because they beat g_0.

(xi) In the quote, "some [models] are useful" because they give you the ability to compute $\hat{y}_* := g(\boldsymbol{x}_*)$.

Problem 2 Let $< \boldsymbol{X}, \boldsymbol{y} >$ be a historical dataset with $n = 300$. Assume the first column of $\boldsymbol{X}$ is $\mathbf{1}_n$ so that there are $p + 1$ columns. Let the sets A, B, C denote random subsets of the indicies such that $A \cup B \cup C = \{1, \ldots, n\}$ but none of the sets have any overlap, i.e., $A \cap B = A \cap C = B \cap C = \varnothing$ and each set is equally sized. Let $\text{nrow}(U)$, $\text{ncol}(U)$ denote the number of rows and columns in U respectively. Let $U = \boldsymbol{QR}$ be the Q-R decompositions of the the matrix on the left hand side. Let the subscript of X and $\boldsymbol{y}$ denote only those rows, e.g., $\boldsymbol{X}_A$ is the submatrix $\boldsymbol{X}$ with only the rows of A. Also let $\boldsymbol{Q}_S$, $\boldsymbol{R}_S$ denote the Q-R decomposition of the matrix $\boldsymbol{X}_S$ where S is a subset of $\{1, \ldots, n\}$. Assume $\boldsymbol{X}_A, \boldsymbol{X}_B, \boldsymbol{X}_C$ are full rank.

(a) [18 pt / 29 pts] For each true statement, circle the roman numeral on the left.

(i) $\text{nrow}(\boldsymbol{X}_A) + \text{nrow}(\boldsymbol{X}_B) + \text{nrow}(\boldsymbol{X}_C) = n$

(ii) $\text{ncol}(\boldsymbol{X}_A) + \text{ncol}(\boldsymbol{X}_B) + \text{ncol}(\boldsymbol{X}_C) = p + 1$

(iii) $\text{colsp}[\boldsymbol{X}_A] = \text{colsp}[\boldsymbol{X}_B] = \text{colsp}[\boldsymbol{X}_C]$

(iv) $\text{rank}[\boldsymbol{X}_A] = \text{rank}[\boldsymbol{X}_B] = \text{rank}[\boldsymbol{X}_C]$

(v) $\text{nrow}(\boldsymbol{Q}_A) + \text{nrow}(\boldsymbol{Q}_B) + \text{nrow}(\boldsymbol{Q}_C) = n$

(vi) $\text{ncol}(\boldsymbol{Q}_A) + \text{ncol}(\boldsymbol{Q}_B) + \text{ncol}(\boldsymbol{Q}_C) = p + 1$

(vii) $\text{colsp}[\boldsymbol{Q}_A] = \text{colsp}[\boldsymbol{Q}_B] = \text{colsp}[\boldsymbol{Q}_C]$

(viii) $\text{rank}[\boldsymbol{Q}_A] = \text{rank}[\boldsymbol{Q}_B] = \text{rank}[\boldsymbol{Q}_C]$

(ix) $\text{rank}[\boldsymbol{R}_A] = \text{rank}[\boldsymbol{R}_B] = \text{rank}[\boldsymbol{R}_C]$

(x) The matrix $\boldsymbol{QR}$ is full rank

(xi) The matrix $\boldsymbol{XR}$ is full rank

(xii) $\boldsymbol{Q}_A^\top \boldsymbol{Q}_A = \boldsymbol{I}_{p+1}$

(xiii) $\boldsymbol{Q}_A^\top \boldsymbol{Q}_B = \boldsymbol{I}_{p+1}$

(xiv) $\boldsymbol{Q}_A^\top \boldsymbol{Q}_C = \boldsymbol{I}_{p+1}$

(xv) $\boldsymbol{Q}_A \boldsymbol{Q}_A^\top = \boldsymbol{I}_{n/3}$

(xvi) $\boldsymbol{Q}_A \boldsymbol{Q}_B^\top = \boldsymbol{I}_{n/3}$

(xvii) $\boldsymbol{Q}_A \boldsymbol{Q}_C^\top = \boldsymbol{I}_{n/3}$

(xviii) $\boldsymbol{Q}_A \boldsymbol{Q}_A^\top \boldsymbol{y}_A = \boldsymbol{Q}_B \boldsymbol{Q}_B^\top \boldsymbol{y}_B = \boldsymbol{Q}_C \boldsymbol{Q}_C^\top \boldsymbol{y}_C$

Let $\mathcal{A}$ be an algorithm and $\mathcal{H}$ be a candidate set. Let $g = \mathcal{A}(< \boldsymbol{X}, \boldsymbol{y} >, \mathcal{H})$ and $g_S = \mathcal{A}(< \boldsymbol{X}_S, \boldsymbol{y}_S >, \mathcal{H})$ where S is a subset of $\{1, \ldots, n\}$. The word "expected" below means on average over many possible datasets.

(b) [3 pt / 32 pts] Which of the following models have the lowest in-sample error? Circle the roman numeral on the left. If they are all equal, or the answer cannot be determined given the information available, do not circle any.

(i) g_A

(ii) $g_{A \cup B}$

(iii) g

(c) [3 pt / 35 pts] Which of the following models have the lowest **expected** in-sample error? Circle the roman numeral on the left. If they are all equal, or the answer cannot be determined given the information available, do not circle any.

(i) g_A

(ii) $g_{A \cup B}$

(iii) g

(d) [3 pt / 38 pts] Which of the following models have the lowest **expected** oos error? Circle the roman numeral on the left. If they are all equal, or the answer cannot be determined given the information available, do not circle any.

(i) g_A

(ii) $g_{A \cup B}$

(iii) g

(e) [3 pt / 41 pts] Which of the following models have the lowest **expected** estimation error? Circle the roman numeral on the left. If they are all equal, or the answer cannot be determined given the information available, do not circle any.

(i) g_A

(ii) $g_{A \cup B}$

(iii) g

(f) [3 pt / 44 pts] Which of the following models have the lowest **expected** misspecification error? Circle the roman numeral on the left. If they are all equal, or the answer cannot be determined given the information available, do not circle any.

(i) g_A

(ii) $g_{A \cup B}$

(iii) g

(g) [3 pt / 47 pts] Which of the following models have the lowest **expected** ignorance error? Circle the roman numeral on the left. If they are all equal, or the answer cannot be determined given the information available, do not circle any.

(i) g_A

(ii) $g_{A \cup B}$

(iii) g

(h) [3 pt / 50 pts] Compare the expected error of $g_A(\boldsymbol{X}_A)$ and $g_A(\boldsymbol{X}_B)$. In the circle below, write $<, >$ or $=$.

$$\text{Error}(g_A(\boldsymbol{X}_A)) \quad \bigcirc \quad \text{Error}(g_A(\boldsymbol{X}_B))$$

(i) [13 pt / 63 pts] Circle the roman numeral on the left for all the error measurements that are *honest* i.e., they represent the generalization error in future predictions without further assumptions.

(i) $\text{Error}(g_A(\boldsymbol{X}_A))$

(ii) $\text{Error}(g_A(\boldsymbol{X}_B))$

(iii) $\text{Error}(g_A(\boldsymbol{X}_C))$

(iv) $\text{Error}(g_A(\boldsymbol{X}_{B \cup C}))$

(v) $\text{Error}(g_A(\boldsymbol{X}))$

(vi) $\text{Error}(g_{A\cup B}(\boldsymbol{X}_A))$

(vii) $\text{Error}(g_{A\cup B}(\boldsymbol{X}_B))$

(viii) $\text{Error}(g_{A\cup B}(\boldsymbol{X}_C))$

(ix) $\text{Error}(g_{A\cup B}(\boldsymbol{X}_{B\cup C}))$

(x) $\text{Error}(g_{A\cup B}(\boldsymbol{X}))$

(xi) $\text{Error}(g(\boldsymbol{X}_A))$

(xii) $\text{Error}(g(\boldsymbol{X}_B))$

(xiii) $\text{Error}(g(\boldsymbol{X}_C))$

Assume SSE is the error metric going forward.

(j) [3 pt / 66 pts] Write a formula to find the in-sample error of the model g_A. Be as explicit as possible using the notation given in this problem.

(k) [3 pt / 69 pts] Write a formula to find the *most accurate* measure of the oos error of the model g_A. Be as explicit as possible using the notation given in this problem.

(l) [3 pt / 72 pts] Now assume n is very large, i.e. $n >> 300$. Write a formula to find the *most accurate* measure of the oos error of the model g_A. Be as explicit as possible using the notation given in this problem.

(m) [3 pt / 75 pts] Write a formula to find an estimate of the oos error of the model g using $K = 3$-fold cross validation. Be as explicit as possible using the notation given in this problem.

(n) [3 pt / 78 pts] Will the estimate in the previous question be conservative? Yes/no and why.

Problem 3

Consider the following data frame `D` illustrated using the `skimr::skim` function:

```
-- Data Summary ------------------------
                           Values
Name                       D
Number of rows             137
Number of columns          8
_______________________
Column type frequency:
  factor                   2
  numeric                  6
________________________
Group variables            None

-- Variable type: factor ----------------------------------------------
  skim_variable n_missing complete_rate ordered n_unique top_counts
1 trt                   0             1 FALSE          2 con: 69, exp: 68
2 celltype              0             1 FALSE          4 sma: 48, squ: 35, ade: 27, lar: 27

-- Variable type: numeric ---------------------------------------------
  skim_variable n_missing complete_rate    mean      sd p0 p25 p50 p75 p100
1 time                  0             1 122.    158.     1  25  80 144  999
2 status                0             1   0.934   0.249  0   1   1   1    1
3 karno                 0             1  58.6    20.0   10  40  60  75   99
4 diagtime              0             1   8.77   10.6    1   3   5  11   87
5 age                   0             1  58.3    10.5   34  51  62  66   81
6 prior                 0             1   2.92    4.56   0   0   0  10   10
```

Let $y :=$ `time`. And let $\boldsymbol{X}$ denote the design matrix constructed from the remaining columns of `D`. Assume the first column of $\boldsymbol{X}$ is $\mathbf{1}_n$.

(a) [2 pt / 80 pts] What type of modeling task is this? In this class, the answer is a one-word answer.

(b) [2 pt / 82 pts] What is the data type of `celltype`?

(c) [3 pt / 85 pts] What is the *most likely* data type of `status`? Be as specific as possible.

(d) [4 pt / 89 pts] What is the value of p?

(e) [4 pt / 93 pts] What is g_0?

(f) [5 pt / 98 pts] How do we know g_0 will perform poorly in-sample for this dataset?

(g) [3 pt / 101 pts] Compute R^2 for g_0.

(h) [4 pt / 105 pts] Without any other information given, do you think OLS will beat g_0 oos? Yes, no or can't be determined. Explain.

(i) [3 pt / 108 pts] We run an OLS on all features and calculate SSE = 2482093. Compute RMSE to once decimal.

(j) [5 pt / 113 pts] We run an OLS on all features and calculate SSE = 2482093. Compute R^2 to three decimals.

(k) [8 pt / 121 pts] Assume there was no overfitting. What is the *least likely* source of error in the OLS model?

Here are the first six rows:

```
> head(D)
             trt  celltype time status karno diagtime age prior
107 experimental smallcell   29      1    40        8  67     0
121 experimental     adeno  186      1    90        3  60     0
51       control     adeno   12      1    50        4  63    10
58       control     large  553      1    70        2  47     0
15       control  squamous   11      1    70       11  48    10
81  experimental  squamous   25      1    20       36  63     0
```

(l) [6 pt / 127 pts] Provide the first six rows of $\boldsymbol{X}$. No need to name the columns or rows.

(m) [8 pt / 135 pts] Consider the scenario where you add junk vectors (full of random values) one-by-one and run OLS for each new junk vector. You keep adding junk vectors until OLS can no longer compute. Assume each iteratively new $\boldsymbol{X}$ is full rank. Graph the expected R^2 from an OLS on the y axis and the number of vectors *in addition* to the features already in $\boldsymbol{X}$ on the x-axis. The y-axis should start at zero and end at 1. Label axes.

Now assume $y :=$ `trt`

(n) [2 pt / 137 pts] What type of modeling task is this? In this class, the answer is a one- or two-word answer.

(o) [4 pt / 141 pts] Explain two reasons why the perceptron will likely not be a good model here.

Now assume you use KNN to model the response.

(p) [4 pt / 145 pts] Provide an explicit distance function $d(\boldsymbol{x}_*, \boldsymbol{x}_i)$ where $\boldsymbol{x}_i$ is a row in $\boldsymbol{X}$ and $\boldsymbol{x}_*$ is a future subject we wish to predict their response for. Be as explicit as possible in your notation

Regardless of what you wrote in the previous question, assume the distance function is reasonable and standard going forward.

(q) [3 pt / 148 pts] Explain why KNN will likely beat the SVM with the Vapnik objective function.

(r) [4 pt / 152 pts] If $K = n$, what is the prediction for the third row in `D`?

(s) [4 pt / 156 pts] If $K = 1$, what is the prediction for the third row in `D`?