

recap...

 $X \sim \text{Bernoulli}(p) := p^x (1-p)^{1-x}$
 $X \sim \text{Binomial}(n, p) := \binom{n}{x} p^x (1-p)^{n-x}$
 $X \sim \text{Hypergeometric}(n, K, N) := \frac{\binom{K}{x} \binom{N-K}{n-x}}{\binom{N}{n}}$
 $X \sim \text{Hypergeometric}(n, p, N) := \frac{\binom{pN}{x} \binom{1-pN}{n-x}}{\binom{N}{n}}$
 $\lim_{N \rightarrow \infty} X \sim \text{Hypergeometric}(n, p, N) = X \sim \text{Binomial}(n, p)$ conceptually...

Binomial distribution Proof

property of PMF: $\sum_{x \in \text{Supp}[X]} p(x) = 1$ if the Binomial is truly a rv.

verify $\sum_{x=0}^n \binom{n}{x} p^x (1-p)^{n-x} = 1$ we must prove = 1 how?

recall the Binomial Theorem $(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$ let $a=p, b=1-p$

reason why we call this distribution 'binomial'

plug in $(p+(1-p))^n = \sum_{x=0}^n \binom{n}{x} p^x (1-p)^{n-x}$

$(p-p+1)^n = (1)^n = 1$ QED

what are "iid" r.v.'s?

X_1 & X_2 are independent r.v. if:

$$\begin{aligned} P(X_1=x_1 | X_2=x_2) &= P(X_1=x_1) \\ P(X_2=x_2 | X_1=x_1) &= P(X_2=x_2) \\ P(X_1=x_1, X_2=x_2) &= P(X_1=x_1) P(X_2=x_2) \end{aligned}$$

remember test for independence?

$$P(A \cap B) = P(A) P(B)$$


Multiplication Rule

$$P(A \cap B) = P(A|B) \cdot P(B)$$

$\forall x_1 \in \text{Supp}[X_1] \quad \forall x_2 \in \text{Supp}[X_2]$

X_1 & X_2 are independent & identically distributed - iid
if X_1 and X_2 are independent & $X_1 \stackrel{\text{iid}}{\sim} X_2$ denoted $X_1, X_2 \stackrel{\text{iid}}{\sim}$

(statistics) patients in a medical trial are considered iid



are X_1, X_2 independent? - Yes, since

$P(X_1=1 | X_2=1) = P(R_1|R_2) = P(R_1) = P(X_1=1)$

$P(X_1=1 | X_2=0) = P(R_1|G_2) = P(R_1) = P(X_1=1)$

$P(X_1=0 | X_2=1) = P(G_1|R_2) = P(G_1) = P(X_1=0)$

$P(X_1=0 | X_2=0) = P(G_1|G_2) = P(G_1) = P(X_1=0)$

thus X_1, X_2 are $\stackrel{\text{iid}}{\sim}$ independent \rightarrow iid (independent & identically distributed)

$X_1, X_2 \stackrel{\text{iid}}{\sim} \text{Bernoulli}(1/3)$

g1, X2 function of two r.v.

$T_2 = X_1 + X_2$ $\text{Supp}[T_2] = \{0, 1, 2\}$

disjoint events can be added

two sets are called 'disjoint' if their intersection is the empty set: \emptyset

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

one is two is

1000 coin flips - 600 Heads

$P(600H) = P(X=600) = \binom{1000}{600} (1/2)^{1000}$

$X \sim \text{Binomial}(1000, 1/2)$ this is how it all connects

a bag so big that, if you

take one ball out, the

change in the ratio of

success is negligible and

observed as fixed

OR

a small bag but you are

sampling with replacement

thus keeping the ratio of

success fixed

a really big bag with a fixed proportion of successes

$T = \lim_{n \rightarrow \infty} X \sim \text{Hypergeometric}(n, p, N)$

$T = X_1, X_2, \dots, X_n$ where $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Bernoulli}(p)$

you're summing a whole bunch of iid Bernoullis and at the end you ask - how many successes did I get?

$\text{Binomial}(n, p) = X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Bernoulli}(p)$

$X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Bernoulli}(p)$

$T_n = X_1 + X_2 + \dots + X_n$ $\text{Supp}[T_n] = \{0, 1, 2, \dots, n\}$

$T_n = \sum_{i=1}^n X_i$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

$T_n \sim$

CDF of Binomial

what is a CDF of a Binomial?

$F(x) = P(X \leq x)$

$F(x) = \sum_{i=0}^x p(i)$

incomplete regularized gamma function

$F(x) = \sum_{i=0}^x \binom{n}{i} p^i (1-p)^{n-i}$

$= I_{1-p}(n-k, 1+k) := (n-k) \binom{n}{k} \int_0^{1-p} t^{n-k-1} (1-t)^k dt$

no closed form! (if you find one you get a Fields Medal)

Geometric r.v. model

we keep trying and as soon as we succeed we stop

$X_1, X_2, \dots \stackrel{\text{iid}}{\sim} \text{Bernoulli}(p)$

infinite series of binary experiments all with probability p of success where all are independent of one another (iid)

infinite sequence of iid r.v.'s

keep on trying until $X_n=1$ (success)

self-note: think of the y-coordinates as indices and x-coordinates as T_0 and T_n .

6-failure 1-success

let $T = \min\{t: X_t=1\}$

first success 'stopping time'

$\{3, 4, 7, 11, 18, \dots\}$

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success

1st success