

Long Acting Injectable Guide

Medication	FDA Approved Indications	Starting Dose	Maintenance Dose	Frequency and Injection Site	Oral Dose Equivalence	Oral Overlap?	Late Doses?	Early Doses?
Abilify Maintena ® Aripiprazole	Schizophrenia Bipolar I	400 mg	400 mg 300 mg - for drug interactions	Every 4 weeks Deltoid or gluteal	10-15 mg per day	Yes - 2 weeks after initial dose	Next dose given ASAP <u>Oral overlap needed after 7 days</u>	No sooner than <u>26 days</u> between
Aristada ® Aristada Initio ® Aripiprazole	Schizophrenia	441 mg 662 mg 882 mg 1064 mg 675 mg (initio)	441 mg 662 mg 882 mg 1064 mg N/A	Every 4 weeks Every 4 weeks Every 4 or 6 weeks Every 8 weeks Once Deltoid - 441, 675 mg Gluteal - all others	10 mg 15 mg 15-20 mg 15 mg	No if patient gets <u>initio + 30 mg tablet</u> Otherwise, oral overlap for 3 weeks after first injection	ASAP <u>Oral overlap/starting dose</u> needed if it has been <u>2-4 weeks</u> since due date	No sooner than <u>14 days</u> between
Zyprexa Relprevv ™ Olanzapine	Schizophrenia	<u>First 8 weeks</u> 210 mg/2 weeks 405 mg/4 weeks 300 mg/2 weeks 405 mg/4 weeks	<u>After 8 weeks</u> 150 mg/2 weeks 300 mg/4 weeks 210 mg/2 weeks 300 mg/2 weeks 405 mg/4 weeks	Every 2-4 weeks Gluteal only	10 mg 15 mg 20 mg 15 mg	No	Not specified	Not specified
Invega Sustenna ® Paliperidone Invega Trinza ® Paliperidone	Schizophrenia Schizoaffective	Day 1: 234 mg Day 8: 156 mg Can switch if patient has been stable on Sustenna for at least 4 months	39 mg 78 mg 117 mg 156 mg 234 mg 273 mg 410 mg 546 mg 819 mg	Every 4 weeks Every 4 weeks Every 4 weeks Every 4 weeks Every 4 weeks Every 12 weeks Every 12 weeks Every 12 weeks Every 12 weeks Starting must be	3 mg (1 mg risperidone) 3 mg (2 mg risperidone) 6 mg (3 mg risperidone) 9 mg (4 mg risperidone) 12 mg (6 mg risperidone) 78 mg Sustenna dose 117 mg Sustenna dose 156 mg Sustenna dose 234 mg Sustenna dose	No	Up to 6 weeks after last injection before catch-up dosing is required 14 days late	7 days early 14 days early

Long Acting Injectable Guide

				deltoid, maintenance can be deltoid or gluteal				
Risperdal Consta ® Risperidone	Schizophrenia Bipolar I	25 mg 37.5 mg 50 mg	25 mg 37.5 mg 50 mg	Every 2 weeks Deltoid or gluteal	2 mg 3 mg 4 mg	Yes - 3 weeks after 1st injection	Not specified	Not specified
Perseris ® Risperidone	Schizophrenia	90 mg 120 mg	90 mg 120 mg	Every 4 weeks Subcutaneous (abd)	3 mg 4 mg	No	ASAP	Not specified
Haldol Decanoate ® Haloperidol	Schizophrenia Tourettes	10 to 20 times daily oral dose - If more than 100 mg, must be in 2 separate injections (100 mg initially, the rest in 3-7 days)	10-15 times previous daily oral dose	Every 3-4 weeks	With overlap: keep oral dose until at least the second injection Without overlap: 20 times oral dose for initial injection		<u>Less than 6 weeks</u> since last injection - <u>resume dose ASAP</u> <u>6-12 weeks</u> since last injection - <u>consider oral overlap</u> if symptoms reemerge <u>Over 13 weeks</u> - needs <u>restart</u>	Not specified
Prolixin Decanoate ® Fluphenazine	Psychotic disorders	1.25 times oral dose	12.5 mg - 50 mg	Every 2 weeks	12.5 mg = 10 mg/day oral 25 mg = 20 mg/day oral 37.5 mg = 30 mg/day oral 50 mg = 40 mg/day oral ^based on every 3 week dosing	After the first decanoate dose, decrease oral fluphenazine dose by half after the initial injection - consider discontinuing oral therapy after second injection	Steady state not reached (<u>less than 4 weeks of therapy</u>) or between <u>6-24 weeks since last dose</u> : administer <u>missed dose ASAP</u> , provide PO antipsychotic supplementation if symptoms reemerge, monitor closely for ADRs Steady state reached (<u>4-6 weeks of therapy</u>) and <u>less than 6 weeks</u> since last injection: <u>administer dose ASAP</u> <u>Over 24 weeks</u> since last injection: stabilize on PO antipsychotic, reinitiate starting dose	Not specified