Foundations of Mathematics and the Foundational Crisis

Kevin Kappelmann

Technical University of Munich

May 22, 2017

Overview

- 1 Causes of the crisis
 - Ancient Mathematics
 - Uncertainties
- 2 The Foundational Crisis
 - Logicism
 - Intuitionism
 - Formalism
 - Peak and End
- 3 Aftermath and Prospects

Ancient Mathematics

First Steps

■ In the beginning, maths served as an applied tool.

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC
 - Axiomatic, deductive treatment of mathematics similar to modern systems

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC
 - Axiomatic, deductive treatment of mathematics similar to modern systems
 - A work of timeless certainty...

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC
 - Axiomatic, deductive treatment of mathematics similar to modern systems
 - A work of timeless certainty...... at least so had been thought

Ancient Mathematics

Euclid's Postulates

"Let the following be postulated:

1 To draw a straight line from any point to any point.

- 1 To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.

- To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.
- 3 To describe a circle with any centre and distance [radius].

- 1 To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.
- 3 To describe a circle with any centre and distance [radius].
- That all right angles are equal to one another.

- To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.
- 3 To describe a circle with any centre and distance [radius].
- 4 That all right angles are equal to one another.
- That, if a straight line falling on two straight lines make the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which are the angles less than the two right angles."

Gauß proved the independence of Euclid's fifth postulate in 1813.

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - \Rightarrow Scepticism towards used systems

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - ⇒ Scepticism towards used systems
- Rigorous axiomatisation of mathematical branches in the late 19th century

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - ⇒ Scepticism towards used systems
- Rigorous axiomatisation of mathematical branches in the late 19th century
 - Arithmetic of natural numbers by Peano
 - Geometry by Hilbert and Pasch
 - Predicate logic by Frege

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - ⇒ Scepticism towards used systems
- Rigorous axiomatisation of mathematical branches in the late 19th century
 - Arithmetic of natural numbers by Peano
 - Geometry by Hilbert and Pasch
 - Predicate logic by Frege
- Desire for a universal and consistent system

Cantor's set theory

- Cantor's set theory
 - "A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought which are called elements of the set."
 Georg Cantor

- Cantor's set theory
 - "A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought which are called elements of the set."
 - Certainly universal, but informal and thus not adequate for a study of consistency

- Cantor's set theory
 - "A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought which are called elements of the set."
 - Certainly universal, but informal and thus not adequate for a study of consistency
 - Nonetheless, it was widely accepted.

- Cantor's set theory
 - "A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought which are called elements of the set."
 - Certainly universal, but informal and thus not adequate for a study of consistency
 - Nonetheless, it was widely accepted.
- Frege tried to build a consistent foundation with logic.
 - Sophisticated work, but not well received

- Cantor's set theory
 - "A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought which are called elements of the set."
 - Certainly universal, but informal and thus not adequate for a study of consistency
 - Nonetheless, it was widely accepted.
- Frege tried to build a consistent foundation with logic.
 - Sophisticated work, but not well received
- And then came Russell...

Consider the set of all sets that are not members of themselves:

$$R := \{X \mid X \notin X\}$$

Consider the set of all sets that are not members of themselves:

$$R := \{X \mid X \notin X\}$$

Question: Is R a member of itself? That is, does $R \in R$ hold?

Consider the set of all sets that are not members of themselves:

$$R := \{X \mid X \notin X\}$$

Question: Is R a member of itself? That is, does $R \in R$ hold?

Answer: $R \in R \iff R \notin R$, a contradiction!

Cantor's as well as Frege's system were victims of this paradox.

- Cantor's as well as Frege's system were victims of this paradox.
- "Hardly anything more unfortunate can befall a scientific writer than to have one of the foundations of his edifice shaken after the work is finished."

- Cantor's as well as Frege's system were victims of this paradox.
- "Hardly anything more unfortunate can befall a scientific writer than to have one of the foundations of his edifice shaken after the work is finished."
- Systems founded on Cantor's set theory were on shaky ground.

- Cantor's as well as Frege's system were victims of this paradox.
- "Hardly anything more unfortunate can befall a scientific writer than to have one of the foundations of his edifice shaken after the work is finished."
- Systems founded on Cantor's set theory were on shaky ground.
- A new foundation of mathematics had to be found.

The Three Schools of Thought

- Three schools of thought tried to establish a new foundation.
 - Logicism
 - Intuitionism
 - Formalism



Source: geopolicraticus.tumblr.com/post/142561195372

A Foundation Made of Logic

Russell and Whitehead revisited the idea of reducing mathematics to logic, as tried by Frege.

A Foundation Made of Logic

- Russell and Whitehead revisited the idea of reducing mathematics to logic, as tried by Frege.
- Mathematics as an extension of logic

A Foundation Made of Logic

- Russell and Whitehead revisited the idea of reducing mathematics to logic, as tried by Frege.
- Mathematics as an extension of logic
- Only few axioms that must pose fundamental logical principles

A Foundation Made of Logic

- Russell and Whitehead revisited the idea of reducing mathematics to logic, as tried by Frege.
- Mathematics as an extension of logic
- Only few axioms that must pose fundamental logical principles
- Chief work: Principia Mathematica

■ Type theory to avoid antinomies

- Type theory to avoid antinomies
- Difficulties in explaining some axioms
 - Axiom of reducibility
 - Axiom of infinity

- Type theory to avoid antinomies
- Difficulties in explaining some axioms
 - Axiom of reducibility
 - Axiom of infinity
- Regarded as "the outstanding example of an unreadable masterpiece."

- Type theory to avoid antinomies
- Difficulties in explaining some axioms
 - Axiom of reducibility
 - Axiom of infinity
- Regarded as "the outstanding example of an unreadable masterpiece."
- Nonetheless, many adopted it as a new foundation.

```
*54·43. \vdash :: \alpha, \beta \in 1 \cdot D : \alpha \cap \beta = \Lambda \cdot \equiv : \alpha \cup \beta \in 2

Dem.

\vdash .*54·26 \cdot D \vdash :: \alpha = \iota^{\iota}x \cdot \beta = \iota^{\iota}y \cdot D : \alpha \cup \beta \in 2 \cdot \equiv : x \neq y \cdot \begin{bmatrix} *51·231 \end{bmatrix} \qquad \equiv : \iota^{\iota}x \cap \iota^{\iota}y = \Lambda \cdot \begin{bmatrix} *13·12 \end{bmatrix} \qquad \equiv : \alpha \cap \beta = \Lambda \qquad (1)
\vdash .(1) .*11·11·35 . D
\vdash :: (\exists x, y) . \alpha = \iota^{\iota}x . \beta = \iota^{\iota}y . D : \alpha \cup \beta \in 2 . \equiv : \alpha \cap \beta = \Lambda \qquad (2)
\vdash .(2) .*11·54 .*52·1 . D \vdash . Prop
```

From this proposition it will follow, when arithmetical addition has been defined, that 1 + 1 = 2.

Principia Mathematica's infamous proof of 1+1=2. It was not until page 379 that this proof was possible.

■ Mathematics is **not** reducible to some formal system.

- Mathematics is **not** reducible to some formal system.
- Mathematics is a constructive process conducted by humans.

- Mathematics is **not** reducible to some formal system.
- Mathematics is a constructive process conducted by humans.
- The existence of any mathematical object ist equivalent to the possibility of its construction, according to Brouwer.

- Mathematics is **not** reducible to some formal system.
- Mathematics is a constructive process conducted by humans.
- The existence of any mathematical object ist equivalent to the possibility of its construction, according to Brouwer.
 - ⇒ No antinomies as paradoxical sets cannot be constructed.

- Mathematics is **not** reducible to some formal system.
- Mathematics is a constructive process conducted by humans.
- The existence of any mathematical object ist equivalent to the possibility of its construction, according to Brouwer.
 - ⇒ No antinomies as paradoxical sets cannot be constructed.
- Consequently, some assumptions of classical logic must be rejected.

$$P \vee \neg P \equiv ?$$

$$P \vee \neg P \equiv ?$$

Proposition: There exist two irrational numbers a and b such that a^b is rational.

$$P \vee \neg P \equiv ?$$

Proposition: There exist two irrational numbers a and b such that a^b is rational.

Proof. It is known that $\sqrt{2}$ is irrational. Let us consider the number $\sqrt{2}^{\sqrt{2}}$.

$$P \vee \neg P = ?$$

Proposition: There exist two irrational numbers a and b such that a^b is rational.

Proof. It is known that $\sqrt{2}$ is irrational. Let us consider the number $\sqrt{2}^{\sqrt{2}}$. If it is rational, our statement is proved.

$$P \vee \neg P \equiv ?$$

Proposition: There exist two irrational numbers a and b such that a^b is rational.

Proof. It is known that $\sqrt{2}$ is irrational. Let us consider the number $\sqrt{2}^{\sqrt{2}}$. If it is rational, our statement is proved. If it is irrational, $(\sqrt{2}^{\sqrt{2}})^{\sqrt{2}} = 2$ proves our statement.

The Price to Pay

Intuitionistic logic aggravates many proofs.

The Price to Pay

- Intuitionistic logic aggravates many proofs.
- "Taking this tertium non datur (law of excluded middle) from the mathematician would be the same as, say, denying the astronomer his telescope and the boxer the use of his fists."
 - David Hilbert

The Price to Pay

- Intuitionistic logic aggravates many proofs.
- "Taking this tertium non datur (law of excluded middle) from the mathematician would be the same as, say, denying the astronomer his telescope and the boxer the use of his fists."
 - David Hilbert
- Consequently, only a few scholars adhered to intuitionism.

Mathematics as a Symbol Modifier

Formalists support the autonomy of mathematics.

Mathematics as a Symbol Modifier

- Formalists support the autonomy of mathematics.
- Mathematics shall be based on symbols and axioms that describe syntactical operations on them.

Mathematics as a Symbol Modifier

- Formalists support the autonomy of mathematics.
- Mathematics shall be based on symbols and axioms that describe syntactical operations on them.
- Mathematics must not justify the existence of its objects as its objects are just meaningless shapes.

- Formalists support the autonomy of mathematics.
- Mathematics shall be based on symbols and axioms that describe syntactical operations on them.
- Mathematics must not justify the existence of its objects as its objects are just meaningless shapes.
- Hilbert, as one of the most renowned mathematicians of given time, was the driving force of formalism.

• Hilbert initiated a research programme called *Hilbert's* program.

- Hilbert initiated a research programme called Hilbert's program.
 - 1 Formalise system that is able to derive all of mathematics using syntactical operations.

- Hilbert initiated a research programme called Hilbert's program.
 - 1 Formalise system that is able to derive all of mathematics using syntactical operations.
 - 2 Prove the system's consistency with metamathematical reasoning.

• Consequently, only a few scholars adhered to intuitionism.

The End



Georg Cantor.

Beiträge zur Begründung der transfiniten Mengenlehre.

Mathematischen Annalen, 46:481, 1895.



Philip J. Davis, Reuben Hersh, and Elena Anne Marchisotto.

The Mathematical Experience.

Modern Birkäuser Classics, 2012.



Gottlob Frege.

Grundgesetze der Arithmetik, volume 2.

1902.



David Hilbert.

Die Grundlagen der Mathematik.

Abhandlungen aus dem Mathematischen Seminar der Hamburger Universität, page 80, 1928.