

Foundations of Mathematics and the Foundational Crisis

Kevin Kappelmann

Technical University of Munich

May 20, 2017

- Mathematics strives for absolute certainty.

- Mathematics strives for absolute certainty.
- For this, we use rigorous proofs.

- Mathematics strives for absolute certainty.
- For this, we use rigorous proofs.
- When shall I accept a proof?

- Mathematics strives for absolute certainty.
- For this, we use rigorous proofs.
- When shall I accept a proof?
- Can we trust each other?

- Mathematics strives for absolute certainty.
- For this, we use rigorous proofs.
- When shall I accept a proof?
- Can we trust each other?
- Can we trust a computer?

- Mathematics strives for absolute certainty.
- For this, we use rigorous proofs.
- When shall I accept a proof?
- Can we trust each other?
- Can we trust a computer?
 - Computer-generated proofs are barely verifiable by hand.

- Mathematics strives for absolute certainty.
 - For this, we use rigorous proofs.
 - When shall I accept a proof?
 - Can we trust each other?
 - Can we trust a computer?
 - Computer-generated proofs are barely verifiable by hand.
- ⇒ Controversies

- Mathematics strives for absolute certainty.
- For this, we use rigorous proofs.
- When shall I accept a proof?
- Can we trust each other?
- Can we trust a computer?
 - Computer-generated proofs are barely verifiable by hand.
 - ⇒ Controversies
- But it is not the first controversy in mathematical history...

Overview

- 1 Causes of the crisis
- 2 The Foundational Crisis
 - Logicism
 - Formalism
 - Intuitionism
 - Peak and End
- 3 Aftermath and Prospects

History

- In the beginning, maths served as an applied tool.

History

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.

History

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.

History

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics

History

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC

History

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC
 - Axiomatic, deductive treatment of mathematics similar to modern systems

History

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC
 - Axiomatic, deductive treatment of mathematics similar to modern systems
 - A work of timeless certainty...

History

- In the beginning, maths served as an applied tool.
- Ancient Greeks developed the notion of a proof in 600 BC.
- The Pythagoreans discovered irrational numbers.
 - ⇒ First controversy of mathematics
- Euclid's Elements around 300 BC
 - Axiomatic, deductive treatment of mathematics similar to modern systems
 - A work of timeless certainty...
 - ...at least so had been thought

Euclid's Postulates

“Let the following be postulated:

Euclid's Postulates

“Let the following be postulated:

- 1 To draw a straight line from any point to any point.

Euclid's Postulates

“Let the following be postulated:

- 1 To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.

Euclid's Postulates

“Let the following be postulated:

- 1 To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.
- 3 To describe a circle with any centre and distance [radius].

Euclid's Postulates

“Let the following be postulated:

- 1 To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.
- 3 To describe a circle with any centre and distance [radius].
- 4 That all right angles are equal to one another.

Euclid's Postulates

“Let the following be postulated:

- 1 To draw a straight line from any point to any point.
- 2 To produce [extend] a finite straight line continuously in a straight line.
- 3 To describe a circle with any centre and distance [radius].
- 4 That all right angles are equal to one another.
- 5 That, if a straight line falling on two straight lines make the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which are the angles less than the two right angles.”

Search for foundations

- Gauß proved the independence of Euclid's fifth postulate in 1813.

Search for foundations

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - ⇒ Scepticism towards used systems

Search for foundations

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - ⇒ Scepticism towards used systems
- Rigorous axiomatisation of mathematical branches in the late 19th century

Search for foundations

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - ⇒ Scepticism towards used systems
- Rigorous axiomatisation of mathematical branches in the late 19th century
 - Arithmetic of natural numbers by Peano
 - Geometry by Hilbert and Pasch
 - Predicate logic by Frege

Search for foundations

- Gauß proved the independence of Euclid's fifth postulate in 1813.
 - ⇒ Scepticism towards used systems
- Rigorous axiomatisation of mathematical branches in the late 19th century
 - Arithmetic of natural numbers by Peano
 - Geometry by Hilbert and Pasch
 - Predicate logic by Frege
- Desire for universal and consistent system

Universal systems

- Cantor's set theory

Universal systems

- Cantor's set theory
 - *"A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought — which are called elements of the set."*[1] — Georg Cantor

Universal systems

- Cantor's set theory
 - *"A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought — which are called elements of the set."*[1] — Georg Cantor
 - Certainly universal, but informal and thus not adequate for a study of consistency

Universal systems

- Cantor's set theory

- *"A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought — which are called elements of the set."*[1] — Georg Cantor
- Certainly universal, but informal and thus not adequate for a study of consistency
- Nonetheless, it was widely accepted.

Universal systems

- Cantor's set theory
 - *"A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought — which are called elements of the set."*[1] — Georg Cantor
 - Certainly universal, but informal and thus not adequate for a study of consistency
 - Nonetheless, it was widely accepted.
- Frege tried to build a consistent foundation with logic.
 - Sophisticated work, but not well received

Universal systems

- Cantor's set theory
 - *"A set is a gathering together into a whole of definite, distinct objects of our perception or of our thought — which are called elements of the set."*[1] — Georg Cantor
 - Certainly universal, but informal and thus not adequate for a study of consistency
 - Nonetheless, it was widely accepted.
- Frege tried to build a consistent foundation with logic.
 - Sophisticated work, but not well received
- And then came Russell...

Russell's antinomy

Consider *the set of all sets that are not members of themselves*:

$$R := \{X \mid X \notin X\}$$

Russell's antinomy

Consider *the set of all sets that are not members of themselves*:

$$R := \{X \mid X \notin X\}$$

Question: Is R a member of itself? That is, does $R \in R$ hold?

Russell's antinomy

Consider *the set of all sets that are not members of themselves*:

$$R := \{X \mid X \notin X\}$$

Question: Is R a member of itself? That is, does $R \in R$ hold?

Answer: $R \in R \iff R \notin R$, a contradiction!

The begin of the crisis

- Cantor's as well as Frege's system were victims of this paradox.

The begin of the crisis

- Cantor's as well as Frege's system were victims of this paradox.
- *"Hardly anything more unfortunate can befall a scientific writer than to have one of the foundations of his edifice shaken after the work is finished."*[2] — Gottlob Frege

The begin of the crisis

- Cantor's as well as Frege's system were victims of this paradox.
- *"Hardly anything more unfortunate can befall a scientific writer than to have one of the foundations of his edifice shaken after the work is finished."* [2] — Gottlob Frege
- Systems founded on Cantor's set theory were on shaky ground



The End



Georg Cantor.

Beiträge zur Begründung der transfiniten Mengenlehre.

Mathematischen Annalen, 46:481, 1895.



Gottlob Frege.

Grundgesetze der Arithmetik, volume 2.

1902.