
Boolean expressions.

1. *Express an English sentence as a Boolean expression.*

In each of the following sentences underline up to three atomic propositions and write them as compound propositions using logical connectives.

- (a) For walks in a forrest to be safe, it is necessary, but not sufficient that there are are no poisonous mushrooms along your path and no active ticks.
- (b) There are no active ticks and walks in a forrest are safe, but there are poisonous mushrooms along your path.
- (c) Walks in a forrest are not safe iff there are no active ticks.
- (d) There are either active ticks or poisonous mushrooms along your path or both, whenever the walks in the forrest are not safe.

2. *Build a truth table for a Boolean Expression.*

$$((p \oplus q) \wedge (p \oplus \neg q)) \oplus (\neg q \oplus \neg r).$$

3. *Build a DNF or a CNF for a given truth table.*

Fill in the truth table for the logical expression $p \rightarrow (\neg q \wedge r)$. Create either a DNF or a CNF for this truth table (which one – is up to you).

4. *Simplify a Boolean Expression using identities; prove, disprove tautologies.*

Simplify the following Boolean expressions:

- (a) $(p \oplus q) \oplus (p \oplus r)$
- (b) $(p \wedge q) \wedge (p \rightarrow q \rightarrow r)$

5. *Shade regions in a Venn diagram corresponding to a Boolean Expression.*

Draw a Venn diagram (the ovals correspond to the regions where the corresponding Boolean variables are true). Shade the region, where the compound Boolean expression is true.

- (a) $(p \rightarrow \neg r) \vee (q \rightarrow \neg r)$
- (b) $(p \vee q) \oplus r$
- (c) $(p \wedge q) \vee (\neg q \wedge r)$

Quantifiers.

1. *Express math and programming concepts with predicates.*

Write a logical expression for a predicate $P(x, y)$ which has value true iff the point (x, y) is inside a unit circle (with radius 1 and center $(0, 0)$).

Note 1. If (x, y) is exactly on the border of the unit circle, $P(x, y)$ should evaluate to false.

Note 2. Express similar predicates for points belonging to the 1st (or 2nd, 3rd, 4th) quadrant, or belonging to some rectangle, etc.

2. *Express an English sentence as a predicate logic expression.*

Introduce 1 or 2-argument predicates and express the following sentences in the predicate logic

- (a) Some student in this class has visited Mexico, but has not visited Argentina.
- (b) All students in this class have learned at least one programming language.
- (c) There is a student in this class who has taken every course offered by one of the departments of this university.
- (d) Some student in this class is a citizen of the same country as exactly one other student in this class.

3. *Restore parentheses, identify free/bound variables in a predicate expression.*

For every variable occurrence determine its scope (and what quantifier bounds it, if any such quantifier exists). Draw an arrow from a variable to its quantifier.

$$\forall x \in S (\exists x \in S R(x) \vee P(x)) \wedge Q(x).$$

Does this expression depend on some parameter (and if so, underline that parameter).

4. *Write the negation for a predicate expression; simplify using De Morgan's laws.*

Let U be a set universe and $P(x, y, z)$ be some predicate with three variables $x, y, z \in U$. Rewrite the statement so that negations are applied only to the predicates directly (no negation is outside a quantifier or applied to an expression containing logical connectives).

$$\neg \forall x \in U (\exists y \in U \forall z \in U P(x, y, z) \wedge \exists z \in U \forall y \in U P(x, y, z)).$$

5. *Read and write set-builder notation.*

Use the set-builder notation to describe the set S of all those positive integers k that can be expressed as the product of two positive integers a, b in just one way: Neither a , nor b equals 1 or k (the order of multiplication does not matter; i.e. product $a \cdot b$ and $b \cdot a$ counts as the same way).

Functions and Relations

1. *Given a function, prove/disprove that it is injective, surjective or bijective.*

A function f switches the places of two digits in a positive integer. For example:

$$f(123) = 213 \text{ and } f(123456) = 123546.$$

(In general the 2nd and the 3rd digits from the right switch their places; no other changes are done. For numbers $n < 100$, $f(n) = n$ as there is no 3rd digit from the right in this case.)

Find if f is injective, surjective and/or bijective function.

2. *Given function definitions, evaluate their compositions and inverses.*

Find the inverse function for f defined in the previous problem. Find the composition $f \circ f \circ f$ for this function.

3. *Given a sequence, identify its properties, is it (eventually) constant/periodic, etc.*

Let $a_1 = 7$ and every next member a_{n+1} is the last digit of $3a_n + 2$. Is the sequence (a_n) constant? Eventually constant? Periodic? Eventually periodic?

4. *Convert a description of a binary relation into another form.*

Write a matrix representation for a binary relation $R \subseteq \{0, 1, 2, 3, 4, 5\}^2$ such that $(a, b) \in R$ iff $|a + b|$ or $|a - b|$ is divisible by 3.

5. *Given a binary relation determine if it is a (injective, surjective, bijective) function.*
 Let $S = \{1, 2, 3, 4, 5, 6\}$. Let $R \subseteq S \times S$ be a relation such that $(a, b) \in R$ iff $|a - b| = 3$.
 - (a) Prove that the relation R is a function relation.
 - (b) Is this function injective, surjective and/or bijective?
6. *Given a binary relation determine if it is reflexive, symmetric, antisymmetric or transitive.*
 Let \mathbf{Z} be the set of all integers, and $R \subseteq \mathbf{Z}^2$ is the relation such that $(a, b) \in R$ iff $|a - b|$ is divisible by 7. Is the relation R a reflexive, symmetric, antisymmetric, transitive and/or equivalence relation?
7. *Compute compositions and powers for relations, find transitive closures.*
 The set S consists of 8×8 squares; it is the regular chess board. Two squares $a, b \in S$ are in relation B (written as $(a, b) \in B$) iff it is possible to go from a to b in a single step using a bishop. (Bishops are going diagonally – they can cross as many squares as they like as long as they stay on the same diagonal.) Let B^t be the transitive closure of the relation B . How many squares are in relation B^t with $c \in S$ (where c is the square located on the bottom-left corner of the chess board).

Proofs.

1. *Prove an implication directly.*
 Prove that for any integer x that has decimal notation ending with digit 7, the cube x^3 has the decimal notation that always ends with the same digit (no matter which x you pick).
2. *Prove an implication by contradiction.*
 Assume that $\alpha + \beta$ is a rational number and $\alpha^2 + \beta^2$ is an irrational number. Prove that $\alpha \cdot \beta$ is an irrational number.
3. *Prove a logical equivalence (if and only if).*
 Prove that an integer number n belongs to the interval $[10000, 999999]$ iff $\lfloor \sqrt{n} \rfloor$ is an integer with exactly 3 digits in its decimal notation.
4. *Prove by counterexample.*
 Prove that there exists a prime number p such that the congruence equation

$$x^2 \equiv 2 \pmod{p}$$

has an integer solution x .

5. *Prove by mathematical induction.*
 Prove the following identity for every positive integer n :

$$1 \cdot 1! + 2 \cdot 2! + 3 \cdot 3! + \dots + n \cdot n! = (n + 1)! - 1.$$

6. *Prove or disprove equality of two numbers or sets.*
 Prove that a connected graph with n vertices is a tree iff it has $n - 1$ edges.