

Plant-based DIETARY SUPPLEMENTS (Botanicals/herbs)

Ilona Benkő M.D., Ph.D.

University of Debrecen, Faculty of Medicine,
Dept. of Pharmacology and Pharmacotherapy



Approximately 40 % of the recently used drugs are originated from plants or other living systems,

e.g. many anticancer drugs, aspirine, digitalis, antibiotics, opioids, atropine, physostigmine e.t.c.

After isolation of active ingredients many of them are used as a synthetic drug today.

Screening for the new active ingredients, potential drug molecules in the nature is important even today.

Advantages of the purified chemicals and drugs:

- 1. the reproducible quality and**
- 2. controlled dose with a dose-depending controlled effects**



Quality and strengths of plant extracts and herbal dietary supplements are very variable !!

Standard plant extracts in European Pharmacopoeia have better quality and the content of only the main active ingredient much less variable.

Factors which influence the amount of the active ingredients:

Culturing conditions, soil, sunshine, water

Seasonal or daily rhythms of the plant's life cycle

Which part of the plant was used

Storage conditions

Technology

Pollutants e.g. agricultural chemicals

Delivery conditions

Many ingredients of the herbal supplements saturate biotransformation in the liver !



Drug interactions !!


Factors which influence the amount of the active ingredients:

Culturing conditions, soil, sunshine, water
Seasonal or daily rhythms of the plant's life cycle
Which part of the plant was used
Storage conditions
Technology
Pollutants e.g. agricultural chemicals
Delivery conditions

Many ingredients of the plant extracts saturate active transport molecules in absorption, distribution and excretion and enzymes in biotransformation in the liver !



Hepatotoxicity



Drug interactions !!

DIETARY SUPPLEMENTS with PLANT ORIGIN

Plant extracts have long been used to stay healthy or treat illnesses in any continents. Traditional Chinese Medicine has early written documents which make it well-known all over the world, but plant extracts have been used in any continents and regions.

The Food and Drug Administration (FDA) classifies herbals as dietary supplements, not drugs (2004).

Dietary supplements are made up of processed or unprocessed plant parts (bark, leaves, flowers, fruits and stems), extracts and essential oils. They may come in the form of tablets and capsules or teas and powders. Herbals may be sold as single substances or combined with other materials, such as vitamins and minerals.

The FDA does not regulate dietary supplements as it does the registered drugs:

1. Herbal products do not have to be pure
2. Labelling of such products is under the purview of the FDA
3. Safety does not have to be demonstrated

This means that the quality and strength of herbals can vary greatly among brands

The major problem with a lack of federal oversight on plant-based dietary supplements is the public perception of herbs. Pharmaceuticals, especially registered drugs, tend to be perceived as potent chemicals. What is more, because they are naturally occurring, the chemicals found in herbs may be perceived by the public as less harmful than a drug produced by chemists in labs.

Everybody should be able to appreciate that

these substances are marketed without review of efficacy and safety by authorities.

By law, dietary supplements can be sold legally as long as they do not make any health claims or say they cure a condition.

In many cases the medical value of these substances has not been demonstrated in controlled clinical studies.

Remember, just because it is “natural” does not mean it is gentler or more beneficial !

Many dietary supplements contain active ingredients that have strong biological effects and their safety is not assured.

Although herbal supplement are not synthetic, herbs contain powerful chemical compounds as well. **Monkshood**, for example, can stop the heart and it is very toxic to CNS, killing a human within a couple of hours of even handling it.

It can be mistaken with horse-radish.

Monkshood



Aconitum variegatum

aconitine is a very toxic compound

Poison used for arrows in Asia

Horseradish



Toxic amanitine- containing mushrooms:

Amanita phalloides



Amanita muscaria contains muscarine

Some herbal medicines were evaluated for their therapeutical and toxic, side effects and they become registered drugs in some countries:

Sylimarine an extract of milk thistle (*Silybum marianum*)

is used in hepatic disorders even in hepatic failure or in poisoning with *Amanita* mushrooms

active ingredients: flavonoids

silybin (also known as silybinin or silibinin),
silychristin (silichristin)
and silydianin (silidianin).

Milk Thistle



Side effects:

laxative because of increased bile flow

Pharmacological effects of sylimarine:

in vitro: reduces lipid peroxidation

free radical scavenger

enhances superoxide dismutase activity

inhibits lipoxygenase and formation leukotrienes

increases RNA polymerase I activity , protein synthesis and regeneration

in animal models:

It protects against liver injury caused by CCl_4 , alcohol or acetaminophene (=paracetamol) and prevent fibrosis

In clinical trials:

It has favorable effect on liver injury caused by poisoning with Amanita mushrooms. It was useful in viral hepatitis.

Echinacea extract from *Echinacea purpurea*

or *E. pallida* and *E. angustifolia*



most popular herbal dietary supplement with the highest selling in USA.

active ingredients:

echinacoside, cichoric acid, alkamides, water soluble polysacharides


water soluble extract has effect on cytokine release

THERAPEUTICAL EFFECTS:

1. immunomodulatory effect by release of cytokines: TNFalpha, interleukines, IFNgamma
2. increased phagocytosis, total circulating monocytes, neutrophils, and natural killer cells
3. immunosuppressive through cannabinoid receptors of immune cells
4. inhibition on lipoxigenase, COX enzymes, hyaluronidase - antiinflammatory effects
5. Virucidal activity against influenza and herpes simplex viruses and bactericidal activity against *Streptococcus pyogenes*, *Haemophilus influenzae*, and *Legionella pneumophila*

CLINICAL INDICATIONS: antiviral effects


especially in the early phase of infections

Name	Popular Use (not proven in many cases)	Adverse Reactions
<p>Echinacea / purple coneflower</p> 	<p>To prevent and remedy colds and flu</p> <p>Mode of action:</p> <p>Increases cytokine deliberation (TNFalpha, ILs)</p> <p>Anti-inflammatory effect by inhibition of lipoxigenase and COX enzymes</p>	<p>Upset stomach and diarrhea; skin rash if used topically. Using regularly may suppress the immune system.</p> <p><u>Taking echinacea for <u>eight weeks or more</u> increases risk of hepatotoxicity.</u></p> <p>Flu-like syndrome after i.v. administration</p> <p>Immunosuppression in long treatments</p> <p>Contraindicated in immune disorders !!</p>

Echinacea extract on CYP enzymes

Echinacea alkamide compounds inhibit several CYP enzymes.

Effect of Echinacea extract:



Hydrastis canadensis

Hydrastis canadensis = goldenseal (tabletta, extract)



active ingredients: hydrastin, berberin

antimicrobial effects, choleric

Toxicity: hepatotoxicity in higher doses
carcinogenic in rats,
hepatocellular adenoma, hepatoblastoma

National Toxicology Program, USA, NTP TR 562, NIH Publication No.
10-5903 Planta Med 2012; 78: 1490–1514



Echinacea + *Hydrastis canadensis* combination on CYP enzymes

In USA Echinacea extract often contains *Hydrastis canadensis*

Hydrastis canadensis has izokinolin alkaloids: **hydrastin and berberin with strong inhibitory effect on CYP enzymes.**

Potentiated inhibitory
effects of the two extract



on CYP3A4
+ CYP2D6



**The most of the
drugs are
metabolized by these
2 enzymes**

eg. beta hydroxylation of testosteron is reduced by 88 %

14 -day -long administration increases significantly AUC of midazolam and cyclosporinA .

INCREASED HEPATOTOXICITY (both extract has hepatotoxicity)

ginseng

Panax ginseng

P. quinquefolium, the American variety

Siberian ginseng
(*Eleutherococcus senticosus*)

and Brazilian ginseng (*Pfaffia paniculata*)

Popular Use (not proven in many cases)

Adverse Reactions



Nootropic, to improve mental and physical performance ?

Ginsenosids
+
Panaxosides

Steroid glikozides

Modulatory effect on :
Ca, Na, K ion channels.
GABA-A, 5HT-3, nicotin, Ach,
NMDA receptors

Weak oestrogenic effect
Insomnia, nervousness, hypertension

CYP3A4 interactions

Interaction with:

warfarin
Increased bleeding, increased INR
The same risk with Thr aggregatory inhibitors



With sympathomimetic drugs -
Hypertension, sleepness

Contraindication in Mania –depressive disease



Beneficial effects

Adverse effects

<p>Garlic / Allium sativum</p> 	<p>To prevent stroke, atherosclerosis, high blood pressure and lower cholesterol</p> <p><u>Allicin</u> <u>Organosulfur compounds</u> (S-allylcysteine)</p> <p>inhibit HMG-CoA reductase, Improve vessel elasticity, Antiplatelete and fibrinolytic effect, relaxes smooth muscle, Antimicrobial effect</p>	<p>Upset stomach, intestinal problems, heartburn, bad breath and body odor.</p> <p>nausea (6%), hypotension (1.3%), allergy (1.1%), and bleeding (rare).</p>
<p>St. John's Wort / Hypericum perforatum Blooming is in June at St. John's feast</p> 	<p>To treat mild depression</p> <p><u>hypericin,</u></p> <p><u>Hyperforin has inhibitory effect on reuptake of biogenic amines</u></p>	<p>Sensitivity to sunlight if taken for long spells, itching, headache, fatigue, bloating, weight gain and constipation. Hypomania, mania, autonomic arousal</p> <p>Cautions! With other antidepressants may result in serotonin syndrome (life-threatening sympathetic outflow)</p>


Importance of interactions of St John's Wort extracts

Hypericum perforatum is the most frequently used plant extract because of antidepressive effect of its active ingredient, **hyperforin**.

OTC product !!

15 % of plant dietary supplements to be selling for therapy of depression, especially **St John's Wort extracts** are used .

Drug- dietary supplement interactions



78 % are the interactions induced by
St John's Wort extracts in the cases
published in scientific literature !!

Pharmacodynamic interactions in CNS

St. John's Wort / *Hypericum perforatum*

Warnings for patients:

With any antidepressant drugs

St. John's Wort extracts are NOT RECOMMENDED!!

As plant extract the concentrations of active ingredient are changing in wide range !!

Ingredients of Herbs brought in as dietary supplements are not controlled by authorities  high amplitudes in amounts

Unpredictable amounts and effects
és hatás miatt



Unpredictable serious
toxicity in CNS/agitation, convulsion
or
malignant serotonin sy !

Pharmacodynamic interactions in CNS

St. John's Wort / *Hypericum perforatum*

Mode of action: hyperforin inhibits serotonin reuptake mechanism



It increases the effects of selective serotonin reuptake inhibitors
Toxicity – serotonin sy !!

Serotonin sy



amfetamin caused disco fever ?

Common leading symptoms:


- malignant hyperthermia
- Malignant hypertension
- Muscle rigidity



Life-threatening !
DEATH WITHOUT TREATMENT

Pregnan X receptor - nuclear interactions

The strongest induction of CYP3A4 is through Pregnane X receptor (PXR = steroid XR nuclear receptor)



St. John's Wort extract results in inefficiency of many drugs e.g. anticoncipients, 5-HT_{1D} agonists in migraine attacks, AIDS therapy, warfarin

Life-threatening interactions with St John's Wort extract + warfarin

St John's Wort extract may lay off effect of anticoagulant therapy

reason: enhanced elimination of warfarin

due to induction of CYP2C9 enzyme which metabolizes warfarin



trombosis, AMI, stroke

St John's Wort extract induced CYP enzymes: **CYP3A4, 2C9, 2C19,1A2**



These all together influence metabolism, elimination and bioavailability
of the **80 % of drugs!!**



**It would be desirable to warn patients for the risk of the use on labels of
packages of St. John's Wort extract !!**

Ephedra sinica

Ephedra sinica: about 0,5 m in height, Northeast-China

Active ingredient:

ephedrin and pseudoephedrin alkaloids.

Mode of action of Ephedrin :

Indirect + direct mixed type sympatomimetics

Effects in CNS+ perifery.

Indirect : neurotransmitter release from Adrenerg neurons –
amfetamin-like effect



CNS sympathetic overflow sy – stimulatory amfetamin-like effect - euphoria

Direct effect:

	alfa1	beta1	beta2
adrenalin	+++	+++++	+++
efedrin	+	+++	++



Ephedra distachya L.



Ephedra sinical Ma Huang

ephedra is one of the most infamous dietary supplement

use: body building, sportmen, obesity,
for weight reducing, enhancement of performance

effects: weight reducing effect within 1-12 months
synergistic effect in combination with caffeine

Acute sympathomimetic effects in CNS+periferally

Reduced sensation of fatigue, enhance breath volume + cardiac output

Adrenergic effects , e.g. enhanced blood sugar  excess energy



Enhancement of performance

Ephedra sinica withdrawal

Ephedrin therapeutic use:

as a drug, the synthetic ephedrin is the component of
Mixture pectoralis FoNo

Bronchodilatation in obstructive bronchitis, asthma bronchiale

Ephedrin-containing dietary supplements: are PROHIBITED to sell

2004. U . S . Food and Drug Administration (FDA) banned dietary supplements, herbs containing *ephedra sinica*

2005. in Utahban tried to delibarate, but several months later it was banned again.

Many DEATH of young men due to the uncontrolled use of *ephedra sinica*

***Ephedra sinica* toxic interactions**

Ephedrin-containing dietary supplements became forbidden products because of unsafety

It was impossible to control their use



hypertension, tachycardia, sympathetic overflow in CNS , arrhythmias, myocardial infarction, stroke even in healthy young men!




Interaction:

With any drugs and energy drinks which may result in
sympathetic stimulation or overflow in CNS

Synergistic effect with caffeine

Even physical activity enhances the risk!!

Biloba- prophylaxis of dementia ?

Name	Popular Use (not proven in many cases)	Adverse Reactions
Ginkgo Biloba  Ginkgo leaves	<p>To treat headache, tinnitus, prophylactic agent to prevent progression to dementia</p> <p><u>Flavone glycosides and terpenoids</u> including ginkgolides A, B, C, J, and bilobalide</p> <p>Increased blood flow : PAF antagonist Inhibits erythrocyte and thr aggregation Decrease capillary permeability and edema Antioxidant</p>	<p>Muscle spasms, anxiety, allergic skin reactions, cramps, and mild digestive problems.</p> <p>Bleeding ! (overdose)</p> <p>Or in interactions with other thr. aggregation inhibitors as aspirine</p>







Remarkable *Ginkgo biloba* in [Tournai](#), ([Belgium](#)).

Name	Popular Use (not proven in many cases)	Adverse Reactions
Feverfew / <i>Tanacetum parthenium</i>	<p>To treat headache, migraine, rheumatoid arthritis</p> <p><u>Parthenolide</u> and other antiinflammatory agents</p>	<p>Mouth ulcers, gastrointestinal upset, contact dermatitis</p> <p>Rebound: nervousness, tension-type headache, insomnia</p>



Name	Popular Use (not proven in many cases)	Adverse Reactions
<p>Aloe vera</p> <p>There are more than 300 species of the aloe plant, but <i>Aloe barbadensis</i> (aloe vera) is the best known. grows naturally in dry, tropical climates in Africa, Asia, Europe, and the southern and western parts of the United States and also known as “burn plant,” “lily of the desert,” and “elephant’s gall” .</p>	<p>Popular herbal remedy that was used to help treat wounds, hair loss, hemorrhoids, and digestive issues or prevent sunburn. Heartburn (GER) relief and lowering blood glucose are also hoped.</p> <p>From leaves bright green gel or juice, capsules</p> <p>The yellow pulp that’s found just under the outer part of the plant leaf. Aloe latex has been shown to have laxative properties, and it’s usually taken orally to treat constipation. Drinks, juice, aloe vera water contain aloe latex. Aloin, which is an anthraquinone is the active ingredient.</p>	<p>Aloe latex, however, can be dangerous. Taking aloe latex orally can lead to cramps and bloody diarrhea, haemorrhoids. Renal toxicity: Taking even just 1 gram orally per day for several days could end up causing kidney damage and may even be fatal</p> <p>! Pregnants are very sensitive to aloe vera !! – haemorrhages, even abortus may develop.</p> <p>whole-leaf aloe vera extract created cancerous tumors within the large intestines of rats</p>



Name	Popular Use (not proven in many cases)	Adverse Reactions
<p>Datura stramonium jimsonweed or devil's snare Dose dependent toxicity of atropine</p>	<p>Hiosciamine and atropin especially in seeds</p> <p>Scopolamine especially in leaves</p>	<p>Atropin intoxication or scopolamine intoxication. OPPOSITE EFFECTS on CNS ! Atropin is sympathomimetics both centrally and peripherally. Excitation, involuntary movements, epileptiform convulsions, hallucinations, coma</p> <p>Scopolamine is sedatohypnotics. Somnolence, coma, respiratory distress</p>
<p>Atropa belladonna</p> <p>HIGH Atropin containing plant ! Only the standard belladonna leaf extracts are good to use as medicine</p>  	<p>It was used in neuralgia, pain killer in rheumatic arthritis.</p> <p>Smoke or cigarette against asthma bronchiale.</p> 	

Name	Popular Use (not proven in many cases)	Adverse Reactions
<p>Viscum album L. commonly known as European mistletoe, common mistletoe or simply as mistletoe. It is native to Europe and Western and Southern Asia.</p>	<p>Active ingredient: Viscotoxin - vasodilation</p> <p>Cold dipping of leaves to obtain tea was used. Mistletoe leaves and young twigs are used by herbalists.</p>	<p>Very dangerous- contraindication in children</p> <p>The toxic lectin viscumin has been isolated from <i>Viscum album</i>. Viscumin is a cytotoxic protein (ribosome inactivating protein, or RIP) that binds to galactose residues of cell surface glycoproteins and may be internalised by endocytosis Viscumin strongly inhibits protein synthesis by inactivating the 60 S ribosomal subunit. The structure of this protein is very similar to other RIPs, showing the most resemblance to ricin and abrin.</p>



Table 63–2. Potential toxicity of selected botanicals.

Name	Intended Use	Potential Toxicity
Aconite (monkshood, wolfsbane)	Analgesic (topical and oral)	Cardiac and CNS toxicity with oral use
Borage (beebread, burrage)	Anti-inflammatory, diuretic	Oral use causes gastrointestinal distress and possible <u>hepatotoxicity</u>
Chaparral (creosote bush, greasewood)	Anti-infective, antioxidant	<u>Hepatotoxicity</u> and renal dysfunction
Coltsfoot (coughwort)	Respiratory tract and oral infections	Allergic reactions, phototoxicity, <u>liver dysfunction</u>
Ephedra (ma huang, sea grape, yellow horse)	Bronchodilator, diet aid, CNS stimulant	Hypertension, cardiac arrhythmias, stroke, seizures
Germander	Diet aid, digestive aid, gastrointestinal dysfunction	Multiple cases of <u>hepatitis and death</u> . (Still used as flavoring agent in <u>United States</u>)
Jimsonweed (<i>Datura</i> , devil's apple, stinkweed)	Respiratory tract diseases, hallucinogen	Marked adverse effects due to atropine and related M-blockers
Pennyroyal	Abortifacient, digestive aid, induction of menstrual flow	Gastrointestinal distress (hematemesis), CNS dysfunction, <u>hepatotoxicity</u> , renal dysfunction, disseminated intravascular coagulation
Pokeweed berries and root (American nightshade)	Berries used as food coloring; root extracts for emesis, rheumatism	Oral use highly toxic: bloody diarrhea, hypotension, coma, blindness, respiratory failure
Royal jelly	Tonic, immune potentiation, hyperlipidemia	Allergic reactions including anaphylaxis and death
Sassafras	Blood thinner, urinary tract disorders; oil used topically as antiseptic	Diaphoresis, hot flushes with oral use of bark; ingestion of sassafras oil may be lethal (coma, cardiovascular collapse and respiratory paralysis)