

Your project title

Last!: Parker Dingman, Ethan Donecoff, Karam Oubari, Pei Yi Zhuo

2021-04-25

Your written report goes here! Before you submit, make sure your code chunks are turned off with `echo = FALSE` and there are no warnings or messages with `warning = FALSE` and `message = FALSE`

Introduction and Data

Research question, background, and data

In this research project, we hope to most accurately predict life expectancy using economic, social, and health-related country level-data in conjunction with the regression analysis method, multiple linear regression. Therefore our research question is: how can we most accurately predict a country's average life expectancy?

By telling us the average age of death in a population, life expectancy is a key metric for understanding a country's health. According to Max Roser, Esteban Ortiz-Ospina and Hannah Ritchie from Our World in Data, "Broader than the narrow metric of the infant and child mortality, which focus solely at mortality at a young age, life expectancy captures the mortality along the entire life course." [1]

Over the course of history, life expectancy has risen dramatically. It is estimated that in pre-modern times, life expectancy worldwide was only about 30 years. Since the Industrial Revolution in the 18th and 19th centuries, many countries had huge gains in this number. And since the beginning of the 20th century, global average life expectancy has risen to about 70 years. However, there remain huge inequalities in this number. Currently (as of 2019), the Central African Republic has the lowest life expectancy of 53 years while Japan has the highest with 83.

In addition to the more obvious health-related connections to life expectancy, numerous pieces of academic literature have delved into the non-medical factors behind life expectancy. A major example is a longitudinal study conducted by Charles Lin, Eugene Rogot, Norman Johnson, Paul Sorlie, and Elizabeth Arias, which examined life expectancy by socioeconomic factors. [2] Academic literature such as this provides us with motivations to examine this topic on an international level, looking at various health-related and non-health-related factors that connect to life expectancy.

In terms of initial hypotheses of model selection, we expect that stronger predictors of life expectancy will be **Adult Mortality**, **infant deaths**, **under-five deaths**, and **Total expenditure**. We also predict that countries that have **Status** equal to "Developed" will have higher life expectancy than those that have **status** equal to "Developing".

Our primary data set is comprised of information that had been gleaned from the websites of the World Health Organization and the United Nations. [3] Each entry describes the health, social, and economic conditions for one of 193 countries in a given year from 2000 to 2015. [3] Because this data set lacks a variable that specifies the region of each country, we joined it with another data set that pairs country and region. [4] This secondary data set compiles information that can be found on the CIA World Factbook. [4] Both data were found on Kaggle and credit is given to Deeksha Russell and Duan Wang for creating the datasets.

Definitions of Relevant Variables:

Life expectancy: Average life expectancy in years (Response variable)

Country: Name of country (Identifier variable)

Region: Area, subcontinent, or continent where a country is located (Possible predictor variable)

Adult Mortality: Probability of dying between 15 and 60 years per 1000 population (Possible predictor variable)

infant deaths: Number of infant deaths per 1000 population (Possible predictor variable)

under-five deaths: Number of under-five deaths per 1000 population (Possible predictor variable)

Total expenditure: General government expenditure on health as a percentage of total government expenditure (Possible predictor variable)

Status: Developed or developing status (Possible predictor variable)

Exploratory Data Analysis

First, we will look at some summary statistics of our response variable, **Life expectancy**. Though it is possible to analyze the entire data set over all years, this creates difficulties with creating models. As one might expect, the average global life expectancy increased from 2000 to 2015. This relationship might make it unclear whether a rise in life expectancy is explained by our predictor variables or if it is simply due to human development over time. As a result, we will only use data from one year to perform our analysis.




We will only analyze life expectancy for the year 2011, because it is the latest year among those years with the least missing data (no variable has a missing rate greater than 22%).

Table 1: Summary Statistics of Response Variable, Life Expectancy


min	max	range	mean	median	Q1	Q3	iqr	sd
48.9	88	39.1	70.654	73.3	63.9	76.1	12.2	8.925

These summary statistics give a rough idea of the distribution of the response variable. The median life expectancy (~73.3 yrs) is almost 3 years larger than the mean (~70.7 yrs). Additionally, the median is closer to the third quartile than the first quartile. This suggests that life expectancy may be left-skewed, which we will evaluate further with visualizations.

Now, we will look at some summary visualizations of life expectancy, as well as other possibly relevant predictor variables.




Here we see that life expectancy is left skewed with a center just over 70 years. Life expectancy ranges from about 50 years to 90 years. Additionally, we see that a strong negative relationship between life expectancy and adult mortality, as well as much higher life expectancies among countries with `status` “developed” than “developing”.



Here we can see the differences in life expectancy across the world. For example, it is clear that life expectancy is generally higher in North American and European countries than African countries.

Correlated Predictors



Methodology

In order to create a model capable of predicting life expectancy, we use model selection to cull the number of potential predictor variables. We then assess whether certain interaction terms should be added to the model before addressing multicollinearity. Finally, with the set of predictor variables finalized, we ascertain whether any variable transformations are necessary through condition checking and identify outliers by examining model diagnostics.

Model Selection

The response variable we will be using in this project is **Life expectancy**. This is a measure of the average age of death in a year for the given country during that year. In terms of regression model technique, we will be using multiple linear regression because our response variable is quantitative.

The goal of our analysis is to calculate the most precise prediction of the response variable (**Life expectancy**). To do this we will perform forward and backward selection using AIC and BIC. Forward selection entails adding variables recursively using AIC or BIC as the criterion while backward selection means dropping variables one at a time that are deemed irrelevant based on AIC or BIC.

Table 2: Potential Models

Selection Method	AIC	BIC	Adjusted R-squared
Backward (AIC)	636.282	699.368	0.918
Backward (BIC)	654.341	674.413	0.895
Forward (AIC)	636.413	688.029	0.915
Forward (BIC)	654.341	674.413	0.895

Based on AIC and adjusted R^2 , we prefer the model we found through backward selection using AIC. This model results in the highest adjusted R^2 out of all four models. Moreover, this model is the only model out of the four that is superior to the remaining three models in more than one metric.

Potential Interaction Terms

Before proceeding with the model, we will look at some potential interaction terms to see if they should be added to the selected model. We will begin by looking at the interaction between Region and HIV/AIDS. This is because we might suspect that the impact of HIV/AIDS on life expectancy changes by region due to disparate access to relatively new treatments. The interaction between Region and Income composition of resources is also shown. The table below shows the selected model including these interaction terms, but only displays the interaction terms.

Table 3: Selected Model Interaction Terms

term	estimate	std.error	statistic	p.value
RegionBALTICS:HIV/AIDS	NA	NA	NA	NA
RegionC.W. OF IND. STATES:HIV/AIDS	-8.530	14.324	-0.595	0.553
RegionEASTERN EUROPE:HIV/AIDS	NA	NA	NA	NA
RegionLATIN AMER. & CARIB:HIV/AIDS	-7.502	6.365	-1.179	0.241
RegionNEAR EAST:HIV/AIDS	NA	NA	NA	NA
RegionNORTHERN AFRICA:HIV/AIDS	NA	NA	NA	NA
RegionNORTHERN AMERICA:HIV/AIDS	NA	NA	NA	NA
RegionOCEANIA:HIV/AIDS	-6.167	6.775	-0.910	0.365
RegionSUB-SAHARAN AFRICA:HIV/AIDS	-4.655	5.696	-0.817	0.416
RegionWESTERN EUROPE:HIV/AIDS	NA	NA	NA	NA
RegionBALTICS:Income composition of resources	24.692	134.802	0.183	0.855
RegionC.W. OF IND. STATES:Income composition of resources	-7.027	17.522	-0.401	0.689
RegionEASTERN EUROPE:Income composition of resources	-32.847	27.206	-1.207	0.230
RegionLATIN AMER. & CARIB:Income composition of resources	-9.491	13.462	-0.705	0.483
RegionNEAR EAST:Income composition of resources	-15.406	16.364	-0.941	0.349
RegionNORTHERN AFRICA:Income composition of resources	-10.469	31.918	-0.328	0.744
RegionNORTHERN AMERICA:Income composition of resources	NA	NA	NA	NA
RegionOCEANIA:Income composition of resources	-4.197	11.762	-0.357	0.722
RegionSUB-SAHARAN AFRICA:Income composition of resources	-1.952	9.103	-0.214	0.831
RegionWESTERN EUROPE:Income composition of resources	-36.022	28.032	-1.285	0.202

As shown in the table above, some of the groups do not have enough data to generate model coefficients. For the model coefficients calculated, there are no significant interaction terms for the variables investigated. Thus, there is not enough evidence to say that there is a significant interaction term between Region and HIV/AIDS or Region and Income composition of resources.

Multicollinearity

Table 4: Predictor Variable VIFs

Variable	VIF
RegionBALTICS	1.408
RegionC.W. OF IND. STATES	1.899
RegionEASTERN EUROPE	1.920
RegionLATIN AMER. & CARIB	2.742
RegionNEAR EAST	1.721
RegionNORTHERN AFRICA	1.260
RegionNORTHERN AMERICA	1.441
RegionOCEANIA	1.630
RegionSUB-SAHARAN AFRICA	4.657
RegionWESTERN EUROPE	3.002
Adult Mortality	2.254
infant deaths	244.366
Hepatitis B	2.972
Measles	3.156
under-five deaths	244.039
Total expenditure	1.328
Diphtheria	2.777
HIV/AIDS	2.016
Income composition of resources	13.166
Schooling	8.937

Variables with a $VIF > 10$ will have issues with multicollinearity. **infant deaths** and **under-five deaths** are clearly highly correlated (this makes a lot of sense in the context of the data). **Income composition of resources** appears to be correlated with **Schooling**. These two relationships were visualized in the EDA.

We should try models without **infant deaths** or without **under-five deaths** and then use model comparison techniques to decide on which of these two variables should be removed. Likewise, we ought to compare models with only **Income composition of resources** or **Schooling** and keep only one of the two.

Table 5: Infant Deaths vs. Under-five Deaths

Included Variable	AIC	BIC	adj.r.squared
Infant Deaths	637.3430	697.5612	0.9164
Under-five Deaths	637.5806	697.7988	0.9162

When comparing two models that are identical except that one includes **under-five deaths** and the other includes **infant deaths**, the one that includes **infant deaths** has lower values of AIC and BIC as well as a higher value of adjusted R^2 .


Table 6: Schooling vs. Income Composition of Resources

Included Variable	AIC	BIC	adj.r.squared
Schooling	695.555	752.905	0.868
Income Composition of Resources	642.335	699.686	0.913


Likewise, given two models that have the same predictors except that one includes **Schooling** while the other

includes Income composition of resources, the one that includes Income composition of resources is superior in terms of AIC, BIC, and adjusted R^2 .

Model Conditions and Diagnostics




Judging from the histogram and QQ plot above, the standardized residuals appear to be normally distributed, thus fulfilling the normality condition.



The plot above depicts standardized residuals on the y-axis and predicted values on the x-axis. We see no major departure from the constant variance assumption. However, two countries, Iraq and Lesotho are moderate outliers and two, Angola and Sierra Leone, are severe outliers in terms of their life expectancy. Specifically, Iraq has a much higher life expectancy than what we would expect given the values of its predictor variables while Angola, Lesotho, and Sierra Leone have far lower values for life expectancy than what was predicted.

Checking Linearity



The standardized residuals do not display a discernible pattern when plotted against any of the quantitative variables. This combined with the similar lack of any relationship in the plot of standardized residuals vs. predicted values above indicates that the linearity assumption holds.

Table 7: Predictor Variable Outliers

Country	Leverage
Algeria	0.345
Canada	1.000
Estonia	0.335
India	0.735
Indonesia	0.379
Latvia	0.339
Lithuania	0.345
Morocco	0.343
Myanmar	0.334
Netherlands	0.390
Tunisia	0.337
Ukraine	0.358

The 12 countries in the table above have high leverage (leverage > leverage_threshold = 0.3230769) while no countries are influential (Cook's Distance > 0.5).

Results

Table 8: Selected Model

term	estimate	std.error	statistic	p.value
(Intercept)	52.013	2.401	21.668	0.000
RegionBALTICS	-3.263	1.807	-1.805	0.074
RegionC.W. OF IND. STATES	-2.618	1.171	-2.237	0.027
RegionEASTERN EUROPE	-0.322	1.298	-0.249	0.804
RegionLATIN AMER. & CARIB	1.054	1.006	1.048	0.297
RegionNEAR EAST	1.110	1.392	0.798	0.427
RegionNORTHERN AFRICA	0.636	1.723	0.369	0.713
RegionNORTHERN AMERICA	4.466	3.187	1.401	0.164
RegionOCEANIA	-0.759	1.199	-0.633	0.528
RegionSUB-SAHARAN AFRICA	-2.576	1.030	-2.500	0.014
RegionWESTERN EUROPE	4.905	1.296	3.784	0.000
Adult Mortality	-0.013	0.003	-4.335	0.000
infant deaths	0.003	0.004	0.828	0.409
Hepatitis B	0.034	0.018	1.881	0.063
Measles	0.000	0.000	-1.679	0.096
Total expenditure	0.125	0.107	1.176	0.242
Diphtheria	-0.035	0.023	-1.517	0.132
HIV/AIDS	-0.620	0.118	-5.271	0.000
Income composition of resources	31.747	2.930	10.834	0.000

92.47% of the variation in the Life expectancy is explained by the regression model above, which contains Region, Adult Mortality, infant deaths, Hepatitis B, Measles, under-five deaths, Total expenditure, Diphtheria, HIV/AIDS, Income composition of resources and Schooling.

Appendix

Table 9: Full Model With All Possible Predictor Variables

term	estimate	std.error	statistic	p.value
(Intercept)	49.126	3.094	15.876	0.000
RegionBALTICS	-1.206	2.052	-0.588	0.558
RegionC.W. OF IND. STATES	-1.763	1.473	-1.197	0.234
RegionEASTERN EUROPE	0.980	1.700	0.577	0.565
RegionLATIN AMER. & CARIB	1.931	1.336	1.445	0.151
RegionNEAR EAST	1.333	1.678	0.795	0.429
RegionNORTHERN AFRICA	1.399	1.855	0.754	0.452
RegionNORTHERN AMERICA	5.640	3.376	1.671	0.098
RegionOCEANIA	0.458	1.641	0.279	0.781
RegionSUB-SAHARAN AFRICA	-1.361	1.225	-1.111	0.269
RegionWESTERN EUROPE	6.388	1.773	3.603	0.000
StatusDeveloping	1.105	1.243	0.888	0.376
Adult Mortality	-0.014	0.003	-4.443	0.000
infant deaths	0.091	0.041	2.249	0.027
Alcohol	0.008	0.106	0.079	0.937
percentage expenditure	0.001	0.001	1.204	0.232
Hepatitis B	0.050	0.019	2.623	0.010
Measles	0.000	0.000	-1.125	0.263
BMI	0.007	0.021	0.324	0.747
under-five deaths	-0.064	0.029	-2.184	0.031
Polio	-0.020	0.014	-1.440	0.153
Total expenditure	0.141	0.117	1.206	0.231
Diphtheria	-0.036	0.024	-1.520	0.132
HIV/AIDS	-0.529	0.121	-4.355	0.000
GDP	0.000	0.000	-1.036	0.303
Population	0.000	0.000	-1.292	0.199
thinness 1-19 years	0.199	0.272	0.731	0.466
thinness 5-9 years	-0.099	0.262	-0.378	0.706
Income composition of resources	43.132	6.029	7.154	0.000
Schooling	-0.548	0.250	-2.192	0.031

Sources

- [1] <https://ourworldindata.org/life-expectancy>
- [2] <https://europepmc.org/article/med/12785422/reload=0#impact>
- [3] <https://www.kaggle.com/kumaraajarshi/life-expectancy-who>.
- [4] <https://www.kaggle.com/fernandol/countries-of-the-world>