

EEE4113F Literature Review



Prepared by:

Karan Abraham

Lundi Mabasa

Mpilo Nxasana

Michael Van Staden

Prepared for:

EEE4113F

Department of Electrical Engineering

University of Cape Town

May 5, 2024

Declaration

1. I know that plagiarism is wrong. Plagiarism is to use another's work and pretend that it is one's own.
2. I have used the IEEE convention for citation and referencing. Each contribution to, and quotation in, this report from the work(s) of other people has been attributed, and has been cited and referenced.
3. This report is my own work.
4. I have not allowed, and will not allow, anyone to copy my work with the intention of passing it off as their own work or part thereof.



Karan Abraham



Lundi Mabasa

May 5, 2024

Date

May 5, 2024

Date



May 5, 2024

Mpilo Nxasana

Date



May 5, 2024

Michael Van Staden

Date

Contents

List of Figures	v
Abbreviations	vi
1 Introduction	1
1.1 Background	1
1.1.1 Problem Statment	1
1.2 Objectives	1
1.3 System Requirements	1
1.4 Scope & Limitations	1
1.5 Report Outline	1
2 Literature Review	2
2.1 Introduction	2
2.2 The Importance of Monitoring Avian Weight Changes	2
2.3 Current Weighing Methods	2
2.3.1 Spring Scales	3
2.3.2 Electronic Scales	3
2.3.3 Perching Scales	3
2.4 Data Transmission and User Interface	3
2.5 Power supply	4
2.5.1 Wall Power	5
2.5.2 Energy Harvesting	5
2.5.3 Limitations	5
2.6 Challenges and Considerations	5
2.6.1 Size and Species	5
2.6.2 Environmental Conditions	5
2.6.3 Material Considerations	6
2.6.4 Ethical Considerations	6
2.7 Conclusion	6
3 Sensing Subsystem (NXSMPI001)	8
3.1 Introduction	8
4 Conclusions	9
5 Recommendations	10

Bibliography

11

List of Figures

2.1 Overall Performance of each Protocol [1] 4

Abbreviations

Chapter 1

Introduction

Philosophers have hitherto only interpreted the world in various ways;
the point is to change it.

—*Karl Marx*

1.1 Background

1.1.1 Problem Statment

Sally, a researcher at the FitzPatrick Institute, needs a way to weigh the red-winged starlings without startling them, because interestingly, they record their weight manually using a kitchen scale.

1.2 Objectives

1.3 System Requirements

1.4 Scope & Limitations

1.5 Report Outline

Chapter 2

Literature Review

2.1 Introduction

Weight is an important metric for evaluating the overall health, behavior and ecology of a bird in ornithological research. However obtaining this weight data presents several challenges due to their size, fragility and often rapid movements. This literature aims to explore and evaluate the different techniques and technologies researchers employ to obtain weight data, the different ways this data could be transmitted and recorded, and the types of devices needed to power such a system. This literature review will also consider the challenges described above and, other environmental and ethical factors that must be taken into account when weighing birds in the wild.

2.2 The Importance of Monitoring Avian Weight Changes

According to Clark, “Weight summarizes the total biomass of an individual and is probably the most convenient standard of energetic comparisons.” [2], which emphasizes how valuable the weight data of an individual bird can be. Baldwin and Kendeigh state that “The weight of birds and the variations and fluctuations of these weights furnish criteria of considerable importance in the understanding of the physiological and ecological reactions of the bird as a living organism [3], which further establishes how important the tracking of an individual bird’s weight data can be. Both of these statements cement how valuable weight data can be for ornithologists studying individual birds.

Clark goes on to say, “Weights have been used in analyses of the factors that influence differences in species diversity between communities.” [2]. They also state that “Weights and census data have often been combined to calculate the total biomass of a particular species or group of species in an area”. These statements add that weight data could also be invaluable for the analysis of an entire community of birds as opposed to just monitoring individuals.

2.3 Current Weighing Methods

This section examines the different methods and tools used to obtain weight data in ornithological research today.

2.3.1 Spring Scales

Spring scales measure weight based on the extension of a spring when a force (the weight of the bird) is applied. Their main advantage, as described by Manolis [4], is that they are “relatively inexpensive and sufficiently portable to suffice for short-term field project[s]”. However, within the same study, the scale was only accurate to within 0,5g and when smaller birds can weigh less than 50g, spring scales may lack the precision for such research applications.

2.3.2 Electronic Scales

Electronic scales utilize load cells or strain gauges to convert the weight of the bird into an electronic signal, which can then be displayed on a digital screen. These scales offer precise measurements as shown by Carpenter et al. [5], where they were able to improve the precision of their measurements from 0,05g to 0,01g, by simply replacing their spring scales with electronic ones. Another advantage over spring scales, is that they do not have to be recalibrated after moving [5] and they tend to come with features such as taring functions to account for the weight of the housing holding the bird.

2.3.3 Perching Scales

Perching scales integrate a weighing platform into an artificial perch or nest. In Poole and Shoukimas’ [6] study, birds landing on perch would deflect a transducer (a metal beam with 4 strain gauges bonded to it), thus generating an electronic signal. Reid et al. [7] used artificial nests rigged with a load cell in much the same way. In both studies, these electronic signals would then be recorded via some kind of electronic storage medium. This meant the birds could be weighed remotely, which minimizes stress and reduces the risk of injury, making perching scales particularly useful for long-term monitoring studies or behavioural observations. However, for such long-term studies, researchers would need to keep track of a large number of birds, which would also result in a large amount of data that needs to be stored.

Manolis [4] provides a solution to these issues by urging other researchers to make use of telemetry. One such technology is Radio Frequency Identification (RFID) which enables researchers to track individual birds and record their weight automatically. Wang et al. [8] made use of RFID by attaching two transponders to each bird, which would be detected by antennas placed under the perches. When a tagged bird interacts with the RFID reader, its unique identifier and weight are recorded electronically, making the data much easier to organise. It also reduces the volume of data created as the weight is only taken when the birds are on top of the perch. This allows researchers to collect data on a larger scale but this data need not be stored locally.

2.4 Data Transmission and User Interface

The method that the FitzPatrick Institute currently have for reading the bird weight is having one of the researchers go up to the scale and read off the screen. They then record the weight into *Cybertracker* [9], a mobile app that creates an Excel spreadsheet for them to analyze later. This highlights a need for a way to access that data remotely; or perhaps, a way to send that data directly to the Cybertracker app.

There are many communication protocols for transmitting data from a microprocessor. In a comparative performance study by Eridani et al., three protocols were compared: Bluetooth, Wi-Fi direct, and ESP-NOW (‘a new protocol that allows multiple devices to communicate with each other without the use of Wi-Fi, with low power consumption’ [1]). Five metrics were used in the tests: maximum range, transmission speed, latency, power usage, and signal resistance to obstructions [1]. A brief summary of the performance of each protocol is shown below in Figure 2.1.




Figure 2.1: Overall Performance of each Protocol [1]

ESP-NOW performs best in range and latency; Bluetooth in power usage; and Wi-Fi in transmission speed. Since the scale must be portable, it is important to keep power usage to a minimum. Bluetooth seems to provide sufficient range and speed. The problem with this; however, is that connecting the system to the user’s phone requires effort, and perhaps expertise that the user may not have. In this case, connecting the system to the internet may be a better option (that is if an internet connection is available, i.e. if *eduroam* is in range).

Budoyo and Andriana used the internet when designing a prototype of a digital scale to measure the weight of onions. [10]. They interfaced the microcontroller (an ATmega2560) to the internet using an ESP8266 Wi-Fi module. The weight data is sent to a website where it is stored in a database. A database is useful in creating an Excel spreadsheet with many fields which is the end product that the client requires.

2.5 Power supply

Traditional weighing scales have relied on either battery or electric power sources for operation. Battery-powered scales utilize internal batteries, commonly alkaline or lithium-ion, to supply the necessary electrical power. This section will discuss the different types of power supplies available for bird scales and the power limitations on what type of power source the final design can use.

2.5.1 Wall Power

For indoor scales, which are fixed to one spot, the electric-type scales are directly connected to a power source via a cord, typically drawing from AC power provided by a wall outlet. These type of scales are used in laboratories and residential environments.

2.5.2 Energy Harvesting

One other method researchers use in low powered bird scales is Energy Harvesting. Energy Harvesting is used to extend the lifespan of the scale and sensing devices, however the process is not always effective [11]. An example of energy harvesting would be using wind or solar as a power source to the scale device. This concept is useful in situations where the scale is left in the field for data capture, and only accessed after a prolonged period of time. It is worth noting that low-power weighing scales are an existing topic, where in some cases there are scales and sensors that are able to take measurements, read and communicate the data in real time [12].

2.5.3 Limitations

The reliance on electrical power poses serious constraints, particularly in terms of mobility. This limitation becomes pronounced in specialized applications such as bird weighing scales, especially for very mobile birds such as the starlings. A better solution is to use a rechargeable battery source, such as the alkaline batteries mentioned above. Environmental conditions also pose a risk to the battery lifespan. Solar panels can pose collision risks for birds, particularly if the panels are highly reflective. Some birds may collide with solar panels while flying, leading to injury or mortality.

2.6 Challenges and Considerations

While modern weighing methods offer significant advantages in terms of accuracy, convenience, and animal welfare, researchers must consider several factors when selecting the most appropriate technique for their study.

2.6.1 Size and Species

The size and behaviour of the target bird species may influence the suitability of different weighing methods. Some birds may become skittish around researchers which would make measurements unreliable. In Manolis' case [4], they had to use binoculars to take readings of the scale from afar; an inconvenience that is entirely removed from the solutions presented by Poole and Shoukimas [6] and Reid et al. [7]. Smaller birds may require scales with higher precision, while larger species may benefit from perching scale systems capable of handling multiple subjects simultaneously.

2.6.2 Environmental Conditions

Field studies often expose equipment to challenging environmental conditions. For example, Manolis [4] had to keep swaying to a minimum to get accurate readings, hence spring scales would not be suitable in windy conditions. Rain can seriously damage electronic components so it was important for Reid et al. [7] to house the amplifier unit in "a small water-resistant case with a sealed connection to

the data logger cable”. Researchers must choose weighing methods that are robust and reliable under these circumstances, with weather-resistant features where necessary.

2.6.3 Material Considerations

An important consideration that our design would need to fulfill is that it is safe for use on birds. Given that they will be in direct contact with the scale, toxicity is a primary concern when considering the materials used to construct the scale.

In the design of an electronic scale, artificial materials such as plastics are an attractive option due to their naturally weatherproof properties. However, caution must be exercised when considering specific materials. Artificial materials such as polytetrafluoroethylene (PTFE) are a common source for airborne toxicity in avians [13]. However, in a paper by Kroshefsky [14] this material is only cause for toxicity concern when exposed under high temperatures. This is because PTFE begins to decompose in air at around 200°C. Even if the scale is place outdoors, the ambient temperature will be well under this temperature limit, making this material a viable choice in the housing.

Heavy metal poisoning is another important concern when it comes to our choice of materials. The most common occurrences of which come from the ingestion of lead [15]. When lead is ingested, it can be absorbed in the gastrointestinal tract and then taken up by soft tissues and eventually bone [15]. The paper by C. Pollock includes a list of common sources of lead in household items, the most notable of which is solder. Thus some kind of insulation is required around soldered circuitry to avoid any trace amounts of lead affecting the birds.

2.6.4 Ethical Considerations

Ethical guidelines emphasize the importance of minimizing stress and harm to the animals being studied. As such researchers should prioritize methods that avoid the need to handle the birds, as this will minimize the risk of injury to the researchers and the birds. This means that the red-winged starlings will need to be lured such that their weight measurements can be taken. This can be achieved using the many perching scale solutions described earlier, but for more traditional weighing methods, Manolis [7] provides a solution. In their study they used sunflower seeds to entice the birds to land on the scale, and since red-winged starlings tend to scavenge for food, this will prove useful for this application as well.

An automatic feeder device could be implemented as a way to streamline this process. However, researchers have found that the introduction of automated feeders tend to reduce species variety in a given area [16]. Automated feeders are also highly inconsistent with seasonal change, which would result in it being a highly inconsistent form of luring birds [16]. The introduction of an automated feeding strategy would have a noticeable effect on the ecological and weight cycles of local birds, especially those that would be studied.

2.7 Conclusion

Based on the reviewed literature, it can be seen that accessing data remotely is very much feasible using accessible technology such as Wi-Fi. The importance of gathering weight data was established

and how it would be of benefit to ornithologists was explored. Various challenges and considerations have been presented in the reviewed literature and the importance thereof will be taken into account going forward.

Chapter 3

Sensing Subsystem (NXSMPI001)

3.1 Introduction

Chapter 4

Conclusions

The same rule holds for us now, of course: we choose our next world through what we learn in this one. Learn nothing, and the next world is the same as this one.

—*Richard Bach, Jonathan Livingston Seagull*

The purpose of this project was to...

This report began with...

The literature review was followed in Chapter...

The bulk of the work for this project followed next, in Chapter...

In Chapter...

Finally, Chapter... attempted to...

In summary, the project achieved the goals that were set out, by designing and demonstrating...

Chapter 5

Recommendations

It is for us the living, rather, to be dedicated here to the unfinished work which they who fought here have thus far so nobly advanced.

—*Abraham Lincoln*

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Bibliography

- [1] D. Eridani, A. F. Rochim, and F. N. Cesara, “Comparative performance study of esp-now, wi-fi, bluetooth protocols based on range, transmission speed, latency, energy usage and barrier resistance,” in *2021 international seminar on application for technology of information and communication (iSemantic)*. IEEE, 2021, pp. 322–328.
- [2] G. A. C. Jr., “Body weights of birds: A review,” *The Condor*, pp. 193–202.
- [3] S. C. K. S. Prentiss Baldwin, “Variations in the weights of birds,” *The Auk*.
- [4] T. Manoils and A. Manolis, “A simple technique for weighing birds at feeders without recapture,” *North American Bird Bander*, vol. 6, no. 3, p. 4, 2024.
- [5] F. L. Carpenter, D. C. Paton, and M. A. Hixon, “Weight gain and adjustment of feeding territory size in migrant hummingbirds,” *Proceedings of the National Academy of Sciences*, vol. 80, no. 23, pp. 7259–7263, 1983.
- [6] A. Poole, “A scale for weighing birds at habitual perches,” *Journal of Field Ornithology*, vol. 53, no. 4, pp. 409–414, 1982.
- [7] K. Reid, G. M. Liddle, P. A. Prince, and J. P. Croxall, “Measurement of chick provisioning in antarctic prions pachyptila desolata using an automated weighing system,” *Journal of Avian Biology*, pp. 127–134, 1999.
- [8] K. Wang, K. Liu, H. Xin, L. Chai, Y. Wang, T. Fei, J. Oliveira, J. Pan, and Y. Ying, “An rfid-based automated individual perching monitoring system for group-housed poultry,” *Transactions of the ASABE*, vol. 62, no. 3, pp. 695–704, 2019.
- [9] CyberTracker. Cybertracker wiki. [Online]. Available: <https://cybertrackerwiki.org/getting-started/>
- [10] Y. Budoyo and A. D. Andriana, “The digital weight scale of iot system using load cell sensor in ud. pangrukti tani,” *Jurnal Ilmiah Komputer dan Informatika (KOMPUTA)*.
- [11] L. Meile, A. Ulrich, and M. Magno, “Wireless power transmission powering miniaturized low power iot devices: A review,” in *2019 IEEE 8th International Workshop on Advances in Sensors and Interfaces (IWASI)*, June 2019, pp. 312–317.
- [12] J. Paek, J. Hicks, S. Coe, and R. Govindan, “Image-based environmental monitoring sensor application using an embedded wireless sensor network,” *Sensors*, vol. 14, no. 9, pp. 15 981–16 002, 2014. [Online]. Available: <https://www.mdpi.com/1424-8220/14/9/15981>
- [13] T. L. Lightfoot, “Pet bird toxicity and related environmental concerns,” *Veterinary Clinics of North America: Exotic Animal Practice*, pp. 229–259, 2008.

- [14] R. D. Kroshefsky, “Teflon poisoning: How dangerous is your cooking to your birds?” *AFA Watchbird Magazine*, pp. 30–33, 1980.
- [15] C. P. Cynthia Chow, “Heavy metal poisoning in birds,” pp. 1–9, 2009.
- [16] J. R. B. K. P. M. C. S. Josie A. Galbraith, Darryl N. Jones, “Urban bird feeders dominated by a few species and individuals,” *Frontiers in Ecology and Evolution*.