

Department of Mathematics
Indian Institute of Technology Bhilai
IC152: Linear Algebra-II
Tutorial Sheet 5

1. Find the matrix of the following inner products relative to given ordered basis \mathcal{B}
 - (i) $V = P_2(\mathbb{R})$ with the inner product $\langle f, g \rangle = \int_0^1 f(x)g(x)dx$, $\mathcal{B} = \{1, x, x^2\}$
 - (ii) $V = \mathbb{C}^3$ with $\langle \alpha, \beta \rangle = \sum_{j=1}^3 \alpha_j \bar{\beta}_j$, with any ordered basis \mathcal{B} of \mathbb{C}^3 . (You can choose any ordered basis of your choice)
2. Apply Gram-Schmidt process to the following subsets S of inner product space V to get an orthonormal basis for span of S .
 - (i) $V = \mathbb{R}^3$ with standard inner product, $S = \{(1, 0, 1), (0, 1, 1), (1, 3, 3)\}$
 - (ii) $V = P_2(\mathbb{R})$ with the inner product $\langle f, g \rangle = \int_0^1 f(x)g(x)dx$, $S = \{1, x, x^2\}$.
 - (iii) $V = \mathbb{C}^4$, with standard inner product, $S = \{(1, i, 2 - i, -1), (2 + 3i, 3i, 1 - i, 2i), (-1 + 7i, 6 + 10i, 11 - 4i, 3 + 4i)\}$.
 - (iv) $V = M_{2 \times 2}(\mathbb{R})$ with standard inner product

$$S = \left\{ \begin{bmatrix} 3 & 5 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 9 \\ 5 & -1 \end{bmatrix}, \begin{bmatrix} 7 & -17 \\ 2 & -6 \end{bmatrix} \right\}$$

3. Prove the following
 - (i) Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be an orthonormal basis for an inner product space V . Prove that for any $\alpha, \beta \in V$

$$\langle \alpha, \beta \rangle = \sum_{i=1}^n \langle \alpha, \alpha_i \rangle \overline{\langle \beta, \alpha_i \rangle}$$

- (ii) Let $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be an orthonormal subset of an inner product space V . Prove that for any $\alpha \in V$

$$\|\alpha\|^2 \geq \sum_{i=1}^n |\langle \alpha, \alpha_i \rangle|^2$$

4. Compute S^\perp for $S = \{(1, 0, i), (1, 2, 1)\}$ in \mathbb{C}^3 .
5. Find the orthogonal projections of the following vectors on the given subspace of the specified inner product space
 - (i) $V = \mathbb{R}^3$ with standard inner product, $\alpha = (2, 1, 3)$, and $W = \{(x, y, z) : x + 3y - 2z = 0\}$.

(ii) $V = P(\mathbb{R})$ with the inner product $\langle f, g \rangle = \int_0^1 f(x)g(x)dx$, $h(x) = 4 + 3x - 2x^2$, $W = P_1(\mathbb{R})$.

6. Let V be an inner product space, S and S_0 are the subsets of V and W is a finite dimensional subspace of V . Prove the following results

(i) $S_0 \subseteq S$ implies that $S^\perp \subseteq S_0^\perp$.

(ii) $S \subseteq (S^\perp)^\perp$ implies $\text{span}(S) \subseteq (S^\perp)^\perp$

(iii) $W = (W^\perp)^\perp$

(iv) $V = W \oplus W^\perp$

7. Let V be an inner product space, and let W be a finite-dimensional subspace of V . If $x \notin W$, prove that there exists $y \in V$ such that $y \in W^\perp$, but $\langle x, y \rangle \neq 0$.

8. Let $V = C([-1, 1]; \mathbb{R})$ be an inner product space with inner product defined as

$$\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx.$$

Suppose that W_e and W_o denote the subspaces of V consisting of the even and odd functions, respectively. Then prove that $W_e^\perp = W_o$.