

SPIDOXAMAT	GROUP	23	INSECTICIDE
FLUPYRADIFURONE	GROUP	4D	INSECTICIDE

PLENEXOS™ SMART

(Note to Reviewer: Optional Text: [Powered by/ PLENEXOS/ technology] denotes the Bayer trademark for the active ingredient spidoxamat.)

Foliar applied insecticide for use in pest management, suppression of listed insects that can transmit disease causing [plant] pathogens and maintenance of plant health.

ACTIVE INGREDIENT(S):

Spidoxamat*7.2%

Flupyradifurone**15.0%

OTHER INGREDIENTS:77.8%

TOTAL:100.0%

Contains 0.072 pounds of spidoxamat and 0.15 pounds of flupyradifurone per pound of formulated product

*CAS No. 907187-07-9

**CAS No: 951659-40-8

EPA Reg. No. 264-REGG

EPA Est.No. _____

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours a Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

See [Back][Side] Panel for First Aid Instructions and [Leaflet][Booklet] for Complete Precautionary Statements and Directions for Use. (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

Net Contents:

[Batch No.: [for non-refillable containers]]

PRODUCED FOR



Bayer CropScience LLC
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)

FIRST AID

If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if available to swallow. • DO NOT induce vomiting unless told to by a poison center or doctor. • DO NOT give anything by mouth to an unconscious person.
If on Skin or Clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call poison control center or doctor for treatment advice.
If Inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
Note to Physician: No specific antidote is available. Treat patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Harmful if swallowed or absorbed through skin.
- Causes moderate eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Terrestrial Use

DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Non-Target Organisms

Flupyradifurone is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Flupyradifurone is highly toxic to adult *Megachile rotundata* (alfalfa leafcutting bee) via direct contact exposure. **DO NOT** apply to foliage when managed alfalfa leafcutting bees are foraging in the treatment area. Flupyradifurone is toxic to adult bees in laboratory studies via oral exposure. Not toxic to honey bees or bumble bees through contact exposure. Spidoxamat is potentially toxic to honey bees through chronic exposure to residues in pollen and nectar. Field studies conducted with flupyradifurone have shown no effects on honeybee colony development.

Ground Water Advisory

Spidoxamat and flupyradifurone have properties and characteristics associated with chemicals detected in groundwater. Spidoxamat and flupyradifurone may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. Spidoxamat is classified as having high potential for reaching surface water via runoff for several days after application. Flupyradifurone and its degradate difluoroacetic acid are classified as having medium and high potential, respectively, for reaching surface water via run-off for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of spidoxamat and flupyradifurone and its degradate difluoroacetic acid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Endangered Species Statement

The use of any pesticide in a manner that may kill or otherwise harm endangered species or adversely modify their habitat is a violation of Federal law.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSOURCE LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling
Read the entire label before using this product.**

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Not for sale, distribution or use in Nassau and Suffolk Counties New York except as permitted under FIFRA Section 24(c),
Special Local Need Registration.
No aerial application in New York State.**

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves (made of any waterproof material)
- Shoes plus socks

PRODUCT INFORMATION

PLENEXOS SMART:

- Is a water dispersible granule (WG) formulation and is active primarily by ingestion against immature target pest life stages. In addition, fertility of adult female target pests, such as aphids and whiteflies, may be reduced.
- Can be applied as a preventative treatment or timed to coincide with an early threshold level in developing insect populations.
- It is widely known that tank mixtures and/or sequential treatments of horticultural spray oil with captan and/or sulfur may cause adverse plant compatibility in tree and vine crops; including [this product/PLENEXOS SMART] in this tank-mix and/or sequential treatment scenario is not recommended.
- Following application to plant foliage, spidoxamat in [this product/PLENEXOS SMART] is fully systemic and flupyradifurone has xylem systemicity, moving through phloem and xylem to new shoot, leaf and root tissues; systemicity and efficacy may be hindered during periods of cold temperatures, under drought conditions, or when plants are not actively growing. Sufficient leaf tissue must be present for uptake and translocation of this product.

USE RESTRICTIONS

- Refer to the specific use directions and restrictions in each Crop, Crop Group or Crop Subgroup table.
- **DO NOT** tank mix with azole fungicides (FRAC group 3) during bloom period.
- Not for use on greenhouse or planthouse crops except where permitted in crop specific directions or allowed by state specific 24(c) labeling.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply prior to petal-fall on stone fruit crops.
- The use of Induce® adjuvant in combination with [this product/PLENEXOS SMART] on grape, pome fruits, and stone fruits is prohibited when fruit is present due to adverse plant compatibility on harvested commodities.
- No aerial application in New York State.
- **DO NOT** apply to crop foliage during foraging by alfalfa leafcutting bees used for pollination services.

APPLICATION INSTRUCTIONS

[This product/PLENEXOS SMART] may be applied as a foliar application using properly calibrated air-blast or similar ground application equipment, or through fixed or rotary winged aircraft. Sufficient spray volume, based on the size and density of the treated crop, must be utilized that allows for good coverage of both young and old foliage without runoff or collection of spray solution on leaf margins, fruit, or other plant tissues. Good coverage will help ensure maximum uptake by leaf surfaces and optimum systemicity within the plant. For optimum control of target pests on tree and vine crops, treating both sides of plant during the same application period is recommended;

for practices such as alternative middles or tops and bottoms, both sides of the trees or vines must be treated within a 72-hours period. Rates for tree and vine crops are based on full-size mature trees and vines.

- Ground application through air-blast or similar ground equipment. For tree and vine crops, application must be made in a minimum of 50 GPA for conventional ground air-blast sprayer and 30 GPA for high air velocity, low volume or air curtain sprayers. Air-assisted or air-blast sprayers move spray droplets into the crop canopy using a forced-air system. The fan must be set up to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area. Equip sprayers with nozzles that provide accurate and uniform application.
- Aerial applications must be made in a minimum of 10 GPA in tree and vine crops. The higher the dosage of [this product/PLENEXOS SMART] within the crop/pest-specific section may be necessary for optimum control of aerial applications.

ENDANGERED AND THREATENED SPECIES PROTECTION REQUIREMENTS

Before using this product, you must obtain any applicable Endangered Species Protection Bulletins ('Bulletins') within six months prior to or on the day of application. To obtain Bulletins, go to Bulletins Live! Two (BLT) at <https://www.epa.gov/pesticides/bulletins>. When using this product, you must follow all directions and restrictions contained in any applicable Bulletin(s) for the area where you are applying the product, including any restrictions on application timing if applicable. It is a violation of federal law to use this product in a manner inconsistent with its labeling, including this labeling instruction to follow all directions and restrictions contained in any applicable Bulletin(s). For general questions or technical help, call 1-844-447-3813, or email ESPP@epa.gov.

MANDATORY SPRAY DRIFT MANAGEMENT [(Continued)]

Applicators must access and search Bulletins Live! Two (BLT) at <https://www.epa.gov/pesticides/bulletins> within six months of the application to determine whether the application site falls within a Pesticide Use Limitation Area (PULA) that has a Bulletin in BLT. If you are located inside a PULA, follow the instructions in the bulletin.

For Aerial and Boomless Ground Applications

- Maintain a wind-directional (downwind) buffer distance (for aerial – 71 mph and for boomless ground – 45 feet) from the edge of the treated area
- The buffer footage distance may include the following managed areas, provided people are not present within the application exclusion zone during the application, and they will not be contacted by the pesticide, either directly or through drift
 - Agricultural fields, including untreated portions of the treated field
 - Roads, paved or gravel surfaces, mowed grassy areas adjacent to field, and areas of bare ground from recent plowing or grading that are contiguous with the treated area
 - Buildings and their perimeters, silos, or other man-made structures with walls and/or roof
 - Areas maintained as a mitigation measure for runoff/erosion or drift control, such as vegetative filter strips (VFS), field borders, hedgerows, Conservation Reserve Program (CRP) lands, and other mitigation measures identified by EPA on the mitigation menu
 - Managed wetlands including constructed wetlands on farm
 - On-farm contained irrigation water resources that are not connected to adjacent water bodies, including on-farm irrigation canals and ditches, water conveyances, managed irrigation/runoff retention basins, and tailwater collection ponds.

Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver coarse or coarser droplets as indicated in nozzle manufacturer's catalogs and in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641). If the windspeed is 10 mph or less, applicators must use ½ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 mph, applicators must use ¾ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Wind speed must be measured at the release height or higher, in an area free from obstructions such as trees, buildings, and farm equipment.
- Wind speed and direction must be measured on location using a windsock or anemometer (including systems to measure wind speed or velocity using application equipment).
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications (Airblast and Handheld Technology)

- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- Wind speed must be measured at the release height or higher, in an area free from obstructions such as trees, buildings, and farm equipment.
- Wind speed and direction must be measured on location using a windsock or anemometer (including systems to measure

MANDATORY SPRAY DRIFT MANAGEMENT [(Continued)]

- wind speed or velocity using application equipment).
- **DO NOT** apply during temperature inversions.
- Sprays must be directed into the canopy.
- User must turn off outward pointing nozzles at row ends and when spraying the outer row.

Buffer Restrictions for Aerial and Boomless Ground Applications

- **Aerial**
 - For foliar applications to all uses, 71 ft buffer is required between application area and closest downwind edge of non-target aquatic AND terrestrial habitat
- **Boomless Ground**
 - For foliar applications to all uses, 45 ft buffer is required between application area and closest downwind edge of non-target aquatic AND terrestrial habitat
- **Ecological Spray Drift Buffer Reduction Options** can be found on EPA's Mitigation Menu webpage (<https://www.epa.gov/pesticides/mitigation-menu#ecological-drift>)

SEE "ADDITIONAL SPRAY DRIFT INFORMATION" SECTION BELOW FOR MORE DETAILS.

ADDITIONAL SPRAY DRIFT INFORMATION:

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

CONTROLLING DROPLET SIZE - AIRCRAFT

Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aurally to crops, **DO NOT** release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.

MANAGEMENT OF RUNOFF/ EROSION

- DO NOT apply during rain.
- DO NOT apply this product when soil is saturated or above field capacity.
- Within 24 hours after application, DO NOT irrigate the treatment area to the point of runoff.

A variety of factors including soil type, slope, and weather conditions (e.g., rainfall) can influence volume and intensity of water running off the treated field. The applicator should evaluate factors and make appropriate adjustments when applying this product. Land

management, agronomic practices, field conditions, and application measures that reduce, to the maximum extent practicable, runoff from treated fields, should be implemented by land managers/users of this product.

RUNOFF/EROSION MITIGATION

Runoff/erosion mitigation is required for the crop uses listed below, unless otherwise specified.

- **Mitigation Points – For all uses on label – 3 points**

You must achieve a minimum of points for the crop uses. Each crop use is assigned a minimum number of mitigation points that must be achieved using one or more runoff/erosion mitigation measures. The mitigation points were determined for each crop use based on the likelihood that runoff or erosion from use of this product in the crop's production could result in harm to non-target plants or animals.

Applicators must access and search Bulletins Live! Two (BLT) at www.epa.gov/pesticides/bulletins within six months of the application to determine whether the application site falls within a Pesticide Use Limitation Area (PULA) that has a Bulletin in BLT. If you are located inside a PULA, follow the instructions in the bulletin.

If the application site is located outside a PULA, runoff/erosion mitigation is required for this product unless certain field/application parameters are present at the time of application (i.e., subsurface or tile drains with controlled outlet, perimeter berm systems, irrigation tailwater return systems, spot treatment, etc.). Access EPA's Mitigation Menu Website at www.epa.gov/pesticides/mitigation-menu for a full list of field/application parameters to evaluate whether your field is subject to runoff/erosion mitigation.

If the application does not meet the specified field/application parameters, a minimum of 3 points for the crop uses listed on the label must be achieved. The applicator must choose among the mitigation and/or mitigation relief measures on EPA's Mitigation Menu Website to meet or exceed these points before applying this product. The website includes the full menu of runoff/erosion mitigation and mitigation relief measures. The following are examples:

- Location in a very low, low, or medium runoff vulnerability county
- Field slope
- Conservation tillage
- Vegetative strips
- Cover crop or continuous ground cover
- Irrigation water management
- Mulching
- Grassed waterway
- Vegetative ditch
- Constructed and natural wetlands
- Water retention systems
- Following recommendations from a runoff specialist or participating in a qualifying conservation program (see www.epa.gov/pesticides/mitigation-menu for minimum elements).

To achieve mitigation points for the application, the mitigation and mitigation relief measures must be:

- Employed in accordance with the instructions and descriptions on EPA's **Mitigation Menu Website**.
- In place during the application unless a different timing (such as before or after application) is specifically provided in the measure's description on EPA's **Mitigation Menu Website**.

EPA may periodically update the **Mitigation Menu Website**, for example, by adding new mitigation measures or updating a mitigation measure description.

COMPATIBILITY TESTING AND TANK MIX PARTNERS

- If [this product/ PLENEXOS SMART] is to be tank mixed with other pesticides, compatibility should be tested prior to mixing.
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

[THIS PRODUCT/ PLENEXOS SMART] must be tank-mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations by crop. However, the use of Induce[®] adjuvant in combination with [this product/ PLENEXOS SMART] on grape, pome fruits, and stone fruits is prohibited when fruit is present due to adverse plant compatibility on harvested commodities. The tank-mixture of [this product/ PLENEXOS SMART] with an adjuvant/ additive having sticking properties or crop protection product formulations containing built-in stickers have been shown to interfere with leaf uptake and should be avoided. Sufficient leaf tissue must be present for uptake and translocation of this product.

It is widely known that tank mixtures and/or sequential treatments of horticultural spray oil with captan and/or sulfur may cause adverse plant compatibility in tree and vine crops; including [this product/ PLENEXOS SMART] in this tank mix and/or sequential treatment scenario is not recommended.

Compatibility

[This product/ PLENEXOS SMART] is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. However, it is known that many components, including crop protection products, fertilizers, micronutrients, and spray adjuvants, may be present in a tank mix combination. There is potential for adverse chemical reactions. It is impossible to determine physical, biological, and plant compatibility for all scenarios that may be encountered; therefore, it is recommended that users determine the chemical, physical, biological and plant compatibility of such mixes prior to application on a broad commercial scale

Order of Mixing

[This product/ PLENEXOS SMART] may be used with other recommended pesticides, fertilizers, and micronutrients. The proper mixing procedure for [this product/ PLENEXOS SMART] alone or in tank mix combinations with other pesticides is:

1. Fill the tank 1/4 to 1/3 full with clean water;
2. While recirculating and with the agitator running, add any products in Polyvinyl acetate (PVA) bags (See Note). Allow time for thorough mixing;
3. Continue to fill tank with water until 1/2 full;
4. Add any wettable powder (WP) or wettable granules (WG) products;
5. Add the required amount of [this product/ PLENEXOS SMART/ (WG)];
6. Add any "flowable" (FL or SC) type products;
7. Allow enough time for thorough mixing of each product added to tank;
8. If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers and micronutrients;
9. Fill tank to desired level and maintain constant agitation to ensure uniformity of mixture.

NOTE: **DO NOT** use PVA packets in a tank mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic that is not soluble in water or solvents.

INSECTICIDE RESISTANCE MANAGEMENT (IRM) RECOMMENDATIONS

For resistance management, [this product/ PLENEXOS SMART] contains a Group 23 and 4D insecticides. Any insect population may contain individuals naturally resistant to [this product/ PLENEXOS SMART/, spidoxamat or flupyradifurone containing products] and other Group 23 and 4D insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take one or more of the following steps:

- **DO NOT** exclusively use [this product/ PLENEXOS SMART/, spidoxamat or flupyradifurone containing products] or other Group 23 or 4D insecticides to control the same pest throughout the season. Avoid exposing consecutive generations of a pest to the same mode of action by using the "application window" approach which rotates products with different MoA chemistries.
- An "application window" is the duration of an insect generation or if unknown, then use an approximate 30-day period. Rotate windows of treatments of [this product/ PLENEXOS SMART/, spidoxamat or flupyradifurone containing products] and other Group 23 or 4D products followed by blocks of treatments with other effective products from different modes of action before returning to Group 23 or 4D products.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Use recommended label rates, timings, and spray intervals with maintained application equipment to reduce insect resistance risk.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Visit the Insecticide Resistance Action Committee (IRAC) website at <http://www.irac-online.org> and or contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937).

ROTATIONAL CROPS

[This product/ PLENEXOS SMART] is labeled for use on the following crops:

Citrus Fruit (crop group 10-10), Pome Fruit (crop group 11-10), Small Fruit Vine Climbing - except Fuzzy Kiwifruit (crop subgroup 13-07F), Stone Fruit (crop group 12-12), and Tree Nuts (crop group 14-12, except Almond).

- Treated areas may be replanted with any crop specified on this label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.
- **DO NOT** plant or transplant any crop not listed on this label within 12 months after the last application.

SPECIFIC CROP DIRECTIONS

Apply specified dosage of [this product/ PLENEXOS SMART] early in the infestation as the population begins to develop or at early threshold for the target insect pest. Apply higher dosages specified within the crop specific sections when applied as a preventive application, for moderate to heavy insect pressure, or where longer residual control is desired. Degree of efficacy against labeled pests will be determined, in part, by the stage of pest development at application and infestation level of those pests.

[This product/ PLENEXOS SMART] must be tank-mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations by crop. However, the use of Induce® adjuvant in combination with [this product/ PLENEXOS SMART] on grape, pome fruits, and stone fruits is prohibited when fruit is present due to adverse plant compatibility on harvested commodities. The tank-mixture of [this product/ PLENEXOS SMART] with an adjuvant/ additive having sticking properties or crop protection product formulations containing built-in stickers have been shown to interfere with leaf uptake and should be avoided. Sufficient leaf tissue must be present for uptake and translocation of this product.

CITRUS FRUIT (Group 10-10)^[*]**FOLIAR**

Australian lime (desert, finger, and round); Brown River finger lime; Calamondin; Citron; Citrus hybrids; Grapefruit; Japanese summer grapefruit; Kumquat; Lemon; Lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; Orange (sour and sweet); Pummelo; Russell River lime; Satsuma mandarin; Sweet lime; Tachibana orange; Tahiti lime; Tangelo; Tangerine (Mandarin); Tangor; Trifoliate orange; Uniq fruit; cultivars, varieties, and/or hybrids of these

Pests Controlled	Product Rate	
	(oz/A)	(lb ai/A)
Aphids ^[*] Asian citrus psyllid ^[*] California red scale ^[*] Citricola scale ^[*] Citrus leafminer ^[*] Citrus bud mite ^[*] Citrus red mite ^[*] Citrus rust mite (silver mite) ^[*] Citrus snow scale ^[*] Citrus thrips ^[*] Florida red scale ^[*] Mealybugs ^[*] Pink citrus rust mite ^[*] Purple scale ^[*] Texas citrus mite ^[*] Whiteflies ^[*]	9.5 - 19.0	0.043 - 0.086 (Spidoxamat) 0.089 - 0.18 (Flupyradifurone)
Pests Suppressed		
Barnacle scale ^{1[*]} Black scale ^[*] Brown scale ^[*] Cottony cushion scale ^[*] Nematodes ^[*]		

Application Instructions:

For production areas in [AL, FL, GA, LA, MS, SC, and TX,] optimum control of target pests is obtained at application volumes up to 250 GPA; application volumes in excess of 350 GPA must be avoided.

¹Time PLENEXOS SMART applications for control of barnacle scale according to crawler stage. Two applications may be required to achieve best efficacy.

Application Restrictions:

Pre-Harvest Interval (PHI): **1 day**

Minimum interval between applications: **10 days**

Minimum Application Volume: **30 or 50 Gallons/Acre** (Ground, See **Application Instructions**); **10 Gallons/Acre** (Aerial)

Maximum [PLENEXOS SMART/ product] single application rate: **19 oz/A** (spidoxamat 0.086 lb ai/A and 96 g ai/ha; flupyradifurone 0.18 lb ai/A)

Maximum number of [PLENEXOS SMART/ product] applications per calendar year: **2**

Maximum [PLENEXOS SMART/ product] allowed per calendar year: **38 oz/A** (spidoxamat 0.172 lb ai/A and 192 g ai/ha; flupyradifurone 0.36 lb ai/A)

DO NOT apply non-ionic surfactants in tank mix combination with [this product/ PLENEXOS SMART] on white grapefruit when fruit is present.

Directions for using soil and foliar application in an IPM program: The combination of one (1) soil application and one (1) foliar application of spidoxamat or flupyradifurone containing products with 14-days minimal interval between applications is allowed per calendar year: **DO NOT** apply more than a total of **0.4 lb ai/A** (444 g ai/ha) spidoxamat or **0.365 lb ai/A** (### g ai/ha) flupyradifurone per calendar year, regardless of formulation or method of application. Always follow label directions.

[*Not for use in CA [without a supplemental label].]

POME FRUIT (Group 11-10) ^[*] FOLIAR Apple; Asian pear; Azarole; Chinese quince; Crabapple; Japanese quince; Loquat; Mayhaw; Medlar; Pear; <i>Pseudocydonia sinensis</i> (Thouin) C.K. Schneid.; Quince; Tejocote; cultivars, varieties, and/or hybrids of these		
Pests Controlled	Product Rate	
Aphids (including Woolly Apple Aphid) ^[*] Apple rust mite ^[*] Leafhoppers ^[*] Mealybugs ^[*] Pear psylla ^[*] Pear rust mite ^[*] San Jose scale ¹ ^[*] Whiteflies ^[*]	(oz/A)	(lb ai/A)
Pests Suppressed	9.5 - 19.0	0.043 - 0.086 (Spidoxamat) 0.089 - 0.18 (Flupyradifurone)
Apple gall midge ^[*] Codling moth ^[*] European red mite ^[*] Micro-lepidoptera leafminers ^[*] Pear leaf midge ^[*] Twospotted spider mite ^[*] White apple leafhopper ^[*]		
Application Information: ¹ For control of San Jose scale west of the Rocky Mountains [including all of MT, WY, CO, and NM], apply at crawler stage followed by a second application 14 – 21 days later. For control of San Jose scale east of the Rocky Mountains; under heavy infestation pressure or where difficult control conditions exist a second application may be necessary.		
Application Restrictions: Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 10 days Minimum Application Volume: 30 or 50 Gallons/Acre (Ground, See Application Instructions); 10 Gallons/Acre (Aerial) Maximum [PLENEXOS SMART/ product] single application rate: 19 oz/A (spidoxamat 0.086 lb ai/A and 96 g ai/ha; flupyradifurone 0.18 lb ai/A) Maximum number of [PLENEXOS SMART/ product] applications per calendar year: 2 Maximum [PLENEXOS SMART/ product] allowed per calendar year: 38 oz/A (spidoxamat 0.172 lb ai/A and 192 g ai/ha; flupyradifurone 0.36 lb ai/A) [This product contains flupyradifurone, DO NOT apply [this product/ PLENEXOS SMART] with horticultural oils to pear in late-season or during periods of above average temperatures.] Directions for using soil and foliar application in an IPM program: The combination of one (1) soil application and one (1) foliar application is allowed per calendar year: DO NOT apply more than a total of 0.276 lb ai/A (308 g ai/ha) spidoxamat or 0.365 lb ai/A (### g ai/ha) flupyradifurone per calendar year, regardless of formulation or method of application. Always follow label directions.		
[*Not for use in CA [without a supplemental label].]		

SMALL FRUIT VINE CLIMBING - Except Fuzzy Kiwifruit (Subgroup 13-07F) ^[*]		
FOLIAR		
Amur river grape; Gooseberry; Grape; Kiwifruit (hardy); Maypop; Schisandra berry; cultivars, varieties, and/or hybrids of these		
Pests Controlled	Product Rate	
Grape tumid gallmaker ^[*] Mealybugs ^[*] Leafhoppers ^[*] Phylloxera ^[*] Sharpshooters ^[*] Willamette mite ^[*] Whiteflies ^[*]	(oz/A)	(lb ai/A)
	9.5 - 19.0	0.043 - 0.086 (Spidoxamat) 0.089 - 0.18 (Flupyradifurone)
Pests Suppressed		
European fruit lecanium scale ^[*] Pacific mite ^[*] Twospotted spider mite ^[*] Nematodes ^[*]		
Application Information: Some adjuvants that may be used with PLENEXOS SMART have caused intolerable damage to grape berries/ clusters when applied alone or in mixes after the initiation of bloom. Testing has shown that PLENEXOS SMART does not increase the potential for damage when used in combination with such adjuvants; however, not all adjuvants have been tested. It is recommended that a high quality spreading-penetrating adjuvant be used at a rate that is known to be safe to the crop.		
Application Restrictions: Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Minimum Application Volume: 30 or 50 Gallons/Acre (Ground, See Application Instructions); 10 Gallons/Acre (Aerial) Maximum [PLENEXOS SMART/ product] single application rate: 19 oz/A (spidoxamat 0.086 lb ai/A and 96 g ai/ha; flupyradifurone 0.18 lb ai/A) Maximum number of [PLENEXOS SMART/ product] applications per calendar year: 2 Maximum [PLENEXOS SMART/ product] allowed per calendar year: 38 oz/A (spidoxamat 0.172 lb ai/A and 192 g ai/ha; flupyradifurone 0.36 lb ai/A) Directions for using soil and foliar application in an IPM program: The combination of one (1) soil application and one (1) foliar application is allowed per calendar year: DO NOT apply more than a total of 0.276 lb ai/A (308 g ai/ha) spidoxamat or 0.365 lb ai/A (### g ai/ha) flupyradifurone per calendar year, regardless of formulation or method of application. Always follow label directions.		
[*Not for use in CA [without a supplemental label].]		

STONE FRUIT (Group 12-12)^[*] FOLIAR Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these		
Pests Controlled		Product Rate
Aphids ^[*] Mealybugs ^[*] San Jose scale ^{1[*]} White peach scale ^[*] Whiteflies ^[*]	(oz/A)	(lb ai/A)
	9.5 - 19.0	0.043 - 0.086 (Spidoxamat) 0.089 - 0.18 (Flupyradifurone)
Pests Suppressed		
Black scale ^[*] Cherry fruit fly ^[*] European fruit lecanium scale ^[*] European red mite ^[*] Nematodes ^[*] Spotted Wing Drosophila ^[*] Twospotted spider mite ^[*]		
Application Information: ¹ For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.		
Application Restrictions: Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 10 days Minimum Application Volume: 30 or 50 Gallons/Acre (Ground, See Application Instructions); 10 Gallons/Acre (Aerial) Maximum [PLENEXOS SMART/ product] single application rate: 19 oz/A (spidoxamat 0.086 lb ai/A and 96 g ai/ha; flupyradifurone 0.18 lb ai/A) Maximum number of [PLENEXOS SMART/ product] applications per calendar year: 2 Maximum [PLENEXOS SMART/ product] allowed per calendar year: 38 oz/A (spidoxamat 0.172 lb ai/A and 192 g ai/ha; flupyradifurone 0.36 lb ai/A) DO NOT apply until after petal fall. Directions for using soil and foliar application in an IPM program: The combination of one (1) soil application and one (1) foliar application is allowed per calendar year: DO NOT apply more than a total of 0.276 lb ai/A (308 g ai/ha) spidoxamat or 0.365 lb ai/A (### g ai/ha) flupyradifurone per calendar year, regardless of formulation or method of application. Always follow label directions. [*Not for use in CA [without a supplemental label].]		

TREE NUTS (Group 14-12, except Almond)^[*] FOLIAR African nut-tree; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these		
Pests Controlled	Product Rate	
Aphids ^[*] Mealybugs ^[*] Phylloxera ^[*] San Jose scale ^{1[*]} Walnut scale ^[*] Whiteflies ^[*]	(oz/A)	(lb ai/A)
Pests Suppressed European fruit lecanium scale ^[*] Nematodes ^[*] Olive scale ^[*] Pacific mite ^[*] Twospotted spider mite ^[*]	9.5 - 19.0	0.043 - 0.086 (Spidoxamat) 0.089- 0.18 (Flupyradifurone)
Application Information: ¹ For control of San Jose scale under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.		
Application Restrictions: Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 10 days Minimum Application Volume: 30 or 50 Gallons/Acre (Ground, See Application Instructions); 10 Gallons/Acre (Aerial) Maximum [PLENEXOS SMART/ product] single application rate: 19 oz/A (spidoxamat 0.086 lb ai/A and 96 g ai/ha; flupyradifurone 0.18 lb ai/A) Maximum number of [PLENEXOS SMART/ product] applications per calendar year: 2 Maximum [PLENEXOS SMART/ product] allowed per calendar year: 38 oz/A (spidoxamat 0.172 lb ai/A and 192 g ai/ha; flupyradifurone 0.36 lb ai/A) Directions for using soil and foliar application in an IPM program: The combination of one (1) soil application and one (1) foliar application is allowed per calendar year: DO NOT apply more than a total of 0.276 lb ai/A (308 g ai/ha) spidoxamat or 0.365 lb ai/A (### g ai/ha) flupyradifurone per calendar year, regardless of formulation or method of application. Always follow label directions. [*]Not for use in CA [without a supplemental label].]		

STORAGE AND DISPOSAL

DO NOT contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE

Store in original container and keep tightly closed when not in use. Store in a cool dry place. **DO NOT** store in direct sunlight. Handle and open container in a manner as to prevent spillage. Avoid cross-contamination with other pesticides.

PESTICIDE DISPOSAL

[Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.]

[Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.]

[To avoid wastes, use all material in this container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.]

CONTAINER HANDLING

[Non-refillable Containers]

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Rigid Non-refillable containers that are too Large to Shake (i.e. with capacities greater than 5 gallons or 50 lbs)

Non-refillable container. **DO NOT** reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

- **Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

- **Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)**

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Non-Refillable Fiber Drums with Liners

Non-refillable container. **DO NOT** reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration. If drum is contaminated and cannot be reused, offer for recycling if available or dispose of in a sanitary landfill or by incineration.

Non-Rigid, Non-refillable Containers

Non-refillable container. **DO NOT** reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

[Refillable Containers]

Refillable container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Contact your Ag retailer or Bayer CropScience for container return, disposal and recycling information.

- **Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

- **Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)**

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container.

PLENEXOS and the Bayer Cross logo are registered trademarks of Bayer Group.

INDUCE is a registered trademark of Helena Holding Company

PLENEXOS™ SMART (PENDING) 09/20/2022, 01/31/2023, 07/18/2023, 12/18/2023, 12/19/2023, 02/02/2024, 04/01/2024, 02/17/2026, 02/20/2026