

**Karan Vora (kv2154)**  
**Machine Learning Assignment 4**

**Problem 1):**

Compute the mean of the image data of shape (100, 1900). And then subtract the mean from all data points to get the de-centered data.

$$\mu = \text{mean}(X)$$

and get x from

$$x = X - \mu$$

Computing the covariance matrix


$$C = x^T x$$

Apply eigenvector Decomposition to covariance matrix


$$C = V \Lambda V^{-1}$$

The 3 most significant eigen value is


Eigen Value = 4.215022



Eigen Value = 3.016759



Eigen Value = 2.099301



The coefficient matrix is obtained by

$$c_{ij} = (X_i - \mu)^T V'_j = x_i^T V'_j$$

This is just the inner product of  $x$  and the top 3 eigenvectors

$$c = \mathbf{xV'}$$

we can calculate  $\hat{X}$  by

$$\hat{X} = \mu + \sum_{j=1}^c c_{ij} V'_j$$

which in simple terms is


$$\hat{X} = \mu + cV'$$

The 10 generated images are


Original image




PCA image




Original image




PCA image




Original image




PCA image




Original image




PCA image




Original image




PCA image




Original image




PCA image




Original image




PCA image







Original image



PCA image





=====

**Problem 2):**

First box 8 Apples 4 Oranges

Second box 10 Apples 2 Oranges

Probability of selecting box =  $P(E_1) = P(E_2) = \frac{1}{2}$

Select Apple from First box

$$P(A/E_1) = \frac{8}{12} = \frac{2}{3}$$

Select Apple from Second box



$$P(A/E_2) = \frac{10}{12} = \frac{5}{6}$$

Applying Bayes Theorem

$$P(E_1/A) = \frac{P(E_1)P(A/E_1)}{P(E_1)P(A/E_1) + P(E_2)P(A/E_2)}$$

$$= \frac{\frac{1}{2} \times \frac{2}{3}}{(\frac{1}{2} \times \frac{2}{3}) + (\frac{1}{2} \times \frac{5}{6})}$$

$$P(E_1/A) = \frac{4}{9}$$

=====

**Problem 3):**

②

parameter,  $\Theta = \{\alpha, \mu_1, \Sigma_1, \mu_2, \Sigma_2\}$

$$P(y|\Theta) = \alpha^y (1-\alpha)^{1-y}$$

$$P(x, y|\Theta) = \mu(x|\mu_y, \Sigma_y)$$

Using the parameter  $\Theta$ , we can generate the IID data from :-

$$P(x|y, \Theta) = P(y|\Theta) P(x|y, \Theta)$$

We will use maximum likelihood, to learn the parameter from data

$$L(\Theta) = \log P(\text{Data}|\Theta)$$

$$= \sum_{i=1}^N \log P(x_i, y_i|\Theta)$$

According to the question there are two classes:-

Class 1:-  $y_i \in 0$

Class 2:-  $y_i \in 1$

$$\therefore L(\Theta) = \sum_{i=1}^N \log (P(y_i|\Theta) P(x_i|y_i, \Theta))$$

$$= \sum_{i=1}^N \log (P(y_i|\Theta)) + \sum_{i=1}^N \log (P(x_i|y_i, \Theta))$$

(2)

$$= \sum_{i=1}^N \log p(y_i | \alpha) + \sum_{y \in 0} \log p(x_i | \mu_0, \Sigma_0) + \sum_{y \in 1} \log p(x_i | \mu_1, \Sigma_1)$$

Estimating the likelihood ~~for the given~~

$$p(y_i | \alpha) = \alpha^{y_i} (1-\alpha)^{1-y_i} \quad \text{--- (1)}$$

$$p(x_i | \mu_0, \Sigma_0) = \frac{1}{2\pi^{D/2} (\sqrt{|\Sigma_0|})} \exp \left( -\frac{1}{2} (\vec{x} - \vec{\mu}_0)^T \Sigma_0^{-1} (\vec{x} - \vec{\mu}_0) \right)$$

$$p(x_i | \mu_1, \Sigma_1) = \frac{1}{2\pi^{D/2} (\sqrt{|\Sigma_1|})} \exp \left( -\frac{1}{2} (\vec{x} - \vec{\mu}_1)^T \Sigma_1^{-1} (\vec{x} - \vec{\mu}_1) \right)$$

Diff. eqn wrt  $\alpha$

$$\frac{\partial}{\partial \alpha} L(\alpha) = \frac{\partial}{\partial \alpha} \sum_{i=1}^N \log (\alpha^{y_i} (1-\alpha)^{1-y_i}) + \frac{\partial}{\partial \alpha} \log \pi = 0$$

$$\frac{\partial}{\partial \alpha} \sum_{i=1}^N (y_i \log \alpha + (1-y_i) \log (1-\alpha)) = 0$$

$$\frac{d}{d\alpha} \left( \sum_{i \in \text{class 1}} \log \alpha + \sum_{i \in \text{class 0}} \log (1-\alpha) \right) = 0$$

$$\sum_{i \in \text{class 0}} \frac{1}{\alpha} - \sum_{i \in \text{class 1}} \frac{1}{1-\alpha} = 0$$

$$\frac{N_1}{\alpha} - \frac{N_0}{1-\alpha} = 0$$

$$\frac{N_1}{\alpha} = \frac{N_0}{1-\alpha}$$

$$\alpha = \frac{N_1}{N_0 + N_1}$$

Differentiating the log-likelihood to  $\mu_0$

$$\frac{\partial(L(\theta))}{\partial \mu_0} = \frac{\partial}{\partial \mu_0} \left( \sum_{i \in \mathcal{O}} \log \left( \frac{1}{(2\pi)^{D/2} \sqrt{|\Sigma_0|}} \right) \right) =$$

$$\exp \left( -\frac{1}{2} (\vec{x}_i - \vec{\mu}_0)^T \Sigma_0^{-1} (\vec{x}_i - \vec{\mu}_0) \right)$$

$$= \frac{\partial}{\partial \mu_0} \left( \sum_{i \in \mathcal{O}} \left( -\frac{D}{2} \log(2\pi) - \frac{1}{2} \log(|\Sigma_0|) - \frac{1}{2} (\vec{x}_i - \vec{\mu}_0)^T \Sigma_0^{-1} (\vec{x}_i - \vec{\mu}_0) \right) \right)$$

(4)

$$\frac{\partial}{\partial \mu_0} \sum_{y_i \in 0} \left[ -\frac{1}{2} (\vec{x}_i - \vec{\mu}_0)^T \Sigma_0^{-1} (\vec{x}_i - \vec{\mu}_0) \right] = 0$$

$$\sum_{y_i \in 0} (\vec{x}_i - \vec{\mu}_0)^T \Sigma_0^{-1} = 0$$

$$\Sigma_0^{-1} \neq 0,$$

$$\sum_{y_i \in 0} (\vec{x}_i - \vec{\mu}_0)^T = 0$$

$$\sum_{y_i \in 0} \vec{x}_i - N_0 \vec{\mu}_0 = 0$$

$$\vec{\mu}_0 = \frac{\sum_{y_i \in 0} \vec{x}_i}{N_0}$$

Similarly,

$$\vec{\mu}_1 = \frac{\sum_{y_i \in 1} \vec{x}_i}{N_1}$$



Now, diff. wrt  $\Sigma_0^{-1}$

$$\frac{\partial (L(\theta))}{\partial \Sigma_0^{-1}} = \frac{\partial}{\partial \Sigma_0^{-1}} \sum_{y_i \in G} \log P(x_i | \mu_0, \Sigma_0) \rightarrow$$

$$\frac{\partial}{\partial \Sigma_0^{-1}} (\text{const})$$

$$= \frac{\partial}{\partial \Sigma_0^{-1}} \sum_{y_i \in G} \log P(x_i | \mu_0, \Sigma_0)$$

$$= \frac{\partial}{\partial \Sigma_0^{-1}} \left( \sum_{y_i \in G} \log \left( \frac{1}{(2\pi)^{D/2} \sqrt{|\Sigma_0|}} \exp \left( -\frac{1}{2} (\vec{x}_i - \vec{\mu}_0)^T \Sigma_0^{-1} (\vec{x}_i - \vec{\mu}_0) \right) \right) \right)$$

$$= \frac{\partial}{\partial \Sigma_0^{-1}} \left[ \frac{N_0}{2} \log |\Sigma_0^{-1}| - \frac{1}{2} \sum_{y_i \in G} \frac{1}{2} (\vec{x}_i - \vec{\mu}_0)^T \Sigma_0^{-1} (\vec{x}_i - \vec{\mu}_0) \right]$$

$$= \frac{\partial}{\partial \Sigma_0^{-1}} \left[ \frac{N_0}{2} \log |\Sigma_0^{-1}| - \frac{1}{2} \sum_{y_i \in G} \text{tr} \left[ (\vec{x}_i - \vec{\mu}_0)^T \Sigma_0^{-1} (\vec{x}_i - \vec{\mu}_0) \right] \right]$$

Now,  $\Sigma_0^{-1} = A$

Therefore we will differentiate w.r.t  $A$

$$\frac{d}{dA} \left( \frac{N_0}{2} \log |A| - \frac{1}{2} \sum_{y_i \in \mathcal{O}} \text{tr} \left[ (\vec{x}_i - \vec{\mu}_0)^T A (\vec{x}_i - \vec{\mu}_0) \right] \right) = 0$$

$$\frac{N_0}{2} (A^{-1})^T - \frac{1}{2} \sum_{y_i \in \mathcal{O}} \left[ (\vec{x}_i - \vec{\mu}_0) (\vec{x}_i - \vec{\mu}_0)^T \right]^T = 0$$

$$\frac{N_0}{2} A^{-1} - \frac{1}{2} \sum_{y_i \in \mathcal{O}} (\vec{x}_i - \vec{\mu}_0) (\vec{x}_i - \vec{\mu}_0)^T = 0$$

~~Therefore when differentiating w.r.t~~

$$\frac{N_0}{2} \Sigma_0 - \frac{1}{2} \sum_{y_i \in \mathcal{O}} (\vec{x}_i - \vec{\mu}_0) (\vec{x}_i - \vec{\mu}_0)^T = 0$$

$$N_0 (\Sigma_0) = \sum_{y_i \in \mathcal{O}} (\vec{x}_i - \vec{\mu}_0) (\vec{x}_i - \vec{\mu}_0)^T$$

$$\Sigma_0 = \frac{1}{N_0} \sum_{y_i \in \mathcal{O}} (\vec{x}_i - \vec{\mu}_0) (\vec{x}_i - \vec{\mu}_0)^T$$

Similarly,

$$\Sigma_1 = \frac{1}{N_1} \sum_{y_i \in 1} (\vec{x}_i - \vec{\mu}_1)(\vec{x}_i - \vec{\mu}_1)^T$$

Now, The given Bayes optimal decision is

$$y = \operatorname{argmax}_{y \in \{0,1\}} P(\hat{y}|x)$$

For a linear decision Boundary of 2 class probth

$$P(y=1|x) = P(y=0|x) = 0.5$$

Using the conditional parameters we get :-

$$P(y=1|x) = \frac{P(x, y=1)}{P(x)}$$

$$= \frac{P(x, y=1) + P(y=1)}{\sum_y P(x, y)}$$

$$= \frac{P(x|y=1)P(y=1)}{P(x, y=1) + P(x, y=0)}$$



(8)

$$= p(x|y=1) p(y=1)$$

$$\frac{p(x|y=1) p(y=1)}{p(x|y=1) p(y=1) + p(x|y=0) p(y=0)}$$

$$p(y=1) = \alpha^1 (1-\alpha)^0$$

$$= \alpha$$

$$p(y=0) = \alpha^0 (1-\alpha)^{1-0}$$

$$= 1-\alpha$$

$$\therefore p(x|y=1) = N(x|\mu_1, \Sigma_1)$$

$$p(y=1|x) = \frac{\alpha N(x|\mu_1, \Sigma_1)}{\alpha N(x|\mu_1, \Sigma_1) + (1-\alpha) N(x|\mu_0, \Sigma_0)}$$

$$p(y=0|x) = \frac{p(x|y=0) \cdot p(y=0)}{p(x|y=0) p(y=1) + p(x|y=0) p(y=0)}$$

$$= \frac{(1-\alpha) N(x|\mu_0, \Sigma_0)}{\alpha N(x|\mu_1, \Sigma_1) + (1-\alpha) N(x|\mu_0, \Sigma_0)}$$

$\therefore$  To calculate decision boundary

$$p(y=1|x) = p(y=0|x)$$

$$+ (1-\alpha) \sum_{i=1}^N (1+\alpha) \frac{1}{2} + \sum_{i=1}^N (1-\alpha) \frac{1}{2}$$

(1)

$$(1-\alpha) N(x|M_0, \Sigma_0) = \alpha (N(x|M_1, \Sigma_1)) \quad (1)$$

Sub. N we get

$$\frac{1}{(2\pi)^{D/2} \sqrt{|\Sigma|}} \exp\left(-\frac{1}{2}(\vec{x}-\vec{M}_1)^T \Sigma_1^{-1} (\vec{x}-\vec{M}_1)\right) =$$

$$\frac{(1-\alpha)}{(2\pi)^{D/2} \sqrt{|\Sigma|}} \exp\left(-\frac{1}{2}(\vec{x}-\vec{M}_0)^T \Sigma_0^{-1} (\vec{x}-\vec{M}_0)\right)$$

~~we~~ simplifying

$$\begin{aligned} & -\frac{1}{2}(\vec{x}-\vec{M}_1)^T \Sigma_1^{-1} (\vec{x}-\vec{M}_1) + \frac{1}{2}(\vec{x}-\vec{M}_0)^T \Sigma_0^{-1} (\vec{x}-\vec{M}_0) \\ & = \log\left(\frac{1-\alpha}{\alpha} \sqrt{\frac{|\Sigma_1|}{|\Sigma_0|}}\right) \end{aligned}$$

Now, for linear classification

$$\Sigma_0, \Sigma_1 = \text{covariance} \quad \alpha = \frac{N_1}{N_0 + N_1}$$

$$\begin{aligned} & -\frac{1}{2}(\vec{x}-\vec{M}_1)^T \Sigma_1^{-1} (\vec{x}-\vec{M}_1) + \frac{1}{2}(\vec{x}-\vec{M}_0)^T \Sigma_0^{-1} (\vec{x}-\vec{M}_0) \\ & = \log\left(\frac{N_0}{N_1}\right) \end{aligned}$$

(10)

$$\Sigma^{-1} (\mu_1 - \mu_0)^T \vec{x} + \frac{1}{2} (\mu_0 + \mu_1)^T \Sigma^{-1} (\mu_0 - \mu_1) +$$

$$\log\left(\frac{N_1}{N_0}\right) = 0$$

$$w = \Sigma (\mu_1 - \mu_0)$$

$$b = \frac{1}{2} (\mu_0 + \mu_1)^T \Sigma^{-1} (\mu_0 + \mu_1) - \log \frac{N_1}{N_0}$$

$$f(x) = \text{sign}(w^T x + b)$$

$$\text{FF}, \Sigma_0 \neq \Sigma_1$$

$$-\frac{1}{2} (\vec{x} - \mu_1)^T \Sigma_1^{-1} (\vec{x} - \mu_1) +$$

$$\frac{1}{2} (\vec{x} - \mu_0)^T \Sigma_0^{-1} (\vec{x} - \mu_0) + \frac{1}{2} \log \frac{\Sigma_0}{\Sigma_1} +$$

$$\log\left(\frac{N_1}{N_0}\right) = 0$$