

18.06 FINAL EXAM. IN 1993.
TO POST

Part I

1. Find all possible values for the determinant of the given type of 3×3 real matrix.

[2] (a) A matrix with independent columns.

[2] (b) A matrix with $A^2 = A$.

[2] (c) A matrix with pivots 1, 2 and 3.

[2] (d) A ~~Markov matrix.~~

[3] 2. Why is there no orthogonal matrix Q such that $Q^{-1}AQ = \Lambda$ if

$$A = \begin{bmatrix} 1 & 4 & 5 \\ 0 & 2 & 6 \\ 0 & 0 & 3 \end{bmatrix}, \quad \Lambda = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}?$$

[2] 3. The left null space $\mathcal{N}(A^T)$ of a matrix A is spanned by $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}$. The set of solutions of

the equation $Ax = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}$ is (circle one)

the empty set, a point, a line, a plane, a three-dimensional hyperplane in \mathbb{R}^4 , all of \mathbb{R}^4 .

- [4] 4. Apply the Gram-Schmidt algorithm to the columns of the matrix $\begin{bmatrix} 0 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$ to obtain three orthonormal vectors.

- [3] 5. For what values of a and b is the quadratic form $ax^2 + 2xy + by^2 = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} a & 1 \\ 1 & b \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$ positive definite?

6. Suppose A is an m by n matrix, with independent columns.

- [2] • What can you deduce about the relation of m and n ?
- [2] • What can you deduce about the set of solutions to $Ax = 0$?
- [2] • For which m and n are there nonzero solutions to $A^T y = 0$?
- [2] • Give two properties of the matrix $A^T A$ (other than the fact that it is square).

[2] 7. Give an example of a matrix with exactly two zero eigenvalues and no zero entries.

- [3] 8. Consider the square matrix $A = \begin{bmatrix} 7 & -3 \\ 9 & -5 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}^{-1}$. Find the solution to the differential equation $\frac{du(t)}{dt} = Au(t)$ with initial condition $u(0) = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

- [3] 9. Find a basis for the orthogonal complement of the subspace of \mathbb{R}^4 spanned by the vectors $\begin{bmatrix} 2 \\ 0 \\ 1 \\ 2 \end{bmatrix}$, $\begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}$, and $\begin{bmatrix} 3 \\ 0 \\ 1 \\ 3 \end{bmatrix}$.

10. Are the following statements true or false? You get 2 points for a correct answer and -2 points for an incorrect answer.

T F (a) If $M^{-1}AM = B$, then A and B must have the same eigenvectors.

T F (b) The matrices A and A^T always have the same rank.

T F (c) There is a matrix with column space is spanned by the vectors $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 4 \end{bmatrix}$, and with row space spanned by vectors $\begin{bmatrix} 1 \\ 4 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$.

T F (d) If a square matrix has a repeated eigenvalue, it cannot be diagonalizable.

T F (e) The set of vectors in \mathbb{R}^3 with integer (whole number) components is a subspace of \mathbb{R}^3 .

Part II

1. Let A be the matrix $\begin{bmatrix} 1 & 2 & 1 & 2 \\ 2 & 1 & 2 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$.

- [3] (a) Find a factorization $A = LU$, where L is a lower triangular matrix, and U is in echelon form.

- [4] (b) Find the general solution of $Ax = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}$.

- [4] 1. (c) The vector $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$ is in the column space of A if a , b and c satisfy what linear conditions?

- [4] 2. The vector space \mathbb{R}^2 has bases $\left\{ \mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\left\{ \mathbf{w}_1 = \begin{bmatrix} 3 \\ 2 \end{bmatrix}, \mathbf{w}_2 = \begin{bmatrix} 4 \\ 3 \end{bmatrix} \right\}$. Write the matrix for the identity linear transformation I from the basis $\{\mathbf{v}_1, \mathbf{v}_2\}$ to the basis $\{\mathbf{w}_1, \mathbf{w}_2\}$.

3. Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 2 \end{bmatrix}$.

- [3] (a) Find a matrix Q with orthonormal columns and an upper triangular matrix R such that $A = QR$.

- [3] (b) Find the closest vector to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ in the column space of A .

- [3] (c) If \mathbf{v} and \mathbf{w} are any two linearly independent vectors, find a nonzero linear combination that is perpendicular to \mathbf{v} .

- [3] (d) Compute the matrix P which projects onto the column space of A .

4. Let \mathbf{u} and \mathbf{v} be vectors in the Euclidean space \mathbb{R}^n , and let A be the square matrix $\mathbf{u}\mathbf{v}^T$.

[3] (a) Describe the row space and nullspace of A in terms of \mathbf{u} and \mathbf{v} .

[2] (b) Show that \mathbf{u} is an eigenvector of A , and find the corresponding eigenvalue.

[2] (c) What condition must be satisfied by \mathbf{u} and \mathbf{v} for A to be skew-symmetric ($A = -A^T$)?

[2] (d) What condition must be satisfied by \mathbf{u} and \mathbf{v} so that $A^2 = A$?

5. Let A_n be the $n \times n$ matrix with entries $a_{ij} = \begin{cases} -2, & i = j - 1, \\ 1, & i = j, \\ 1, & i = j + 1, \\ 0, & \text{other entries.} \end{cases}$ For example,

$$A_1 = [1], \quad A_2 = \begin{bmatrix} 1 & -2 \\ 1 & 1 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 1 & -2 & 0 \\ 1 & 1 & -2 \\ 0 & 1 & 1 \end{bmatrix}, \quad A_4 = \begin{bmatrix} 1 & -2 & 0 & 0 \\ 1 & 1 & -2 & 0 \\ 0 & 1 & 1 & -2 \\ 0 & 0 & 1 & 1 \end{bmatrix}.$$

- [4] (a) Let $d_n = \det(A_n)$. Find numbers a and b such that for $n = 3, 4, 5, \dots$,

$$d_n = ad_{n-1} + bd_{n-2}.$$

- [1] (b) What is d_4 ?

- [4] (c) Write the matrix A such that $\begin{bmatrix} d_{n+1} \\ d_n \end{bmatrix} = A \begin{bmatrix} d_n \\ d_{n-1} \end{bmatrix}$, and calculate its eigenvalues and eigenvectors.

[2] 5. (d) Find the number λ such that d_n/λ^n tends to a non-zero, finite limit as n tends to infinity.

[5] 6. (a) If a, b, c are real numbers, find the eigenvalues and null space of the skew-symmetric matrix $A = \begin{bmatrix} 0 & c & -b \\ -c & 0 & a \\ b & -a & 0 \end{bmatrix}$.

[2] (b) Explain why the matrix A in part (a) of this problem cannot be orthogonal.