

3'end RNA-seq using template switching protocol

Status: [?] | partially migrated / unconfirmed

Status legend: [OK] working [?] unconfirmed / partial [X] broken

About

Protocol for 3' end RNA-seq using a template switching reverse transcription workflow.

Contents

1. RNA extraction
2. RNA fragmentation and template switching RT
3. Template switching RT incubation and cleanup
4. Optional PCR test
5. Indexing PCR
6. PCR beads purification
7. Buffers
8. Migration notes

1. RNA extraction

Input / starting material: RNA lysate

- Maxwell RNA extraction
- Alternative: Add 3 volumes of LS-Trizol to lysate
- Extract RNA using the Zymo MicroSpin Columns
- Elute with 6 μ L of H₂O

2. RNA fragmentation and template switching RT

- Prepare fragmentation buffer.
- Fragmentation buffer: 80 mM Tris-HCl pH 7.5, 40 mM MgCl₂.

Note: have more recently been using milder fragmentation buffer conditions with input concentration of 10 mM Tris-HCl pH 7.5, 5 mM MgCl₂ (final concentration 1.25 mM Tris-HCl and 0.625 mM MgCl₂).

- Use 500 ng of RNA (whenever possible).

- Bring to 3.5 μL using water.
 - Add 0.5 μL of fragmentation buffer (10 mM Tris-HCl pH 7.5, 5 mM MgCl final concentration).
 - Fragment RNA by incubating 12 min at 95 $^{\circ}\text{C}$.
 - Add:
 - 0.5 μL 10 mM dNTPs
 - 0.5 μL 5 μM primer RT1
 - Mix and incubate at 65 $^{\circ}\text{C}$ for 3 min then cool down to 42 $^{\circ}\text{C}$ and hold.
-

3. Template switching RT incubation and cleanup

- Make a mastermix and add 5.5 μL of the following:
 - 2 μL of SSIV buffer
 - 2 μL 5 M betaine
 - 0.5 μL 0.1 M DTT
 - 0.25 μL 40 μM TSO
 - 0.25 μL 200 mM MgCl
 - 0.25 μL SuperRNasin
 - 0.25 μL SSIV
 - Place the mastermix into a PCR strip and into the cycler so that it pre-warms to 42 $^{\circ}\text{C}$ before you start adding it to the samples.
 - Add mastermix to samples while keeping them at 42 $^{\circ}\text{C}$ in the cycler (i.e., don't take the samples out of the cycles while adding the mix).
 - Incubate at 42 $^{\circ}\text{C}$ for 10 min.
 - Incubate at 50 $^{\circ}\text{C}$ for 30 min.
 - Purify using Magbind beads (1.8 μL per 1 μL of PCR reaction). Elute in 8 μL H₂O.
-

4. Optional PCR test

- Use 1 μL RT product.
-

5. Indexing PCR

- Prepare the following 20 μL indexing PCR reaction for each sample:
 - 10 μL of 2 $\times$ Q5
 - 2 μL RT product
 - 1 μL dual indexed primers

- * Use a different index for each sample.
 - * Make sure each sample will have a different index if submitting multiple libraries at the same time.
 - 7 μL of ddH₂O
 - Incubate for the PCR reaction using the Q5 program:
 - Extension time – 45 seconds
 - T_m – 65 °C
 - Typically try 10 and 14 cycles
 - Move samples to post-PCR room.
 - Add 2 μL of purple 6X loading buffer to 5 μL of PCR reaction. Keep the rest of the 20 μL .
 - Load on a 6% TBE gel together with 2 μL of ready-to-use low molecular weight ladder (NEB).
 - Run at 180 V for 25 min.
 - Stain with Sybr Green (1:10000 dilution) in 1X TBE.
 - Visualise.
-

6. PCR beads purification

- Mix together approx. 10 μL of each PCR reaction.
 - Add a suitable amount of Magbind beads (e.g. 0.9 μL per 1 μL of PCR reaction) and mix.
 - Incubate at room temperature for 5 minutes.
 - Remove supernatant and wash with 500 μL of 70% Ethanol for 30 sec. Repeat this step.
 - Air dry the beads.
 - Resuspend in 30 μL of H₂O.
-

7. Buffers

- Fragmentation buffer: 80 mM Tris-HCl pH 7.5, 40 mM MgCl₂.
 - Milder fragmentation buffer conditions: input concentration of 10 mM Tris-HCl pH 7.5, 5 mM MgCl₂ (final concentration 1.25 mM Tris-HCl and 0.625 mM MgCl₂).
-