Characterizing galaxy-dark matter offsets in galaxy clusters of the Illustris simulation

Karen Y. Ng, ¹ Annalisa P. Pillepich, ² William A. Dawson, ³ D. Wittman, ¹ Lars Hernquist, ² etc. [order TBD]

arXiv

ABSTRACT

Key words: - methods: statistical

- 1 INTRODUCTION
- 2 DATA
- 2.1 Test data from Gaussian mixture(s)
- 2.2 Data from the Illustris simulation
- 2.2.1 Characterizing the non-relaxedness of the galaxy clusters / groups
- 2.3 Data proprocessing / selection
- 3 METHODS
- 3.1 Galaxy Centers
- $3.1.1 \quad Cross-validated \ Kernel \ Density \ Estimation \ (KDE) \\ and \ peak \ finder$
- 3.1.2 Shrinking aperture
- 3.1.3 Weighted and unweighted centroids
- 3.1.4 Brightest Cluster Galaxies (BCG)
- 3.2 DM Centers
- 3.3 Finding offsets
- 4 RESULTS
- 4.1 Benchmark test results from Gaussian mixtures
- 4.2 Galaxy-DM Offset in Illustris
- 5 DISCUSSION
- 5.1 Comparison to other simulations
- 5.2 Comparison to other observational studies
- 5.3 Galaxy-DM Offset in Merging Galaxy Clusters
- 6 ACKNOWLEDGEMENTS

APPENDIX A: KDE

2 Karen Y. Ng et al.

This paper has been typeset from a TEX/ IATEX file prepared by the author.