Galaxy-dark matter offsets in galaxy clusters / groups of the Illustris simulation

Karen Y. Ng,¹ Annalisa P. Pillepich,² William A. Dawson,³ D. Wittman,¹ Lars Hernquist,² etc. [order TBD]

arXiv

ABSTRACT

Key words: - methods: statistical

1 INTRODUCTION

- 2 DATA
- 2.1 Test data from Gaussian mixture(s)
- 2.2 Data from the Illustris simulation
- 2.2.1 Properties of the galaxy clusters / groups
- 2.2.2 Data proprocessing / selection
- 2.2.3 Data with and without noise
- 2.2.4 Volume Selection
- 2.2.5 Galaxy Selection
- 2.2.6 Galaxy weights

3 METHODS

3.1 Galaxy Centers

- 3.1.1 Cross-validated Kernel Density Estimation (KDE) and peak finder
- 3.1.2 Shrinking aperture
- 3.1.3 Weighted and unweighted centroids
- 3.1.4 Brightest Cluster Galaxies (BCG)
- 3.2 DM Centers
- 3.3 Finding offsets
- 4 RESULTS
- 4.1 Benchmark results from Gaussian mixtures
- 4.2 Galaxy-DM Offset in Illustris
- 5 DISCUSSION
- 5.1 Comparison to other simulations
- 5.2 Comparison to other observational studies
- 5.3 Galaxy-DM Offset in Merging Galaxy
 Clusters

6 ACKNOWLEDGEMENTS

APPENDIX A: KDE

2 Karen Y. Ng et al.

This paper has been typeset from a TeX/ IATeX file prepared by the author.