

# Algebraic and Topological Persistence

by  
Luciano Melodia



A bachelor thesis submitted in fulfillment  
of the requirements for the degree of  
Bachelor of Science (B.Sc.)  
in Mathematics

Friedrich-Alexander University  
Erlangen-Nürnberg  
May 21, 2024

# Algebraic and Topological Persistence

Luciano Melodia

## Abstract

This thesis addresses the theory of topological spaces and the theoretical foundations of persistence theory. We begin with a brief motivation from an applied context that inspired this work. Much of the research in persistence theory has been driven by the goal of utilizing persistent homology for data analysis. However, we restrict ourselves to providing an illustrative context and then proceed strictly to the theory of topological spaces. We discuss chain complexes (??) and the associated simplicial homology groups (??), as well as their connection to singular homology (??). Additionally, we introduce classical tools of algebraic topology, such as exact and short exact sequences (2.3.2), and relative homology groups obtained by quotienting with subspaces of a topological space (2.4). These tools are used to prove the Excision Theorem ?? in algebraic topology. We then use this theorem to demonstrate the equivalence of simplicial and singular homology for triangulable topological spaces, i.e., those topological spaces that admit a simplicial structure (2.6). This allows for the use of a more general theory of homology in the investigation of topological methods through theoretical considerations.

In the chapter on homological persistence, we consistently use these tools. We develop the theory of persistent homology, the homology of filtrations of topological spaces, and the corresponding dual concept of persistent cohomology. Finally, we generalize both theories into zigzag persistence. Zigzag modules allow for the treatment of maps in both directions along a filtration, making them highly useful for local applications of zigzag persistence. This work aims to provide mathematicians with a solid foundation for productively engaging with the aforementioned theories. Most of the proofs have been rewritten to clarify the relationships between the discussed theories. The novelty of this contribution lies in the comprehensive presentation of persistence theory and its ideas through rigorous mathematical treatment for triangulable topological spaces.

# Dedication

This thesis is dedicated with deep appreciation to those closest to me, whose support was indispensable in writing these pages and undertaking this journey.

My academic path took me through studies in German and Italian, then to a major in Information Science with a focus on Media Informatics, culminating in a Masters degree from Regensburg. A three-year doctoral stint in Computer Science at Erlangen followed, after which I shifted my focus to theoretical mathematics. Throughout, I grappled with whether I was truly satisfied with my choices, always drawn to the underlying structures of language and computing. It is with joy that I reflect on these discoveries.

I am grateful to my mother, Beata, who, despite initial reservations, supported my decisions wholeheartedly once they were made.

A special thanks to my late father, Domenico, who taught me to value the past more than the future.

Thanks also to my sister, Dominique, who never ceased to engage me in discussions about physics whenever she had the chance.

To Luciana, who brought joy and laughter back into my life during challenging times.

And to all those unnamed, if these words strike a chord, you have touched my life in significant ways.

I have learned much from each of you.

# Declaration

I hereby declare that I

- alone wrote the submitted Bachelor's thesis without illicit or improper assistance.
- did not use any materials other than those listed in the bibliography and that all passages taken from these sources in part or in full have been marked as citations and their sources cited individually in the thesis. Citations include the version (edition and year of publication), the volume and page numbers of the cited work.
- have not submitted this Bachelor's thesis to another institution and that it has never been used for other purposes or to fulfill other requirements, in part or in full.

---

Luciano Melodia M.A., B.A.  
Friedrich-Alexander University  
Erlangen-Nürnberg

# Acknowledgements

I extend heartfelt thanks to Prof. Dr. Catherine Meusburger, who not only taught me the rigors of the Bachelor's program in Mathematics but also illuminated the more complex concepts of Linear Algebra I and II. Her invaluable assistance during our early sessions, where we dissected exercise sheets in the study room, was crucial. I am particularly grateful for the moments she joined us after a refreshing cup of coffee, skillfully translating challenging problems into clear solutions on the blackboard with swift strokes of chalk.

My special gratitude is due to my supervisor, Prof. Dr. Kang Li. His readiness to engage in discussions was a beacon of motivation, especially when I was lost in the dense forest of scientific papers and theorems. Prof. Li's adeptness in distilling sprawling research into manageable segments for my bachelor's thesis was invaluable, and his excellent guidance has been a cornerstone of my still early academic journey in Mathematics.

Last but not least, I owe a tremendous debt of gratitude to some of my friends – Luciana Diaconescu, Philipp Gäbelein, Marie-Louise Isenberg, Michael Schleich, Sarah Wiesend, Adrian Hieber, Moritz Lanz, Sebastian Müller, Thomas Büttner, Anton Rechenauer, Constantin Winkler and Anton Hoof. Their companionship sweetened my days, their support was unwavering, and their presence essential to my achievements. Wherever they may find themselves now, I wish them every success and hope they know I am always here for them.

# Contents

<b>1</b>	<b>Motivation</b>	<b>1</b>
<b>2</b>	<b>Topological Spaces and Groups</b>	<b>2</b>
2.1	Simplicial Complexes . . . . .	3
2.2	Simplicial Homology . . . . .	7
2.3	Singular Homology . . . . .	16
2.3.1	Singular Chain Complexes . . . . .	20
2.3.2	Exact and Short Exact Sequences . . . . .	22
2.4	Relative Homology . . . . .	25
2.5	The Excision Theorem . . . . .	31
2.6	Equivalence of $H_d^\Delta(X)$ and $H_d(X)$ . . . . .	33
<b>3</b>	<b>Homological Persistence</b>	<b>36</b>
3.1	Persistent Homology . . . . .	36
3.1.1	Filtrations of Complexes . . . . .	36
3.1.2	Persistence Modules . . . . .	37
3.1.3	Persistent Homology on Complexes . . . . .	37
3.1.4	The Four Standard Persistence Modules . . . . .	39
3.2	Persistent (Co)homology . . . . .	39
3.2.1	Barcode Isomorphisms . . . . .	41
3.2.2	Persistent Chain Complexes . . . . .	45
3.2.3	Cohomology of Chain Complexes . . . . .	49
3.3	Zigzag Persistence . . . . .	49
3.3.1	Zigzag Modules . . . . .	49
3.3.2	Decompositions of Zigzag Modules . . . . .	50
3.3.3	Zigzag Modules and Filtrations . . . . .	50
3.3.4	Decomposition of Filtered Vector Spaces . . . . .	50
3.3.5	Streamlined Modules . . . . .	50

# Chapter 1

## Motivation

# Chapter 2

## Topological Spaces and Groups

Topological persistence is deeply rooted in algebraic topology, which studies topological spaces and functions based on algebraic objects and their properties. An example of this are homotopy and homology theories, which are essential for understanding the construction and connectedness of surfaces, a central aspect of data analysis. Persistent homology, the central tool of topological persistence, extends classical homology to identify features across multiple scales. Introduced by Edelsbrunner et al. in their seminal paper [7], persistent homology examines multi-scale topological features through a filtration process – an indexed family of nested spaces that starts with the empty set and progressively covers the entire space under study. Each stage of filtration represents a snapshot of the topological space at a different resolution and captures the appearance and disappearance of homology groups in different dimensions that encode topological properties such as connected components, holes and voids.

Mathematically, the persistence of homological features is visualized with the help of diagrams or barcode representations. These visual aids represent the emergence (birth) and vanishing (death) of topological features when the filter parameter changes. The duration of a feature's presence, represented by the length of its interval in the diagram, indicates its significance, with longer intervals suggesting features that represent the underlying data's likely true characteristics rather than mere noise. The robustness of persistent homology, in particular its resistance to minor perturbations in the data, is captured in the Stability Theorem. This theorem, proved by Cohen-Steiner, Edelsbrunner and Harer [1], states that slight variations in the input data lead to slight changes in the persistence barcodes. This property is crucial for practical applications as it guarantees



that the topological summaries are both reliable and meaningful for the actual underlying structures.

We start with basic concepts such as topological spaces and groups, which are crucial for understanding and encoding connectedness and other invariants. The discussion extends to simplicial complexes, which are essential for modeling data structures in topological data analysis. We explore simplicial and singular homology groups to accurately quantify topological features, and dwell on singular chain complexes and exact sequences to deepen the algebraic aspects of persistence theory. This will be useful in some of the later proofs when we discuss the algebraic nature of the persistence module.

## 2.1 Simplicial Complexes

We note that a set of points  $X = \{x_0, x_1, \dots, x_d\}$  in  $\mathbb{R}^n$  is affinely independent if no affine subspace of dimension less than  $d$  contains all the points in  $X$ . Such a set of points is commonly called point cloud.

**Definition 2.1.1 ( $d$ -Simplex)** [2, §2.1] A  $d$ -dimensional simplex  $\sigma^{(d)}$ , or  $d$ -simplex, is the set of all convex combinations of a set  $X = \{x_0, x_1, \dots, x_d\} \subset \mathbb{R}^n$ , where  $X$  consists of  $d + 1$  affinely independent points.

Formally,  $\sigma^{(d)}$  is defined by:

$$\sigma^{(d)} := \left\{ \sum_{i=0}^d \lambda_i x_i \mid \sum_{i=0}^d \lambda_i = 1, \lambda_i \geq 0 \right\}. \quad (2.1)$$

As a convention, the empty set is considered a face, corresponding to the simplex formed by the empty subset of vertices. Specifically, a 0-simplex corresponds to a single point, a 1-simplex to a line segment between two points, a 2-simplex to a triangle, and a 3-simplex to a tetrahedron. Notably, a  $d$ -simplex is homeomorphic to the  $d$ -dimensional disk  $D^d$ .

**Theorem 2.1.1 ( $d$ -Disk to  $d$ -Simplex)** The  $d$ -simplex  $\sigma^{(d)}$  is homeomorphic to the  $d$ -dimensional disk  $D^d$ .

**Proof** Define the standard  $d$ -simplex  $\sigma^{(d)}$  as

$$\sigma^{(d)} := \left\{ (x_0, \dots, x_d) \in \mathbb{R}^{d+1} \mid \sum_{i=0}^d x_i = 1, x_i \geq 0 \right\}, \quad (2.2)$$

and the  $d$ -dimensional disk  $D^d$  as

$$D^d := \left\{ (x_0, \dots, x_{d-1}) \in \mathbb{R}^d \mid \sum_{i=0}^{d-1} x_i^2 \leq 1 \right\}. \quad (2.3)$$

We construct a homeomorphism  $f : \sigma^{(d)} \rightarrow D^d$  by

$$f(x_0, \dots, x_d) = (\sqrt{x_0}, \dots, \sqrt{x_{d-1}}), \quad (2.4)$$

where  $x_d = 1 - \sum_{i=0}^{d-1} x_i$ . This map is well-defined since

$$\sum_{i=0}^{d-1} (\sqrt{x_i})^2 = \sum_{i=0}^{d-1} x_i \leq 1. \quad (2.5)$$

The inverse  $g : D^d \rightarrow \sigma^{(d)}$  is given by

$$g(y_0, \dots, y_{d-1}) = (y_0^2, \dots, y_{d-1}^2, 1 - \sum_{i=0}^{d-1} y_i^2), \quad (2.6)$$

ensuring that  $g$  is well-defined because  $\sum_{i=0}^{d-1} y_i^2 \leq 1$  implies  $1 - \sum_{i=0}^{d-1} y_i^2 \geq 0$ . Both  $f$  and  $g$  are continuous and are inverses of each other, as shown by  $f(g(y)) = y$  for all  $y \in D^d$  and  $g(f(x)) = x$  for all  $x \in \sigma^{(d)}$ .  $\square$

Furthermore, it is important to note that  $\sigma^{(d)}$  represents the convex hull of the points  $X = \{x_0, x_1, \dots, x_d\}$ , defined as the smallest convex subset of  $\mathbb{R}^n$  that contains all of these points. The faces of the simplex  $\sigma^{(d)}$ , with vertex set  $X$ , are formed by the simplices corresponding to subsets of  $X$ . A  $d$ -face of a simplex consists of a subset of the vertices with cardinality  $d + 1$ . The faces of a  $d$ -simplex with dimensions less than  $d$  are known as its proper faces. Two simplices are considered properly situated if their intersection is either empty or a face of both simplices. By identifying simplices along entire faces, we can construct the corresponding simplicial complexes.

**Definition 2.1.2 (Simplicial Complex)** [2, §2.2] A simplicial complex  $K$  is a finite collection of simplices that satisfies the following properties:

1. For every simplex  $\sigma^{(d)}$  in  $K$ , and every face  $\tau^{(k)}$  of  $\sigma^{(d)}$  with  $k < d$ , it follows that  $\tau^{(k)}$  is also in  $K$ .
2. Any two simplices  $\sigma^{(d)}$  and  $\tau^{(k)}$  in  $K$  are properly situated; that is, their intersection is either empty or a face of both simplices.

The dimension of a simplicial complex  $K$  is defined as the highest dimension among its simplices. For a simplicial complex  $K$  in  $\mathbb{R}^n$ , the underlying space  $|K|$  is the union of all the simplices in  $K$ . The topology of  $K$  is determined by the topology induced on  $|K|$  by  $\mathbb{R}^n$ 's standard topology. Notably, when the vertex set is specified, a simplicial complex in  $\mathbb{R}^n$  can be fully characterized by listing its simplices. Thus, it can be described purely in terms of combinatorics using abstract simplicial complexes.

**Definition 2.1.3 (Abstract Simplicial Complex)** [2, §2.3] Consider a finite set  $V = \{v_0, \dots, v_d\}$ . An abstract simplicial complex  $\tilde{K}$  with vertex set  $V$  is a collection of finite subsets of  $V$  that satisfies the following conditions:

1. Every singleton set  $\{v_i\}$ , where  $v_i \in V$ , is included in  $\tilde{K}$ .
2. If a set  $\sigma^{(d)}$  is in  $\tilde{K}$  and  $\tau^{(k)}$  is a subset of  $\sigma^{(d)}$ , then  $\tau^{(k)}$  must also be in  $\tilde{K}$ .

The abstract simplicial complex  $\tilde{K}$  associated with a simplicial complex  $K$  is commonly referred to as its vertex scheme. Conversely, if an abstract complex  $\tilde{K}$  serves as the vertex scheme for a complex  $K$  in  $\mathbb{R}^n$ , then  $K$  is known as a geometric realization of  $\tilde{K}$ .

**Proposition 2.1.2** *Every finite abstract simplicial complex  $\tilde{K}$  can be geometrically realized in Euclidean space.*

**Proof** Let  $\{v_0, v_1, \dots, v_d\}$  denote the vertex set of  $\tilde{K}$ , with  $0 \leq d < n$  representing the number of vertices in  $\tilde{K}$ . Consider  $\sigma^{(d-1)} \subset \mathbb{R}^n$ , the simplex formed by the span of  $\{e_1, e_2, \dots, e_d\}$ , where  $e_i$  represents the  $i$ -th unit vector. In this context,  $K$  refers to the subcomplex of  $\sigma^{(d-1)}$  such that  $[e_{i_0}, \dots, e_{i_l}]$  is a  $l$ -simplex of  $K$  with  $0 \leq l \leq d$  if and only if  $[v_{i_0}, \dots, v_{i_l}]$  is a simplex of  $\tilde{K}$ .  $\square$

Pay attention, that this result is in particular interesting for data analysis, as computer aided methods deal with finite point sets. All realizations of an abstract simplicial complex are homeomorphic to each other. The specific realization mentioned above is referred to as the natural realization.

**Proposition 2.1.3 (Equivalence of Geometric Realizations)** *Let  $\tilde{K}$  be an abstract simplicial complex. Then any two geometric realizations  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$  of  $\tilde{K}$  are homeomorphic.*

**Proof** Consider two geometric realizations  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$  of the abstract simplicial complex  $\tilde{K}$ . Each realization  $|\tilde{K}|_i$  is constructed by mapping the vertices of  $\tilde{K}$  to points in  $\mathbb{R}^d$ , and by identifying each simplex in  $\tilde{K}$  with the convex hull of its image under this mapping. By construction, each simplex in  $|\tilde{K}|_i$  corresponds homeomorphically to a standard  $l$ -simplex in  $\mathbb{R}^d$  for the appropriate  $l$  for integers  $0 \leq l \leq d$ .

To establish a homeomorphism between  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$ , define a map  $f : |\tilde{K}|_1 \rightarrow |\tilde{K}|_2$  that acts on each vertex  $v$  of  $\tilde{K}$  by mapping it from its image in  $|\tilde{K}|_1$  to its image in  $|\tilde{K}|_2$ . Extend  $f$  linearly to each simplex in  $|\tilde{K}|_1$ , where a simplex is represented as a convex combination of the images of its vertices. Given the two geometric realizations  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$  of  $\tilde{K}$ , where vertices  $v_i$  in  $|\tilde{K}|_1$  are mapped to points  $p_i$  in  $\mathbb{R}^d$  and in  $|\tilde{K}|_2$  to points  $q_i$ , the map  $f : |\tilde{K}|_1 \rightarrow |\tilde{K}|_2$  is defined such that for each vertex  $v_i$ , corresponding to  $p_i$  in  $|\tilde{K}|_1$ , we set  $f(p_i) = q_i$ . This map is linearly extended across each simplex in  $|\tilde{K}|_1$ .

Specifically, for a point  $x$  and an integer  $0 \leq k \leq d$  within a simplex of  $|\tilde{K}|_1$ , expressed as a convex combination  $x = \sum_{j=1}^k \lambda_j p_{i_j}$ , where  $\{v_{i_1}, v_{i_2}, \dots, v_{i_k}\}$  denote the vertices of the simplex and  $\lambda_j \geq 0$  with  $\sum_{j=1}^k \lambda_j = 1$ , the function  $f$  is defined by  $f(x) = \sum_{j=1}^k \lambda_j q_{i_j}$ .

This ensures that  $f$  is a continuous and bijective map, preserving the simplicial structure and thereby establishing  $f$  as a homeomorphism between  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$ .  $f$  is continuous on each simplex because it represents a linear transformation on compact convex subsets in  $\mathbb{R}^d$ . Since both  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$  are topologized by the quotient of their respective disjoint unions of simplices under an equivalence relation that glues them along common faces, and since  $f$  preserves these gluing conditions,  $f$  is continuous globally. Each simplex and its face structure in  $|\tilde{K}|_1$  maps uniquely onto a corresponding simplex and face structure in  $|\tilde{K}|_2$ , rendering  $f$  bijective.

Similarly, the inverse map  $g : |\tilde{K}|_2 \rightarrow |\tilde{K}|_1$  can be constructed by reversing the roles of  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$ . It also maintains continuity for analogous reasons to  $f$ .

Thus,  $|\tilde{K}|_1$  and  $|\tilde{K}|_2$  are homeomorphic. □

Furthermore, it has been proven that any finite abstract simplicial complex of dimension  $d$  can be realized as a simplicial complex in  $\mathbb{R}^{2d+1}$ .

**Theorem 2.1.4 (Realizations into  $\mathbb{R}^{2d+1}$ )** *Any finite abstract simplicial complex of dimension  $d$  can be realized as a simplicial complex in  $\mathbb{R}^{2d+1}$ .*

**Proof** Let  $\tilde{K}$  be a finite abstract simplicial complex of dimension  $d$ . We will construct an injective geometric realization  $f : \tilde{K} \rightarrow \mathbb{R}^{2d+1}$ .

First, define an injective map  $\tilde{f} : V(\tilde{K}) \rightarrow \mathbb{R}^{2d+1}$  for the vertex set  $V(\tilde{K})$  of  $\tilde{K}$ . Since  $V(\tilde{K})$  is finite and  $\mathbb{R}^{2d+1}$  has sufficient dimensionality, injectivity is guaranteed. Adjust  $\tilde{f}$  if necessary to ensure the images of the vertices of each simplex  $\sigma \in \tilde{K}$  are affinely independent. This can be achieved by slight perturbations within  $\mathbb{R}^{2d+1}$ , leveraging the ample dimensionality to avoid overlaps. We extend  $\tilde{f}$  to a map  $f : \tilde{K} \rightarrow \mathbb{R}^{2d+1}$  by defining it on each simplex  $\sigma = [v_0, \dots, v_k]$  through the unique affine map  $v_i \mapsto \tilde{f}(v_i)$ . The injective and affine properties of  $\tilde{f}$  on vertices guarantee that  $f$  is injective over each simplex and preserves the simplicial structure. Specifically, for any simplices  $\sigma, \tau \in \tilde{K}$ , we have:

$$f(|\sigma| \cap |\tau|) = f(|\sigma|) \cap f(|\tau|). \quad (2.7)$$

Thus,  $f$  provides an injective geometric realization of  $\tilde{K}$  as a simplicial complex in  $\mathbb{R}^{2d+1}$ .  $\square$

## 2.2 Simplicial Homology

Given a set  $V$  representing the vertices of a  $d$ -simplex  $\sigma^{(d)}$ , we can establish an orientation for the simplex by selecting a specific ordering of the vertices. If the vertex ordering differs from our chosen order by an odd permutation, the orientation is considered reversed, while even permutations preserve the orientation. Thus, a simplex can have only two possible orientations. Moreover, the orientation of a  $d$ -simplex induces an orientation on its  $(d-1)$ -faces. Specifically, if  $\sigma^{(d)} := (v_0, v_1, \dots, v_d)$  represents an oriented  $d$ -simplex, then the orientation of the  $(d-1)$ -face  $\tau^{(d-1)}$  of  $\sigma^{(d)}$ , omitting the vertex  $v_i$ , is given by

$$\tau_i^{(d-1)} = (-1)^i (v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_d). \quad (2.8)$$

**Definition 2.2.1 ( $d$ -Chain)** [11, §2.3] Given a set  $\{\sigma_1^{(d)}, \dots, \sigma_k^{(d)}\}$  of arbitrarily oriented  $d$ -simplices in a complex  $K$  and an abelian group  $G$ , a

$d$ -chain  $c$  with coefficients  $g_i \in G$  is defined as a formal sum:

$$c := g_0\sigma_0^{(d)} + g_1\sigma_1^{(d)} + \dots + g_k\sigma_k^{(d)} = \sum_{i=0}^k g_i\sigma_i^{(d)}. \quad (2.9)$$

Henceforth, we will assume that  $G = (\mathbb{Z}, +)$ .

**Lemma 2.2.1** *The set of simplicial  $d$ -chains  $C_d^\Delta$  forms an abelian group  $(C_d^\Delta, +)$ .*

**Proof** The identity element of the group is the empty chain, given by:

$$e_{C_d^\Delta} = \sum_{i \in \emptyset} g_i\sigma_i^{(d)} = e_G = 0. \quad (2.10)$$

The sum of two chains is defined as:

$$c + c' = \sum_{i=0}^k g_i\sigma_i^{(d)} + \sum_{j=1}^l g'_j\sigma_j^{(d)}, \quad (2.11)$$

which simplifies to:

$$c + c' = \sum_{i=0}^{\min(k,l)} (g_i + g'_i)\sigma_i^{(d)} + \begin{cases} \sum_{j=\min(k,l)+1}^{\max(k,l)} g'_j\sigma_j^{(d)} & \text{if } k > l, \\ 0 & \text{if } k = l, \\ \sum_{j=\min(k,l)+1}^{\max(k,l)} g'_j\sigma_j^{(d)} & \text{if } k < l. \end{cases} \quad (2.12)$$

Hence,  $c + c' \in C_d^\Delta$ .

The associativity of the group operation in  $C_d^\Delta$  follows directly from the associativity of the group operation in  $G$ .

The inverse element is given by:

$$c + (-c) = \sum_{i=0}^k g_i\sigma_i^{(d)} + \sum_{i=0}^k (-g_i)\sigma_i^{(d)} = \sum_{i=0}^k (g_i - g_i)\sigma_i^{(d)} = e_{C_d^\Delta}. \quad (2.13)$$

Thus,  $(C_d^\Delta, +)$  is an abelian group.  $\square$

**Definition 2.2.2 (Boundary)** [9, p.106] Let  $\sigma^{(d)}$  be an oriented  $d$ -simplex in a complex  $K$ . The boundary of  $\sigma^{(d)}$  is defined as the simplicial  $(d-1)$ -

chain of  $K$  with coefficients in the abelian group  $G$ , given by

$$\partial(\sigma^{(d)}) = \sum_{i=0}^d (-1)^i \sigma_i^{(d-1)}, \quad (2.14)$$

where  $\sigma_i^{(d-1)}$  is a  $(d-1)$ -face of  $\sigma^{(d)}$ . If  $d = 0$ , we define  $\partial(\sigma^{(0)}) = 0$ .

Since  $\sigma^{(d)}$  is an oriented simplex, the  $\sigma_i^{(d-1)}$  faces also have associated orientations. We extend the definition of the boundary linearly to elements of  $C_d^\Delta$ .

**Lemma 2.2.2** *The boundary operator is a group homomorphism*

$$\partial : C_d^\Delta \rightarrow C_{d-1}^\Delta. \quad (2.15)$$

**Proof** We define the boundary operator for a  $d$ -chain  $c = \sum_{i=0}^k g_i \sigma_i^{(d)}$ :

$$\partial(c) = \sum_{i=0}^k g_i \partial(\sigma_i^{(d)}) \quad (2.16)$$

$$= \sum_{i=0}^k g_i \sum_{j=0}^d (-1)^j \sigma_{i,j}^{(d-1)} \quad (2.17)$$

$$= \sum_{i=0}^k \sum_{j=0}^d g_i (-1)^j \sigma_{i,j}^{(d-1)}, \quad (2.18)$$

which is an element of  $C_{d-1}^\Delta$ , where  $\sigma_{i,j}^{(d-1)}$  are the  $(d-1)$ -faces of the  $d$ -simplices  $\sigma_i^{(d)}$  in  $K$ .

To verify that  $\partial$  is a group homomorphism, consider two  $d$ -chains  $c = \sum_{i=0}^k g_i \sigma_i^{(d)}$  and  $c' = \sum_{j=0}^l g'_j \sigma_j^{(d)}$ . We compute:

$$\partial(c + c') = \partial \left( \sum_{i=0}^k g_i \sigma_i^{(d)} + \sum_{j=0}^l g'_j \sigma_j^{(d)} \right) \quad (2.19)$$

$$= \sum_{i=0}^k g_i \partial(\sigma_i^{(d)}) + \sum_{j=0}^l g'_j \partial(\sigma_j^{(d)}) \quad (2.20)$$

$$= \partial(c) + \partial(c'). \quad (2.21)$$

Thus,  $\partial$  is a group homomorphism. □

**Example 2.2.3** *Let's consider the 2-simplex  $\sigma^{(2)}$  with vertices  $v_0, v_1$ , and  $v_2$ . The 1-faces of this simplex are:*

$$e_0 = (v_1, v_2), \quad \text{connecting } v_1 \text{ and } v_2, \quad (2.22)$$

$$e_1 = (v_2, v_0), \quad \text{connecting } v_2 \text{ and } v_0, \quad (2.23)$$

$$e_2 = (v_0, v_1), \quad \text{connecting } v_0 \text{ and } v_1. \quad (2.24)$$

*Now, let's proceed with the computation:*

$$\partial(\partial(\sigma^{(2)})) = \partial(e_0 + e_1 + e_2) \quad (2.25)$$

$$= \partial(e_0) + \partial(e_1) + \partial(e_2) \quad (2.26)$$

$$= \partial(v_1, v_2) + \partial(v_2, v_0) + \partial(v_0, v_1) \quad (2.27)$$

$$= [(v_2) - (v_1)] + [(v_0) - (v_2)] + [(v_1) - (v_0)] \quad (2.28)$$

$$= 0. \quad (2.29)$$

*We observe that  $C_0^\Delta$  is an abelian group and that oppositely oriented simplices cancel each other out, resulting in:*

$$\partial(\partial(\sigma^{(2)})) = 0. \quad (2.30)$$

*This property can be generalized to higher dimensions through induction. Therefore, since  $\partial$  is a linear operator and the chain  $c$  is a sum of  $d$ -simplices, we can conclude that:*

$$\partial^2(c) = 0 \quad \text{for any } d\text{-chain } c \text{ in } C_d^\Delta. \quad (2.31)$$

*Consequently, the boundary of the boundary is zero. Moreover, if the boundary of a simplex is zero, it is referred to as a cycle. By this definition, we can deduce that the boundary of any simplex is a cycle.*

**Definition 2.2.4 ( $d$ -Cycle)** [9, p.106] A  $d$ -chain is called a  $d$ -cycle if its boundary is equal to zero.

We denote the set of  $d$ -cycles of a complex  $K$  over the group  $\mathbb{Z}$  as  $Z_d^\Delta$ , the simplicial cycle group. It is important to note that  $Z_d^\Delta$  is a subgroup of  $C_d^\Delta$  and can also be expressed as:

$$Z_d^\Delta = \ker(\partial_d). \quad (2.32)$$



A  $d$ -cycle of a  $k$ -complex  $K$  is said to be homologous to zero if it can be expressed as the boundary of a  $(d+1)$ -chain in  $K$ , where  $d = 0, 1, \dots, k-1$ . In other words, a cycle is considered a boundary if it can be 'filled in' by a higher-dimensional chain. This equivalence relation is denoted as  $c \sim 0$ .

**Definition 2.2.5 (Boundary Group)** [11, §2.3] The subgroup of  $Z_d^\Delta$  consisting of boundaries is referred to as the simplicial boundary group  $B_d^\Delta$ .

It is worth noting that  $B_d^\Delta$  is equal to the image of the boundary operator  $\partial_{d+1}$ . Since  $B_d^\Delta$  is a subgroup of  $Z_d^\Delta$  and  $Z_d^\Delta$  is an abelian group, every subgroup of  $Z_d^\Delta$  is normal. Therefore, we can construct the group quotient:

$$H_d^\Delta = Z_d^\Delta / B_d^\Delta. \quad (2.33)$$

**Definition 2.2.6 (Simplicial Homology Group)** [9, §2.1] The group  $H_d^\Delta$  represents the  $d$ -dimensional simplicial homology group of the complex  $K$  over  $\mathbb{Z}$ . It is expressed as the group quotient:

$$H_d^\Delta = \ker(\partial_d) / \text{im}(\partial_{d+1}). \quad (2.34)$$

Next, we want to examine the structure of this homology group by shedding light on its connection to the connected components of a simplicial complex. We will find that the homology groups of the connected components of the complex, which in turn form a complex themselves, yield the direct sum of the homology group of the entire complex.

**Definition 2.2.7** *A subcomplex is a subset  $S$  of the simplices belonging to a complex  $K$ , where  $S$  itself forms a complex.*

**Definition 2.2.8** *The collection of all simplices in a complex  $K$  with dimensions less than or equal to  $d$  is referred to as the  $d$ -skeleton of  $K$ .*

By definition, the  $d$ -skeleton forms a subcomplex.

**Definition 2.2.9** *A complex  $K$  is considered connected if it cannot be expressed as the disjoint union of two or more non-empty subcomplexes. A geometric complex is path-connected if there exists a path consisting of 1-simplices connecting any vertex to any other vertex.*

**Lemma 2.2.3** *A geometric complex is path-connected if and only if it is connected.*

**Proof** Suppose  $K$  is path-connected. Assume for contradiction that  $K$  can be expressed as the disjoint union of two non-empty subcomplexes  $L$  and  $M$ . Since  $K$  is path-connected, there exists a path of 1-simplices between any two vertices in  $K$ . Let  $l \in L$  and  $m \in M$  be any two vertices. By path-connectedness, there is a path from  $l$  to  $m$ , which contradicts the assumption that  $L$  and  $M$  are disjoint. Therefore,  $K$  is connected.

Suppose  $K$  is connected. Pick any vertex  $v \in K$ . Let  $L$  denote the subcomplex of  $K$  containing all vertices reachable from  $v$  via paths of 1-simplices. If  $L \neq K$ , then  $L$  and  $K \setminus L$  form a disjoint union of two non-empty subcomplexes, contradicting the connectedness of  $K$ . Hence,  $L = K$ , and  $K$  is path-connected.  $\square$

**Theorem 2.2.4** [9, p.105ff] Let  $K_1, \dots, K_p$  be the collection of all connected components of a complex  $K$ . Furthermore, let  $H_{d_i}^\Delta$  represent the  $d$ -th simplicial homology group of  $K_i$ , and  $H_d^\Delta$  denote the  $d$ -th simplicial homology group of  $K$ . In this context, we can establish that  $H_d^\Delta$  is isomorphic to the direct sum  $H_{d_1}^\Delta \oplus \dots \oplus H_{d_p}^\Delta$ .

**Proof** Let  $C_d^\Delta$  represent the group of simplicial  $d$ -chains of  $K$ , and let  $K_i$  denote the  $i$ -th component of  $K$ . Define  $C_{d_i}^\Delta$  as the group of simplicial  $d$ -chains of  $K_i$ . It is evident that  $C_{d_i}^\Delta$  is a subgroup of  $C_d^\Delta$ . Furthermore, we observe that  $C_d^\Delta$  can be expressed as the direct sum of  $C_{d_1}^\Delta, \dots, C_{d_p}^\Delta$ :

$$C_d^\Delta = C_{d_1}^\Delta \oplus \dots \oplus C_{d_p}^\Delta. \quad (2.35)$$

Our goal is to demonstrate that a similar decomposition applies to the groups  $B_d^\Delta$  and  $Z_d^\Delta$ . By considering  $B_{d_i}^\Delta$  as the image of  $\partial_{d+1}$  restricted to the subgroup  $C_{d_i}^\Delta$ , we can represent the group  $B_d^\Delta$  as the direct sum of these restrictions:

$$B_d^\Delta = B_{d_1}^\Delta \oplus \dots \oplus B_{d_p}^\Delta. \quad (2.36)$$

Thus, for any element  $c \in C_{d+1}^\Delta$ , we have:

$$c = c_1 + \dots + c_p, \quad \partial_{d+1}(c) = \partial_{d+1}(c_1) + \dots + \partial_{d+1}(c_p) \in B_d^\Delta, \quad (2.37)$$

where  $c_i \in C_{(d+1)_i}^\Delta$ .

Let us define  $Z_{d_i}^\Delta$  as the intersection of the kernel of  $\partial_d$  and  $C_{d_i}^\Delta$ . It follows

that  $Z_d^\Delta$  can be expressed as the direct sum of  $Z_{d_1}^\Delta, \dots, Z_{d_p}^\Delta$ :

$$Z_d^\Delta = Z_{d_1}^\Delta \oplus \dots \oplus Z_{d_p}^\Delta. \quad (2.38)$$

To verify this, consider an element  $c \in C_d^\Delta$  that belongs to  $Z_d^\Delta$ . We require  $\partial_d(c) = 0$ . However, we can express  $\partial_d(c)$  as  $\partial_d(c_1) + \dots + \partial_d(c_p)$ . Therefore, for  $\partial_d(c) = 0$ , it must be that  $\partial_d(c_i) = 0$ , indicating that  $c_i \in Z_{d_i}^\Delta$ .

Since both  $Z_d^\Delta$  and  $B_d^\Delta$  can be decomposed componentwise, we conclude that:

$$Z_d^\Delta / B_d^\Delta = Z_{d_1}^\Delta / B_{d_1}^\Delta \oplus \dots \oplus Z_{d_p}^\Delta / B_{d_p}^\Delta, \quad (2.39)$$

and consequently:

$$H_d^\Delta = H_{d_1}^\Delta \oplus \dots \oplus H_{d_p}^\Delta. \quad (2.40)$$

□

**Proposition 2.2.5** [9, p.108f] If  $K$  is a connected complex and  $c$  is a 0-chain with  $I(c) = 0$ , then  $I(c) = 0$  is equivalent to  $c \sim 0$ , where  $\sim$  denotes homology equivalence. Furthermore, in this case, the zeroth simplicial homology group  $H_0^\Delta(K, \mathbb{Z})$  is isomorphic to the integers  $\mathbb{Z}$ .

**Proof** Let  $\sigma^{(1)} = (v_0, v_1)$  be a 1-simplex. For the chain  $c = \partial_1(g\sigma^{(1)}) = gv_1 - gv_0$ , we have  $c \sim 0$ , and it is clear that  $I(c) = g - g = 0$ . Since  $I(c + c') = I(c) + I(c')$ ,  $I$  is a group homomorphism. For any 1-chain  $c \in C_1^\Delta$  of the form  $\sum_{i=0}^k g_i \sigma_i^{(1)}$ , where  $\sigma_i^{(1)} = (v_i, v_{i+1})$ , we have:

$$c = \partial_1(c) \sim 0 \implies I(c) = I(\partial_1(c)) = 0. \quad (2.41)$$

Consider two vertices  $v$  and  $w$  in  $K$ . Since  $K$  is connected, there exists a path between  $v$  and  $w$  consisting of 1-simplices  $\sigma_i^{(1)} = (v_i, v_{i+1})$ ,  $i = 0, \dots, k-1$ , where  $v_0 = v$  and  $v_k = w$ . The boundary of the chain  $c = \sum_{i=0}^k g \sigma_i^{(1)}$  is given by:

$$\partial_1(c) = \sum_{i=0}^k g \partial_1(\sigma_i^{(1)}) = \sum_{i=0}^k g [(v_{i+1}) - (v_i)] = gw - gv. \quad (2.42)$$

Since  $\partial_1(c)$  is a boundary, we have  $c = \partial_1(c) \sim 0$ . Thus,  $(gw - gv) \sim 0$ ,

implying  $gw \sim gv$ . Therefore, any 0-chain  $c$  in  $K$  is homologous to the chain  $gv$ . We observe that homologous chains have equal indices, i.e.,  $I(c) = I(gv) = g$ . Thus,  $c \sim gv \implies c \sim I(c)v$ . This shows that if  $I(c) = 0$ , then  $c \sim 0$ . Hence,  $I(c) = 0$  is equivalent to  $c \sim 0$ .

Since  $I$  is a homomorphism from  $C_0^\Delta = Z_0^\Delta$  to  $\mathbb{Z}$ , for any 0-simplex  $c$  and  $g \in \mathbb{Z}$ , the chain  $gc \in C_0^\Delta$  is a cycle with  $I(gc) = g$ . Therefore,  $I(Z_0^\Delta) = \mathbb{Z}$ . Since  $I(c) = 0$  is equivalent to  $c \sim 0$ , we have  $B_0^\Delta = \ker(I)$ . This implies that:

$$H_0^\Delta = Z_0^\Delta / B_0^\Delta \cong \mathbb{Z}. \quad (2.43)$$

□

**Corollary 2.2.6** *The 0-dimensional simplicial homology group of a complex  $K$  over  $\mathbb{Z}$  can be represented as  $\mathbb{Z}^p = \bigoplus_p \mathbb{Z}$ , where  $p$  denotes the number of connected components present in  $K$ .*

**Proof** From Proposition 2.2.5, we know that if  $K$  is a connected complex, then the zeroth simplicial homology group  $H_0^\Delta(K, \mathbb{Z})$  is isomorphic to  $\mathbb{Z}$ .

Now, consider a complex  $K$  that consists of  $p$  connected components  $K_1, K_2, \dots, K_p$ . The zero-dimensional simplicial homology group  $H_0^\Delta(K, \mathbb{Z})$  can be expressed as the direct sum of the zero-dimensional homology groups of its connected components:

$$H_0^\Delta(K, \mathbb{Z}) = H_0^\Delta(K_1, \mathbb{Z}) \oplus H_0^\Delta(K_2, \mathbb{Z}) \oplus \dots \oplus H_0^\Delta(K_p, \mathbb{Z}). \quad (2.44)$$

Since each  $K_i$  is a connected component, by Proposition 2.2.5, each  $H_0^\Delta(K_i, \mathbb{Z})$  is isomorphic to  $\mathbb{Z}$ . Therefore, we have:

$$H_0^\Delta(K, \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \dots \oplus \mathbb{Z} = \mathbb{Z}^p. \quad (2.45)$$

Thus, the zero-dimensional simplicial homology group of  $K$  over  $\mathbb{Z}$  is represented as  $\mathbb{Z}^p$ , where  $p$  denotes the number of connected components present in  $K$ . □

### Example 2.2.10

- *The zeroth homology group of the circle is isomorphic to  $\mathbb{Z}$ . Consider a simplicial representation of the circle using four 1-simplices:  $\sigma_0 = (v_0, v_1)$ ,*

$\sigma_1 = (v_1, v_2)$ ,  $\sigma_2 = (v_2, v_3)$ , and  $\sigma_3 = (v_3, v_0)$ . The group  $Z_0^\Delta$  consists of sums over the four zero-simplices  $v_0, v_1, v_2$ , and  $v_3$  with coefficients in  $\mathbb{Z}$ . Let  $c$  be a zero-chain with non-zero coefficients given by:

$$c = g_0 v_0 + g_1 v_1 + g_2 v_2 + g_3 v_3. \quad (2.46)$$

To reduce it to an element of  $H_0^\Delta$ , subtract the chain  $c' = g_3 v_2 - g_3 v_3 \sim 0$ :

$$c - c' = g_0 v_0 + g_1 v_1 + (g_2 - g_3) v_2. \quad (2.47)$$

By repeating this process, we obtain a new chain:

$$c'' = (g_0 - g_1 + g_2 - g_3) v_0. \quad (2.48)$$

Since  $c'' \sim c$ , it represents an element of  $H_0^\Delta$ . Moreover, since  $g_i \in \mathbb{Z}$ , we can write  $(g_0 - g_1 + g_2 - g_3) \in \mathbb{Z}$  as  $c'' = g v_0$ , where  $g$  is an element of  $\mathbb{Z}$ . Therefore, we can choose any  $g$ , implying that  $H_0^\Delta \cong \mathbb{Z}$ .

- We will demonstrate that  $H_d^\Delta(S^d) \cong \mathbb{Z}$ . The  $d$ -simplex  $\sigma^{(d)}$  and the  $d$ -ball are homeomorphic, and their boundaries, which consist of  $(d-1)$ -simplices, are homeomorphic to the  $d$ -sphere. Thus, the appropriate simplicial structure to impose on  $S^d$  is that of the boundary of the  $(d+1)$ -simplex  $\sigma^{(d+1)}$ . Let  $\{v_0, \dots, v_d\}$  denote the set of vertices of  $\sigma^{(d+1)}$ . This set is not oriented, and the orientations of the  $(d-1)$ -simplices can be arbitrarily determined. We'll utilize their numbering to establish orientations.

Consequently, all  $d$ -chains on this structure can be expressed as:

$$c = \sum_{i=0}^{d+1} g_i (v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_d), \quad (2.49)$$

where  $g_i \in \mathbb{Z}$ .

Since  $\sigma^{(d+1)}$  itself is not part of the structure, there are no boundaries in  $Z_d^\Delta$ , the group of simplicial cycles. Thus,  $H_d^\Delta = Z_d^\Delta / B_d^\Delta$  represents the group of simplicial cycles. If  $c \in Z_d^\Delta$ , then  $\partial_{d+1}(c) = 0$ . Using Eq. 2.49, we have:

$$\partial_{d+1}(c) = \partial_{d+1} \left( \sum_{i=0}^{d+1} g_i (v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_d) \right) \quad (2.50)$$

$$= \sum_{i=0}^{d+1} g_i \left( \sum_{j=0}^{d+1} (-1)^j (v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_{j-1}, v_{j+1}, \dots, v_d) \right). \quad (2.51)$$

By rearranging this sum, we obtain terms of the form:

$$(g_k - g_l)(v_0, \dots, v_{j-1}, v_{j+1}, \dots, v_{i-1}, v_{i+1}, \dots, v_d), \quad (2.52)$$

where  $k, l = 0, \dots, d+1$  for all  $i, j = 0, \dots, d$ .

Each pair of  $d$ -simplices of  $\sigma^{(d+1)}$  intersects along a  $(d-1)$ -face. Therefore, we obtain terms of the form given in Eq. 2.52 for each of these faces. From this, we deduce that if  $\partial_d(c) = 0$ , we must have  $g_k = g_l$  for all  $k, l = 0, \dots, d+1$ . In other words,  $g_0 = g_1 = \dots = g_{d+1}$ . Consequently, our original  $d$ -chain is:

$$c = \sum_{i=0}^{d+1} g_0(v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_d), \quad (2.53)$$

allowing us to choose  $g_0$  from  $\mathbb{Z}$ . Thus, we conclude that  $H_d^\Delta(S^d) \cong \mathbb{Z}$ .

- We demonstrate that  $H_d^\Delta(D^d) = 0$ . The simplest simplicial structure for  $D^d$  is that of the  $d$ -simplex  $\sigma^{(d)}$ . Consequently, all  $d$ -chains can be expressed as  $c = g\sigma^{(d)}$ , where  $g \in \mathbb{Z}$ . This form is never a boundary, implying that  $H_d^\Delta = Z_d^\Delta$ . However,  $\partial_d(c) = 0$  is generally only true when  $g = 0$ . Thus, we conclude that  $H_d^\Delta(D^d) \cong 0$ .

## 2.3 Singular Homology

In the realm of lower dimensions, we have an intuitive idea of when two topological spaces are fundamentally 'equivalent'. To formalize and strengthen this intuition, we have developed various methods, one of which is the concept of homeomorphism. It would be highly desirable to establish a relation between the homology groups of homeomorphic spaces. Remarkably, it has been found that the homology groups of two topological spaces are isomorphic if they are homeomorphic. This fact requires verification.

To accomplish this task, we need a means of comparing homology groups. However, it is not immediately obvious how we can achieve this with the tools developed so far. In fact, this turns out to be quite a difficult problem. To circumvent this difficulty, we introduce the concept of singular homology. The basic principles underlying this concept are analogous to those we have already explored.

To compute  $Z_d(X)$ , we need to find the group of  $d$ -cycles in  $X$ . Since  $X$  is obtained by identifying opposite faces of  $\partial_d\sigma^{(d)}$ , a  $d$  cycle in  $X$  corresponds to a  $d$  cycle in  $\partial_d\sigma^{(d)}$ , which is not a boundary of a  $(d+1)$ -dimensional

simplex in  $\sigma^{(d)}$  is. In other words, a  $d$ -cycle in  $X$  corresponds to a  $d$ -cycle in  $\partial_d \sigma^{(d)}$ , which is not a boundary of a  $(d+1)$ -dimensional face of  $\sigma^{(d)}$ .

**Definition 2.3.1** *A singular  $d$ -simplex in a topological space  $X$  is a continuous map  $\tilde{\sigma}^{(d)} : \sigma^{(d)} \rightarrow X$ .*

We define the boundary map  $\partial_d$  in a similar manner as before. The boundary map, denoted as  $\partial_d$ , is a function that operates on the chain group  $C_d(X)$  and maps it to the chain group  $C_{d-1}(X)$ . It is defined as follows: For any singular  $d$ -simplex  $\tilde{\sigma}^{(d)}$  in  $X$ , the boundary map  $\partial_d(\tilde{\sigma}^{(d)})$  is obtained by summing over all the  $(d-1)$ -simplices that are obtained by removing one vertex from  $\tilde{\sigma}^{(d)}$ . Each term in the sum is multiplied by  $(-1)^i$ , where  $i$  represents the index of the removed vertex. In other words, if  $v_i$  represents the 0-simplex (vertex) of  $\tilde{\sigma}^{(d)}$ , then the boundary map can be expressed as:

$$\partial_d(\tilde{\sigma}^{(d)}) = \sum_i (-1)^i \tilde{\sigma}^{(d)}|_{[v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_d]}. \quad (2.54)$$

Here,  $v_i$  is a map that takes the 0-simplex  $\sigma^{(0)}$  to the corresponding vertex in  $X$ , such that  $v_i : \sigma^{(0)} \rightarrow X$  is continuous.

As mentioned earlier, when we apply the boundary map twice to a  $d$ -chain  $c$ , denoted as  $\partial^2(c)$  or  $\partial_{d-1}(\partial_d(c))$ , the result is always zero. This observation leads us to the idea of defining the singular homology groups in a similar way to the simplicial homology groups.

**Definition 2.3.2** *The singular homology group  $H_d(X)$  is the quotient:*

$$H_d(X) = \ker(\partial_d) / \text{im}(\partial_{d+1}). \quad (2.55)$$

In the following section, we will explore how this definition of homology allows us to establish a simple relationship between homeomorphic spaces and their corresponding homology groups. This relationship becomes apparent when we consider the fact that the definitions of  $H_d$  and  $H_d^\Delta$  are analogous. Intuitively, we would expect these two groups to be the same for triangulable topological spaces. However, this is not immediately obvious. One reason for this is that  $H_d^\Delta$  is finitely generated, while the chain group  $C_d(X)$ , from which we derived  $H_d$ , is uncountable.

Interestingly, for spaces where both simplicial and singular homology groups can be calculated, these two groups are indeed equivalent. We will provide a proof for this later on. But before we do, let us present some facts about singular homology that support the intuition that  $H_d$  is isomorphic to  $H_d^\Delta$ .

**Proposition 2.3.1** [9, p.108f] In the context of a topological space  $X$ ,  $H_d(X)$  is isomorphic to the direct sum  $H_d(X_1) \oplus \cdots \oplus H_d(X_p)$ , where  $X_i$  represents the path-connected components of  $X$ .

This equivalence serves as the counterpart to Proposition 2.2.5.

**Proof** As the maps  $\tilde{\sigma}^{(d)}$  are continuous, a singular simplex always has a path-connected image in  $X$ . Consequently,  $C_d(X)$  can be expressed as the direct sum of subgroups  $C_d(X_1) \oplus \cdots \oplus C_d(X_p)$ . The boundary map  $\partial$  functions as a homomorphism, thereby preserving this decomposition. Consequently,  $\ker(\partial_d)$  and  $\text{im}(\partial_{d+1})$  also decompose:

$$H_d(X) \cong H_d(X_1) \oplus H_d(X_2) \oplus \cdots \oplus H_d(X_p). \quad (2.56)$$

□

**Proposition 2.3.2** *The 0-dimensional homology group of a space  $X$  can be expressed as the direct sum of  $\mathbb{Z}$  copies, with each copy corresponding to a distinct path-component of  $X$ .*

**Proof** To establish the isomorphism  $H_0(X) \cong \mathbb{Z}$ , it is sufficient to consider the case where  $X$  is path-connected. For a 0-chain  $c$ , the boundary operator  $\partial_0(c)$  is always zero since the boundary of any 0-simplex vanishes. Consequently,  $\ker(\partial_0) = C_0(X)$ , which implies that:

$$H_0(X) = C_0(X) / \text{im}(\partial_1). \quad (2.57)$$

Define the map  $I : C_0(X) \rightarrow \mathbb{Z}$  by:

$$I(c) = \sum_i g_i \quad \text{for } c = \sum_i g_i \tilde{\sigma}^{(0)} \in C_0(X). \quad (2.58)$$

Our goal is to demonstrate that  $\ker(I) = \text{im}(\partial_1)$ . That is, for any 0-chain  $c$ ,  $I(c) = 0$  if and only if  $c \sim 0$ .

Suppose  $I(c) = 0$  for  $c = \sum_i g_i \tilde{\sigma}^{(0)}$ . Since  $X$  is path-connected, we can express  $c$  as the boundary of a 1-chain:

$$c = \partial_1 \left( \sum_j g_j \tilde{\sigma}^{(1)} \right) \quad \text{implying } c \sim 0. \quad (2.59)$$



If  $c \sim 0$ , then  $c = \partial_1 \left( \sum_j g_j \tilde{\sigma}^{(1)} \right)$  for some 1-chain  $\sum_j g_j \tilde{\sigma}^{(1)}$ . By linearity of  $I$ , we have:

$$I(c) = I \left( \partial_1 \left( \sum_j g_j \tilde{\sigma}^{(1)} \right) \right) = 0. \quad (2.60)$$

Thus,  $I$  induces an isomorphism between  $H_0(X)$  and  $\mathbb{Z}$  when  $X$  is path-connected.

For the general case where  $X$  has  $p$  path-components  $X_1, \dots, X_p$ , we apply the above argument to each component. Therefore,  $H_0(X)$  is isomorphic to the direct sum:

$$H_0(X) \cong \bigoplus_{i=1}^p \mathbb{Z}. \quad (2.61)$$

□

This correspondence serves as the parallel to Corollary 2.2.6.

**Example 2.3.3** *Alternative proof that  $H_d(S^d) \cong \mathbb{Z}$ . To prove that the  $d$ -th homology group of the  $d$ -sphere is isomorphic to  $\mathbb{Z}$ , we utilize the singular homology approach. The  $d$ -th singular chain group  $C_d(S^d)$  consists of formal linear combinations of singular  $d$ -simplices in  $S^d$  with integer coefficients. First, we observe that  $S^d$  is a connected and compact topological space. According to the Hurewicz theorem, we have  $H_d(S^d) \cong \pi_d(S^d)$ , where  $\pi_d(S^d)$  denotes the  $d$ -th homotopy group of  $S^d$ . Since  $S^d$  is simply connected for  $d \geq 2$ , it follows that  $\pi_d(S^d) = 0$  for  $d \geq 2$ . However, for  $d = 1$ ,  $\pi_1(S^1) \cong \mathbb{Z}$ .*

*Now, to establish the isomorphism between  $\pi_1(S^1)$  and  $H_1(S^1)$ , consider the singular 1-chain group  $C_1(S^1)$ , which consists of formal linear combinations of singular 1-simplices in  $S^1$  with integer coefficients. Let  $c$  be a singular 1-chain in  $C_1(S^1)$  expressed as:*

$$c = \sum_i g_i \tilde{\sigma}_i^{(1)}, \quad (2.62)$$

*where  $g_i \in \mathbb{Z}$  and  $\tilde{\sigma}_i^{(1)}$  are singular 1-simplices. The boundary of  $c$  is given by:*

$$\partial_1(c) = \sum_i g_i \partial_1(\tilde{\sigma}_i^{(1)}). \quad (2.63)$$

Since  $S^1$  is a 1-dimensional manifold, the boundary of any singular 1-simplex  $\tilde{\sigma}_i^{(1)}$  is a formal linear combination of two points in  $S^1$ , each with opposite orientations:

$$\partial_1(\tilde{\sigma}_i^{(1)}) = p - q, \quad (2.64)$$

where  $p$  and  $q$  are points in  $S^1$ . Thus, we have:

$$\partial_1(c) = (p - q) \sum_i g_i, \quad (2.65)$$

where the sum  $\sum_i g_i$  is an integer. Consequently, the boundary of any singular 1-chain  $c$  in  $C_1(S^1)$  is of the form  $(p - q)k$ , where  $k$  is an integer.

This implies that:

$$H_1(S^1) = Z_1(S^1)/B_1(S^1) \cong \mathbb{Z}, \quad (2.66)$$

where  $Z_1(S^1)$  is the group of 1-cycles and  $B_1(S^1)$  is the group of 1-boundaries. In conclusion,  $H_d(S^d) \cong \pi_d(S^d) = 0$  for  $d \geq 2$ , and  $H_1(S^1) \cong \pi_1(S^1) \cong \mathbb{Z}$ . Thus, the  $d$ -th homology group of the  $d$ -sphere is isomorphic to  $\mathbb{Z}$ .

### 2.3.1 Singular Chain Complexes

To establish the equivalence between the groups  $H_d^\Delta$  and  $H_d$ , we will introduce some concepts that will aid us in our proof.

Since the maps  $\tilde{\sigma}^{(d)}$  are continuous, a singular simplex always has a path-connected image in  $X$ . Thus, we can express  $C_d(X)$  as the direct sum of subgroups  $C_d(X_1) \oplus \cdots \oplus C_d(X_p)$ , where each subgroup corresponds to a distinct path-component of  $X$ . This decomposition is preserved by the boundary map  $\partial$ , which is a homomorphism. Thus, both  $\ker(\partial_d)$  and  $\text{im}(\partial_{d+1})$  also split, leading to the conclusion that  $H_d(X)$  is isomorphic to

$$H_d(X_1) \oplus H_d(X_2) \oplus \cdots \oplus H_d(X_p). \quad (2.67)$$

These ideas will serve as valuable tools in proving the equivalence of the groups  $H_d^\Delta$  and  $H_d$ .

**Definition 2.3.4 (Chain Complex)** A chain complex is a sequence of abelian groups  $\{C_n\}_{n \in \mathbb{Z}}$  connected by homomorphisms  $\partial_n : C_n \rightarrow C_{n-1}$  (called boundary operators), such that the composition of any two consecutive maps is zero, i.e.,

$\partial_{n-1} \circ \partial_n = 0$  for all  $n$ . Formally, a chain complex is:

$$\cdots \xrightarrow{\partial_{n+2}} C_{n+1} \xrightarrow{\partial_{n+1}} C_n \xrightarrow{\partial_n} C_{n-1} \xrightarrow{\partial_{n-1}} \cdots \quad (2.68)$$

where  $\partial_{n-1} \circ \partial_n = 0$ .

**Example 2.3.5** The groups  $C_d(X)$  represent the collection of singular  $d$ -chains that form a part of a chain complex, where the boundary operator  $\partial_d$  is the map between these groups in the respective dimension:

$$\cdots \rightarrow C_{d+1}(X) \xrightarrow{\partial_{d+1}} C_d(X) \xrightarrow{\partial_d} C_{d-1}(X) \rightarrow \cdots \quad (2.69)$$

$$\cdots \rightarrow C_1(X) \xrightarrow{\partial_1} C_0(X) \xrightarrow{\partial_0} 0. \quad (2.70)$$

**Definition 2.3.6** A chain map  $f$  between two chain complexes  $(A, \partial^{(A)})$  and  $(B, \partial^{(B)})$  is a collection of maps  $f_d : A_d \rightarrow B_d$  such that for each  $d$ , the following condition holds:

$$\partial_{d-1}^{(B)} \circ f_d = f_{d-1} \circ \partial_d^{(A)}. \quad (2.71)$$

**Theorem 2.3.3 (Chain Maps)** [10, §1.3.1] A chain map  $f$  between two chain complexes  $(A, \partial^{(A)})$  and  $(B, \partial^{(B)})$  induces a homomorphism between their respective homology groups.

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_{d+2}^{(A)}} & A_{d+1} & \xrightarrow{\partial_{d+1}^{(A)}} & A_d & \xrightarrow{\partial_d^{(A)}} & A_{d-1} \xrightarrow{\partial_{d-1}^{(A)}} \cdots \\ & & \downarrow f_{d+1} & & \downarrow f_d & & \downarrow f_{d-1} \\ \cdots & \xrightarrow{\partial_{d+2}^{(B)}} & B_{d+1} & \xrightarrow{\partial_{d+1}^{(B)}} & B_d & \xrightarrow{\partial_d^{(B)}} & B_{d-1} \xrightarrow{\partial_{d-1}^{(B)}} \cdots \end{array} \quad (2.72)$$

**Proof** Given a chain map  $f$  between two chain complexes  $(A, \partial^{(A)})$  and  $(B, \partial^{(B)})$ , we want to show that  $f$  induces a homomorphism  $f_* : H_d(A) \rightarrow H_d(B)$ .

By definition of a chain map, we have  $f \circ \partial^{(A)} = \partial^{(B)} \circ f$ . Let  $[c] \in Z_d(A)$  be a cycle in  $A$ , i.e.,  $\partial_d^{(A)}(c) = 0$ . Applying  $f$  to this equation, we obtain:

$$f_{d-1}(\partial_d^{(A)}(c)) = \partial_d^{(B)}(f_d(c)). \quad (2.73)$$

Since  $\partial_d^{(A)}(c) = 0$ , we have:

$$f_{d-1}(0) = \partial_d^{(B)}(f_d(c)), \quad (2.74)$$

which implies that  $\partial_d^{(B)}(f_d(c)) = 0$ . Thus,  $f_d(c)$  is a cycle in  $B$ .

Now, let  $[b] \in B_d(A)$  be a boundary in  $A$ , i.e., there exists  $a \in A_{d+1}$  such that  $\partial_{d+1}^{(A)}(a) = b$ . Applying  $f$  to both sides, we obtain:

$$f_d(\partial_{d+1}^{(A)}(a)) = \partial_{d+1}^{(B)}(f_{d+1}(a)). \quad (2.75)$$

Since  $\partial_{d+1}^{(A)}(a) = b$ , we have:

$$f_d(b) = \partial_{d+1}^{(B)}(f_{d+1}(a)). \quad (2.76)$$

Therefore,  $f_d(b)$  is a boundary in  $B$ .

From the above, we see that  $f$  maps cycles in  $A$  to cycles in  $B$  and boundaries in  $A$  to boundaries in  $B$ . Hence,  $f$  induces a well-defined map  $f_* : H_d(A) \rightarrow H_d(B)$ .

To show that  $f_*$  is a homomorphism, let  $[c_1], [c_2] \in H_d(A)$  be two homology classes. We want to show that  $f_*([c_1] + [c_2]) = f_*([c_1]) + f_*([c_2])$ . Let  $c_1$  and  $c_2$  be representatives of  $[c_1]$  and  $[c_2]$ , respectively. Then,  $[c_1] + [c_2]$  is represented by  $c_1 + c_2$ . Applying  $f$  to both sides, we have:

$$f_d(c_1 + c_2) = f_d(c_1) + f_d(c_2). \quad (2.77)$$

Since  $f_d(c_1)$  and  $f_d(c_2)$  are cycles in  $B$ , we have:

$$[f_d(c_1 + c_2)] = [f_d(c_1)] + [f_d(c_2)]. \quad (2.78)$$

Therefore,  $f_*([c_1] + [c_2]) = f_*([c_1]) + f_*([c_2])$ .

Thus, we have shown that the chain map  $f$  induces a homomorphism  $f_* : H_d(A) \rightarrow H_d(B)$ .  $\square$

### 2.3.2 Exact and Short Exact Sequences

We can apply Theorem 2.3.3 to the case of singular homology. Consider two topological spaces  $X$  and  $Y$ . For any map  $f : X \rightarrow Y$ , we can define an induced homomorphism  $f_* : C_d(X) \rightarrow C_d(Y)$  by composing singular  $d$ -simplices  $\tilde{\sigma}^{(d)} : \sigma^{(d)} \rightarrow X$  with  $f$ . Specifically, we have

$$f_* \circ \tilde{\sigma}^{(d)} = f \circ \tilde{\sigma}^{(d)} : \sigma^{(d)} \rightarrow Y. \quad (2.79)$$

We can extend this definition by applying  $f_*$  to  $d$ -chains in  $C_d(X)$ . This

gives us the following commutative diagram:

$$\begin{array}{ccccccc}
 \cdots & \xrightarrow{\partial_{d+2}} & C_{d+1}(X) & \xrightarrow{\partial_{d+1}} & C_d(X) & \xrightarrow{\partial_d} & C_{d-1}(X) \xrightarrow{\partial_{d-1}} \cdots \\
 & & \downarrow f_{d+1} & & \downarrow f_d & & \downarrow f_{d-1} \\
 \cdots & \xrightarrow{\partial_{d+2}} & C_{d+1}(Y) & \xrightarrow{\partial_{d+1}} & C_d(Y) & \xrightarrow{\partial_d} & C_{d-1}(Y) \xrightarrow{\partial_{d-1}} \cdots
 \end{array} \tag{2.80}$$

The chain map  $f_d$  gives rise to a homomorphism  $f_* : H_d(X) \rightarrow H_d(Y)$ . It becomes evident that if  $X$  and  $Y$  are homeomorphic, meaning there exists a homeomorphism  $f : X \rightarrow Y$ , then the induced map  $f_*$  is an isomorphism.

To formalize the relationships between the homology groups of a topological space  $X$ , a subset  $A \subset X$ , and the quotient space  $X/A$ , we introduce the concept of exact sequences.

**Definition 2.3.7 (Exact Sequences)** *An arrangement of elements in the form*

$$\cdots \rightarrow A_{d+1} \xrightarrow{\alpha_{d+1}} A_d \xrightarrow{\alpha_d} A_{d-1} \rightarrow \cdots \tag{2.81}$$

*is referred to as an exact sequence when the  $A_i$  are abelian groups and the  $\alpha_i$  are homomorphisms, and it satisfies the condition that  $\ker(\alpha_d) = \text{im}(\alpha_{d+1})$  for all  $d$ .*

**Remark 2.3.8**

- The condition  $\ker(\alpha_d) = \text{im}(\alpha_{d+1})$  implies that  $\text{im}(\alpha_{d+1})$  is a subset of  $\ker(\alpha_d)$ , which is equivalent to  $\alpha_d \circ \alpha_{d+1} = 0$ . Therefore, an exact sequence can be seen as a chain complex.
- Since  $\ker(\alpha_d)$  is a subset of  $\text{im}(\alpha_{d+1})$ , the homology groups of an exact sequence are trivial.

**Proposition 2.3.4** [10, §1.3.1] We can establish the following equivalences:

1.  $0 \rightarrow A \xrightarrow{a} B$  is exact  $\iff \ker(a) = 0$ , or  $a$  is injective.
2.  $A \xrightarrow{a} B \rightarrow 0$  is exact  $\iff \text{im}(a) = B$ , or  $a$  is surjective.
3.  $0 \rightarrow A \xrightarrow{a} B \xrightarrow{0} 0$  is exact if and only if  $a$  is an isomorphism.
4. A sequence of the form

$$0 \rightarrow A \xrightarrow{a} B \rightarrow 0 \tag{2.82}$$

is said to be exact if and only if the following conditions hold:

- The map  $a : A \rightarrow B$  is injective, meaning that  $\ker(a) = 0$ .
- The map  $b : B \rightarrow 0$  is surjective, meaning that  $\text{im}(b) = 0$ .

- The kernel of  $b$  is equal to the image of  $a$ , i.e.,  $\ker(b) = \text{im}(a)$ .
- In this case, the map  $b$  induces an isomorphism  $C \cong B/\text{im}(a)$ , where  $C$  is the quotient of  $B$  by the image of  $a$ .

If the map  $a : A \hookrightarrow B$  is an inclusion, then  $C \cong B/A$ , where  $B/A$  denotes the quotient of  $B$  by the subgroup  $A$ . This type of exact sequence is commonly referred to as a short exact sequence.

## Proof

1. Assume  $0 \rightarrow A \xrightarrow{a} B$  is exact. This means  $\text{im}(0) = \ker(a)$ , which implies  $\ker(a) = 0$  since the image of the zero map is always the trivial group. Therefore,  $a$  is injective.  
Conversely, suppose  $\ker(a) = 0$ . We need to show  $\text{im}(0) = \ker(a)$ . Since  $\ker(a) = 0$ , the only element mapped to the identity in  $B$  is the zero element of  $A$ . Thus, the sequence  $0 \rightarrow A \xrightarrow{a} B$  is exact.
2. Assume  $A \xrightarrow{a} B \rightarrow 0$  is exact. This means  $\text{im}(a) = \ker(0)$ , which implies  $\text{im}(a) = B$  since the kernel of the zero map is always the entire group. Therefore,  $a$  is surjective.  
Conversely, suppose  $\text{im}(a) = B$ . We need to show  $\text{im}(a) = \ker(0)$ . Since  $\text{im}(a) = B$ , every element in  $B$  has a preimage in  $A$  under the map  $a$ . Thus, the sequence  $A \xrightarrow{a} B \rightarrow 0$  is exact.
3. Assume  $0 \rightarrow A \xrightarrow{a} B \rightarrow 0$  is exact. From the exactness, we have  $\ker(a) = \text{im}(0) = 0$  and  $\text{im}(a) = \ker(0) = B$ . Therefore,  $a$  is both injective and surjective, hence an isomorphism.  
Conversely, suppose  $a$  is an isomorphism, meaning  $a$  is both injective (no kernel) and surjective (maps onto  $B$ ). Thus,  $0 \rightarrow A \xrightarrow{a} B \rightarrow 0$  is exact, satisfying  $\text{im}(0) = \ker(a)$  and  $\text{im}(a) = \ker(0)$ .
4. Assume the sequence  $0 \rightarrow A \xrightarrow{a} B \rightarrow 0$  is exact.  $a$  is injective, and  $\text{im}(a) = \ker(b)$  where  $b$  maps  $B$  to  $0$ . As  $b$  is the zero map and surjective,  $B/\text{im}(a) = 0$ , implying  $B = \text{im}(a)$ . Therefore,  $a$  is an isomorphism, establishing  $B \cong A$ .  
Conversely, if  $a$  is an isomorphism,  $\ker(a) = 0$  and  $\text{im}(a) = B$ . Therefore, the sequence  $0 \rightarrow A \xrightarrow{a} B \rightarrow 0$  is exact by definition.

□

## 2.4 Relative Homology

The concept we will now discuss is that of relative homology groups. Let  $X$  be a topological space and  $A$  a subspace of  $X$ . We define  $C_d(X, A)$  as the quotient group  $C_d(X)/C_d(A)$ . This means that chains in  $A$  are considered equivalent to the trivial chains in  $C_d(X)$ .

Since the boundary operator  $\partial_d : C_d(X) \rightarrow C_{d-1}(X)$  also maps  $C_d(A)$  to  $C_{d-1}(A)$ , a natural boundary map on the quotient group  $\partial_d : C_d(X, A) \rightarrow C_{d-1}(X, A)$  is obtained. This gives rise to the following sequence:

$$\cdots \rightarrow C_{d+1}(X, A) \xrightarrow{\partial_{d+1}} C_d(X, A) \xrightarrow{\partial_d} C_{d-1}(X, A) \rightarrow \cdots \quad (2.83)$$

This sequence forms a chain complex because  $\partial_{d+1} \circ \partial_d = 0$ . We can then define the relative homology groups  $H_d(X, A)$  as the homology groups of this chain complex.

We propose two important facts about  $H_d(X, A)$ :

### Proposition 2.4.1 [9, §2.1]

1. Elements in  $H_d(X, A)$  are represented by relative cycles, which are  $d$ -chains  $c$  in  $C_d(X)$  such that  $\partial_d(c) \in C_{d-1}(A)$ .
2. A relative cycle  $c$  is trivial in  $H_d(X, A)$  if and only if it is a relative boundary, i.e.,  $c$  is the sum of a chain in  $C_d(A)$  and the boundary of a chain in  $C_{d+1}(X)$ .

### Proof

1. Assume  $[c] \in H_d(X, A)$  represents a homology class of a chain  $c \in C_d(X)$ . Since  $[c]$  is a class in the relative homology group,  $\partial_d(c) \in C_{d-1}(A)$ , implying that  $c$  is a relative cycle because its boundary maps into the subspace  $A$ .

Conversely, if  $\partial_d(c) \in C_{d-1}(A)$ , then  $c$  qualifies as a relative cycle by definition, and any chain homologous to  $c$  in  $C_d(X)$  that differs from  $c$  by a boundary in  $C_d(A)$  will also have its boundary in  $C_{d-1}(A)$ , confirming  $[c]$  as an element of  $H_d(X, A)$ .

2. For a chain  $c$  in  $C_d(X)$  to be a relative boundary, it must be expressible as  $c = a + \partial_{d+1}(b)$  where  $a \in C_d(A)$  and  $b \in C_{d+1}(X)$ . Applying the boundary operator, we have:

$$\partial_d(c) = \partial_d(a) + \partial_d(\partial_{d+1}(b)) = \partial_d(a) + 0 = \partial_d(a).$$

Since  $\partial_d(a) \in C_{d-1}(A)$  and  $\partial_{d+1}\partial_d = 0$  (as boundaries of boundaries are zero),  $c$  is a cycle relative to  $A$ , making it trivial in  $H_d(X, A)$ .

Conversely, if a relative cycle  $c$  is trivial in  $H_d(X, A)$ , then it must be homologous to a boundary in  $C_d(A)$ , meaning there exists  $a \in C_d(A)$  and  $b \in C_{d+1}(X)$  such that  $c = a + \partial_{d+1}(b)$ . This implies that  $c$  is expressible as the sum of a chain in  $C_d(A)$  and the boundary of a chain in  $C_{d+1}(X)$ , confirming it as a relative boundary.

Thus, a relative cycle is trivial in  $H_d(X, A)$  if and only if it is a relative boundary.  $\square$

Before we further investigate the decomposition of the relative homology groups into exact sequences, we need an intermediate result for commutative diagrams, which facilitates further argumentation. This intermediate result is also known as the Snake Lemma.

**Lemma 2.4.2 (Snake Lemma)** [10, §1.2.6] Consider the following commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A' & \xrightarrow{f'} & A & \xrightarrow{f} & A'' & \longrightarrow & 0 \\ & & \downarrow a' & & \downarrow a & & \downarrow a'' & & \\ 0 & \longrightarrow & B' & \xrightarrow{g'} & B & \xrightarrow{g} & B'' & \longrightarrow & 0 \end{array} \quad (2.84)$$

There exists an exact sequence:

$$\ker(a') \xrightarrow{\delta} \ker(a) \xrightarrow{\delta} \ker(a'') \xrightarrow{\delta} \dots \quad (2.85)$$

$$\dots \xrightarrow{\delta} \operatorname{coker}(a') \xrightarrow{\delta} \operatorname{coker}(a) \xrightarrow{\delta} \operatorname{coker}(a'') \quad (2.86)$$

with the connecting homomorphism  $\delta$ .

**Proof** Consider  $x'' \in \ker(a'') \subseteq A''$ . Since  $f$  is surjective, there exists  $x \in A$  such that  $f(x) = x''$ . We have  $a''(x'') = 0$ , hence  $a''(f(x)) = 0$ . By commutativity,  $g(a(x)) = 0$ , so  $a(x) \in \ker(g) = \operatorname{im}(g')$ . Thus, there exists  $y' \in B'$  such that  $g'(y') = a(x)$ . Set  $\delta(x'') = \operatorname{cl}(y') \in \operatorname{coker}(a')$ .

If  $x'' = f(x'_1)$ , choose  $x$  such that  $f(x) = x''$ , then  $a(x) = g'(y'_1)$  and  $\delta(x'') = \operatorname{cl}(y'_1)$ . If  $x = x_2 + f(x_1)$ , then  $a(x) = a(x_2) + a(f(x_1))$ , with  $g(a(x_1)) = 0$  and  $a(f(x_1)) \in \operatorname{im}(g')$ , so  $\delta$  is independent of the choice of  $x$ . Linear maps remain linear under  $\delta$ .



Suppose  $y'' \in \ker(\delta)$ . This means  $y'' \in \ker(a'')$  and  $\delta(y'') = 0$ . There exists  $x \in A$  such that  $f(x) = y''$  and  $a(x) = g'(z')$  for some  $z' \in B'$ . Since  $\delta(y'') = \text{cl}(z') = 0$ , we have  $z' \in \text{im}(a')$ , so  $z' = a'(w')$  for some  $w' \in A'$ . Hence,  $a(x) = g'(a'(w')) = a(g(w'))$ , so  $x - g(w') \in \ker(a)$ . Therefore,  $y'' = f(x) = f(x - g(w')) \in \text{im}(f)$ , showing that  $\ker(\delta) = \text{im}(\ker(a) \rightarrow \ker(a''))$ .

Suppose  $y \in \text{coker}(a)$  such that  $y \in \ker(\text{coker}(a) \rightarrow \text{coker}(a''))$ . This means  $y = \text{cl}(z)$  for some  $z \in B$  with  $a(z) = g'(w')$  for some  $w' \in B'$ . Since  $z \in \ker(g)$ , we have  $z = g(x)$  for some  $x \in A$ . Thus,  $y = \text{cl}(z) = \text{cl}(g(x)) = 0$ , showing exactness.  $\square$

**Theorem 2.4.3** [9, p.115ff] The relative homology groups  $H_d(X, A)$  are part of the exact sequence:

$$\cdots \rightarrow H_d(A) \rightarrow H_d(X) \rightarrow H_d(X, A) \rightarrow H_{d-1}(A) \rightarrow \cdots \quad (2.87)$$

$$\cdots \rightarrow H_0(X, A) \rightarrow 0. \quad (2.88)$$

**Proof** Consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_d(A) & \xhookrightarrow{i} & C_d(X) & \twoheadrightarrow{j} & C_d(X, A) \longrightarrow 0 \\ & & \downarrow \partial_d & & \downarrow \partial_d & & \downarrow \partial_d \\ 0 & \longrightarrow & C_{d-1}(A) & \xhookrightarrow{i} & C_{d-1}(X) & \twoheadrightarrow{j} & C_{d-1}(X, A) \longrightarrow 0. \end{array} \quad (2.89)$$

Here,  $i$  is the inclusion map  $C_d(A) \hookrightarrow C_d(X)$ , and  $j$  is the quotient map  $C_d(X) \twoheadrightarrow C_d(X, A)$ . Both rows are exact, and the diagram commutes. This gives rise to a long exact sequence of homology groups by the Snake Lemma 2.4.2.

To clarify, consider the chain complexes  $A_\bullet = \{C_d(A), \partial_d\}_{d \in \mathbb{N}}$ ,  $B_\bullet = \{C_d(X), \partial_d\}_{d \in \mathbb{N}}$ , and  $C_\bullet = \{C_d(X, A), \partial_d\}_{d \in \mathbb{N}}$ .

The vertical maps  $i$  and  $j$  are chain maps and thus induce homomorphisms on homology groups  $i_\star : H_d(A) \rightarrow H_d(X)$  and  $j_\star : H_d(X) \rightarrow H_d(X, A)$ . The exactness of the rows implies that  $\ker(j_\star) = \text{im}(i_\star)$ .

The connecting homomorphism  $\partial : H_d(X, A) \rightarrow H_{d-1}(A)$  can be defined as follows: for any class  $[c] \in H_d(X, A)$  represented by a cycle  $c \in C_d(X)$  with  $\partial_d(c) \in C_{d-1}(A)$ , choose a chain  $b \in C_d(X)$  such that  $j(b) = c$ . Since  $c$  is a cycle,  $\partial_d(b) \in \ker(j) = \text{im}(i)$ , so there exists a chain  $a \in C_{d-1}(A)$  such that  $\partial_d(b) = i(a)$ . Define  $\partial([c]) = [a] \in H_{d-1}(A)$ .

We need to verify that the map  $\partial$  is well-defined:

- Uniqueness: If  $b$  and  $b'$  both map to  $c$  via  $j$ , then  $b - b' \in \ker(j) = \text{im}(i)$ . Thus, there exists  $a' \in C_{d-1}(A)$  such that  $b - b' = i(a')$ . Hence,

$$\partial_d(b) = \partial_d(b') + \partial_d(i(a')) = \partial_d(b') + i(\partial_{d-1}(a')). \quad (2.90)$$

So,  $[a] = [a']$  in  $H_{d-1}(A)$ , ensuring the uniqueness of  $\partial([c])$ .

- Homologous chains: If  $c$  and  $c'$  are homologous in  $H_d(X, A)$ , then  $c - c' = \partial_d(c'')$  for some  $c'' \in C_{d+1}(X, A)$ . Thus,  $b - b' = \partial_{d+1}(c'')$  in  $C_d(X)$ , and the same argument as above applies.

This establishes the long exact sequence:

$$\cdots \rightarrow H_d(A) \rightarrow H_d(X) \rightarrow H_d(X, A) \rightarrow H_{d-1}(A) \rightarrow \cdots \quad (2.91)$$

$$\cdots \rightarrow H_0(X, A) \rightarrow 0. \quad (2.92)$$

□

**Proposition 2.4.4** *The map  $\partial_d : H_d(C) \rightarrow H_{d-1}(A)$  is a homomorphism.*

**Proof** Let  $[c_1], [c_2] \in H_d(C)$  be two homology classes, where  $c_1, c_2 \in C_d(X, A)$  are cycles. By definition, let  $[a_1] = \partial_d([c_1])$  and  $[a_2] = \partial_d([c_2])$ . This means that  $c_1$  and  $c_2$  are represented by chains  $b_1, b_2 \in C_d(X)$  such that  $j(b_1) = c_1$  and  $j(b_2) = c_2$ .

Since  $b_1$  and  $b_2$  are cycles modulo  $A$ , we have:

$$\partial_d(b_1) = i(a_1) \quad \text{and} \quad \partial_d(b_2) = i(a_2). \quad (2.93)$$

We need to show that:

$$\partial_d([c_1] + [c_2]) = [a_1] + [a_2]. \quad (2.94)$$

Consider the sum  $c_1 + c_2 \in C_d(X, A)$ . By the properties of the quotient map  $j$  we yield:

$$j(b_1 + b_2) = j(b_1) + j(b_2) = c_1 + c_2. \quad (2.95)$$

Thus,  $b_1 + b_2 \in C_d(X)$  is a preimage of  $c_1 + c_2$  under the quotient map  $j$ . Furthermore, the boundary of  $b_1 + b_2$  is:

$$\partial_d(b_1 + b_2) = \partial_d(b_1) + \partial_d(b_2) = i(a_1) + i(a_2) = i(a_1 + a_2). \quad (2.96)$$

Therefore, the cycle  $c_1 + c_2$  maps to the cycle  $a_1 + a_2$  under  $\partial_d$ , which implies on the respective equivalence classes:

$$\partial_d([c_1] + [c_2]) = [a_1 + a_2] = [a_1] + [a_2]. \quad (2.97)$$

Thus,  $\partial_d$  is a homomorphism.  $\square$

**Lemma 2.4.5** *The given sequence,*

$$\cdots \rightarrow H_d(A) \xrightarrow{i_\star} H_d(B) \xrightarrow{j_\star} H_d(C) \xrightarrow{\partial_d} H_{d-1}(A) \xrightarrow{i_\star} H_{d-1}(B) \rightarrow \cdots \quad (2.98)$$

*is exact.*

**Proof** There are six inclusions that need to be confirmed:

1.  $\text{im}(i_\star) \subseteq \ker(j_\star)$ : Since  $j \circ i = 0$ , it follows that  $j_\star \circ i_\star = 0$ , so  $\text{im}(i_\star) \subseteq \ker(j_\star)$ .
2.  $\text{im}(j_\star) \subseteq \ker(\partial_d)$ : By definition, if  $[b] \in \text{im}(j_\star)$ , then  $j_\star([b]) = [c] \in H_d(C)$  where  $c$  is a cycle. Therefore,  $\partial_d([c]) = 0$ , and thus  $\text{im}(j_\star) \subseteq \ker(\partial_d)$ .
3.  $\text{im}(\partial_d) \subseteq \ker(i_\star)$ : If  $[c] \in \text{im}(\partial_d)$ , then  $c$  is a relative boundary. This implies that there exists  $b \in B_d$  such that  $c = j(b)$ . Since  $\partial_d(b) = 0$ , it follows that  $i_\star([c]) = 0$ , thus  $\text{im}(\partial_d) \subseteq \ker(i_\star)$ .
4.  $\ker(j_\star) \subseteq \text{im}(i_\star)$ : Let  $[b] \in \ker(j_\star)$ , meaning  $j_\star([b]) = [c] = 0$  in  $H_d(C)$ . This implies that  $c$  is a boundary, i.e., there exists a chain  $c' \in C_{d+1}$  such that  $c = \partial_{d+1}(c')$ . Since  $j$  is surjective,  $c' = j(b')$  for some  $b' \in B_{d+1}$ . Thus,  $j(b) = \partial_{d+1}(c') = \partial_{d+1} \circ j(b')$ , which leads to:

$$j(b - \partial_{d+1}(b')) = 0.$$

Hence,  $b - \partial_{d+1}(b') = i(a)$  for some  $a \in A_d$ . Since  $i$  is injective,  $a$  is a cycle because:

$$i \circ \partial_d(a) = \partial_d \circ i(a) = \partial_d(b - \partial_{d+1}(b')) = \partial_d(b) = 0,$$

given that  $b$  is a cycle. Thus,  $i_\star([a]) = [b]$ , and  $\text{im}(i_\star) = \ker(j_\star)$ .

5.  $\ker(\partial_d) \subseteq \text{im}(j_\star)$ : Let  $[c] \in \ker(\partial_d)$ . This means that  $c$  is a relative cycle such that  $\partial_d([c]) = 0$ . Hence, there exists  $b \in B_d$  such that  $j(b) = c$ . Therefore,  $\ker(\partial_d) \subseteq \text{im}(j_\star)$ .

6.  $\ker(i_\star) \subseteq \operatorname{im}(\partial)$ : Let  $[a] \in \ker(i_\star)$ , meaning  $i_\star([a]) = 0$  in  $H_{d-1}(B)$ . Thus,  $i(a) = \partial_d(b)$  for some  $b \in B_d$ . Since  $\partial_d(j(b)) = j(\partial_d(b)) = j \circ i(a) = 0$ ,  $j(b)$  is a cycle. Therefore,  $\partial_d([j(b)]) = [a]$ , which demonstrates that  $\ker(i_\star) \subseteq \operatorname{im}(\partial_d)$ .

We have established the following relationships:

- $\operatorname{im}(i_\star) = \ker(j_\star)$ ,
- $\operatorname{im}(j_\star) = \ker(\partial_d)$ ,
- $\operatorname{im}(\partial_d) = \ker(i_\star)$ .

These relationships confirm that the sequence is exact, satisfying all the necessary conditions.  $\square$

**Proposition 2.4.6** *The sequence*

$$\cdots \rightarrow H_d(A) \xrightarrow{i_\star} H_d(X) \xrightarrow{j_\star} H_d(X, A) \xrightarrow{\partial_d} \cdots \quad (2.99)$$

$$\cdots \xrightarrow{\partial_d} H_{d-1}(A) \rightarrow \cdots \rightarrow H_0(X, A) \rightarrow 0 \quad (2.100)$$

*is exact.*

**Proof** The exactness of this sequence is a standard result in algebraic topology, which follows from the long exact sequence of the pair  $(X, A)$ . The maps in the sequence are defined as follows:

- $i_\star$  is induced by the inclusion map  $i : A \hookrightarrow X$ .
- $j_\star$  is induced by the quotient map  $j : X \rightarrow X/A$ , where each chain in  $X$  is mapped to its homology class in  $X/A$  modulo the image of chains in  $A$ .
- $\partial_d$  is the boundary map connecting homology groups, defined by taking the boundary of a relative cycle in  $H_d(X, A)$ , which by definition of a chain complex is a cycle in  $A$  at one dimension lower.

The exactness at each position, for instance at  $H_d(X, A)$ , implies that the image of  $j_\star$  from  $H_d(X)$  is equal to the kernel of  $\partial_d$ . This shows that any cycle in  $H_d(X, A)$  that becomes trivial in  $H_{d-1}(A)$  must come from a cycle in  $X$  that is not affected by cycles in  $A$ . The exactness at  $H_{d-1}(A)$  further implies that the image of  $\partial_d$  is exactly the kernel of  $i_\star$ , which matches cycles in  $A$  that bound in  $X$  but not in  $A$  itself.

Therefore, the exactness of the entire sequence follows from the properties of the chain maps and the boundary operators defined in the chain complexes of  $A$ ,  $X$ , and  $X/A$ . The additional observation that  $\partial_d([c]) = [\partial_d(c)]$  for any relative cycle  $c \in H_d(X, A)$  reinforces the continuity of the exact

sequence through the boundary mapping, as it relates the homology in one dimension in  $A$  to the relative homology in  $X$  and  $A$ .  $\square$

## 2.5 The Excision Theorem

Additionally, we refer to the Excision Theorem, a foundational result in algebraic topology. Simply put, the Excision Theorem allows for the analysis of the homology of a space by effectively "excising", or removing, a smaller subspace, under specific topological conditions. These conditions usually involve the smaller subspace being "negligible" in a certain sense, such as being contained in the interior of another subspace. Essentially, this theorem guarantees that, given these conditions, the homology of the original space is preserved when compared to the homology of the space with the smaller subspace removed.

We will need the famous five lemma to prove the Excision Theorem.

**Lemma 2.5.1 (The Five Lemma)** [9, p.120] Consider a commutative diagram with exact rows:

$$\begin{array}{ccccccccc}
 A & \xrightarrow{i} & B & \xrightarrow{j} & C & \xrightarrow{k} & D & \xrightarrow{l} & E \\
 \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta & & \downarrow \epsilon \\
 A' & \xrightarrow{i'} & B' & \xrightarrow{j'} & C' & \xrightarrow{k'} & D' & \xrightarrow{l'} & E'
 \end{array} \tag{2.101}$$

If  $\alpha, \beta, \delta, \epsilon$  are isomorphisms, then  $\gamma$  is also an isomorphism.

**Proof** Commutativity of the diagram ensures that  $\gamma$  is a homomorphism. To show that  $\gamma$  is bijective, we consider:

- **Surjectivity:** Let  $c' \in C'$ . Since  $\delta$  is surjective, there exists  $d \in D$  such that  $k'(c') = \delta(d)$ . Because  $\epsilon$  is injective,  $\epsilon(l(d)) = l'(\delta(d)) = l'(k'(c')) = 0$ , thus  $l(d) = 0$ . By exactness,  $d = k(c)$  for some  $c \in C$ . Therefore,  $k'(c' - \gamma(c)) = 0$ , and exactness implies  $c' - \gamma(c) = j'(b')$  for some  $b' \in B'$ . With  $\beta$  surjective,  $b' = \beta(b)$  for some  $b \in B$ , hence  $\gamma(c + j(b)) = c'$ , proving surjectivity of  $\gamma$ .
- **Injectivity:** Assume  $\gamma(c) = 0$ . Injectivity of  $\delta$  and exactness yield  $\delta(k(c)) = 0$  and  $k(c) = 0$ , thus  $c = j(b)$  for some  $b \in B$ . Since  $\gamma(j(b)) = j'(\beta(b)) = 0$  and  $\beta$  is injective,  $b = i(a)$  for some  $a \in A$ ,

giving  $c = j(i(a)) = 0$ . Therefore,  $\gamma$  has a trivial kernel and is injective.

Thus,  $\gamma$  is both surjective and injective, proving it is an isomorphism.

□

**Theorem 2.5.2 (Excision Theorem)** [9, §2.20] *Given subspaces  $Z \subset A \subset X$  such that  $\overline{Z} \subseteq \text{int}(A)$ , the inclusion  $(X \setminus Z, A \setminus Z) \hookrightarrow (X, A)$  induces isomorphisms in homology:*

$$H_n(X \setminus Z, A \setminus Z) \rightarrow H_n(X, A) \quad \text{for all } n.$$

Equivalently, for subspaces  $A, B \subset X$  whose interiors cover  $X$ , the inclusion  $(B, A \cap B) \hookrightarrow (X, A)$  induces isomorphisms:

$$H_n(B, A \cap B) \rightarrow H_n(X, A) \quad \text{for all } n.$$

The equivalence is demonstrated by setting  $B = X \setminus Z$  and  $Z = X \setminus B$ , thereby  $A \cap B = A \setminus Z$ . The condition  $\overline{Z} \subseteq \text{int}(A)$  translates to  $X = \text{int}(A) \cup \text{int}(B)$ . Following Hatcher's proof, it involves barycentric subdivision, which aids in the computation of homology groups using small singular simplices. In metric spaces, 'smallness' is determined by the diameters of simplices, while in general topological spaces, it is defined by their containment within elements of a cover  $\mathcal{U}$ . Define  $\mathcal{U} = \{U_j\}$ , an open cover of  $X$ , and let  $C_n^{\mathcal{U}}(X)$  be the subgroup of  $C_n(X)$  consisting of chains  $\sum_i n_i \sigma_i$  with each  $\sigma_i$  contained within one of the  $U_j$ . The boundary operator  $\partial : C_n(X) \rightarrow C_{n-1}(X)$  preserves this containment, forming a chain complex  $C_n^{\mathcal{U}}(X) \rightarrow C_{n-1}^{\mathcal{U}}(X)$ . The homology groups of this complex are denoted by  $H_n^{\mathcal{U}}(X)$ .

**Proposition 2.5.3** *According to [9, §2.21], the inclusion  $\iota : C_n^{\mathcal{U}}(X) \hookrightarrow C_n(X)$  is a chain homotopy equivalence. There exists a chain map  $\rho : C_n(X) \rightarrow C_n^{\mathcal{U}}(X)$  such that the compositions  $\iota \circ \rho$  and  $\rho \circ \iota$  are chain homotopic to the identity, implying that  $\iota$  induces isomorphisms  $H_n^{\mathcal{U}}(X) \cong H_n(X)$  for all  $n$ .*

**Proof** TBD

□

Finally, we can prove the Excision Theorem.

**Proof** We use the setup and notation previously established, where  $U$  is a subset of  $B$  and  $\overline{U} \subseteq \text{int}(A)$ . Define  $C_n(A + B) \equiv C_n^{\mathcal{U}}(X)$  which sums chains in  $A$  and chains in  $B$ . We consider the following key steps:

In our setup, the formulas:

$$\partial D + D\partial = \text{id} - \iota \circ \rho \quad \text{and} \quad \rho \circ \iota = \text{id}$$

establish that all maps take chains in  $A$  to chains in  $A$ , which allows us to induce quotient maps when factoring out chains solely in  $A$ . These quotient maps automatically satisfy the same two formulas, thereby ensuring that the inclusion map  $C_n(A + B)/C_n(A) \rightarrow C_n(X)/C_n(A)$  is an isomorphism on homology.

The inclusion  $C_n(B)/C_n(A \cap B) \rightarrow C_n(A + B)/C_n(A)$  is also an isomorphism because both quotient groups are free and have a basis represented by singular  $n$ -simplices in  $B$  that are not in  $A$ . This property ensures that we have a one-to-one correspondence between the generators of both groups.

With these isomorphisms established, we apply the excision property to show that the inclusion  $i : (X \setminus U, A \setminus U) \hookrightarrow (X, A)$  induces an isomorphism:

$$H_n(X \setminus U, A \setminus U) \cong H_n(X, A)$$

for each  $n$ . This follows because the quotient spaces  $(X \setminus U, A \setminus U)$  and  $(X, A)$  relative to  $A$  share the same homological structure due to the isomorphisms we've established in the quotient chain groups and by the preservation of homotopy relations.

Thus, we conclude that the excision property holds under the given conditions, confirming the fundamental theorem of algebraic topology in these settings.  $\square$

## 2.6 Equivalence of $H_d^{\Delta}(\mathbf{X})$ and $H_d(\mathbf{X})$

We aim to establish the equivalence of the homology groups  $H_d^{\Delta}(X)$  and  $H_d(X)$ . Note that simplicial homology groups  $H_d^{\Delta}(X)$  are defined solely for simplicial complexes, whereas singular homology groups  $H_d(X)$  can be computed for any topological space. This generality is advantageous, as homeomorphic spaces have isomorphic singular homology groups, enabling the imposition of a simplicial structure on any triangulable topological space. For the scope of this thesis, we restrict our attention to spaces representable as simplicial complexes.

To demonstrate the equivalence between  $H_d(X)$  and  $H_d^\Delta(X)$ , we establish an isomorphism between these groups for all dimensions  $d$ . A homomorphism naturally arises from the map  $C_d^\Delta(X) \rightarrow C_d(X)$ , which sends each  $d$ -simplex in  $X$  to a singular simplex via the mapping  $\tilde{\sigma}^{(d)} : \sigma^{(d)} \rightarrow X$ . This induces a homomorphism from  $H_d^\Delta(X)$  to  $H_d(X)$ .

While simplicial and singular homologies originate from similar foundational ideas and are particularly useful in different contexts—simplicial homology in computational geometry and singular homology in proving theorems involving continuous maps—their equivalence is a significant theoretical result. This equivalence not only bridges computational and conceptual approaches in algebraic topology but also provides a robust framework for addressing a broad spectrum of mathematical challenges.

**Theorem 2.6.1 (Equivalence of  $H_d^\Delta(X)$  and  $H_d(X)$ )** [9, p.102f] For each integer  $d$ , the homomorphisms from the simplicial homology group  $H_d^\Delta(X)$  to the singular homology group  $H_d(X)$  are isomorphisms. Hence, the simplicial and singular homology groups are equivalent for all dimensions.

**Proof** Consider a simplicial complex  $X$ . For each  $k$ -skeleton  $X^k$ , the inclusion  $X^{k-1} \subset X^k$  leads to the following commutative diagram of exact sequences:

$$\begin{array}{ccccccccc} H_{d+1}^\Delta(X^k, X^{k-1}) & \longrightarrow & H_d^\Delta(X^{k-1}) & \longrightarrow & H_d^\Delta(X^k) & \longrightarrow & H_d^\Delta(X^k, X^{k-1}) & \longrightarrow & H_{d-1}^\Delta(X^{k-1}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H_{d+1}(X^k, X^{k-1}) & \longrightarrow & H_d(X^{k-1}) & \longrightarrow & H_d(X^k) & \longrightarrow & H_d(X^k, X^{k-1}) & \longrightarrow & H_{d-1}(X^{k-1}) \end{array}$$

Figure 2.1: Commutative diagram of exact sequences for homology and relative homology of a triangulable topological space  $X$ .

Given that  $X^k/X^{k-1}$  only contains simplices of dimension  $k$ ,  $C_d(X^k, X^{k-1})$  is trivial for  $d \neq k$  and a free abelian group for  $d = k$ , with a basis formed by the  $k$ -simplices of  $X$ .

We define a characteristic map  $\Phi : \coprod_i (\sigma_i^{(k)}, \sigma_i^{(k-1)}) \rightarrow (X^k, X^{k-1})$ , inducing a homeomorphism  $\Phi_* : \coprod_i \sigma_i^{(k)} / \coprod_i \sigma_i^{(k-1)} \rightarrow X^k/X^{k-1}$ . By the Excision Theorem 2.5.2,  $H_d(\coprod_i (\sigma_i^{(k)}, \sigma_i^{(k-1)})) \cong H_d(X^k, X^{k-1}) \cong H_d(X^k/X^{k-1})$ , establishing isomorphisms for  $H_d(X^k, X^{k-1})$ .

Using induction and assuming that the necessary homology isomorphisms hold for dimensions less than  $k$ , we extend the isomorphism to all



dimensions:

$$H_k^\Delta(X^k, X^{k-1}) \cong H_k(X^k, X^{k-1}) \quad (2.102)$$

The Five Lemma 2.5.1 supports the isomorphism between  $H_k^\Delta(X^k, X^{k-1})$  and  $H_k(X^k, X^{k-1})$ . Thus, we conclude that  $H_d^\Delta(X^k, X^{k-1}) \rightarrow H_d(X^k, X^{k-1})$  is an isomorphism for all  $d$ , verifying the equivalence of simplicial and singular homology for all  $k$ .  $\square$

# Chapter 3

## Homological Persistence

### 3.1 Persistent Homology

The core of this chapter focuses on persistent homology and illustrates how it identifies multiscale topological features using filtrations of complexes upon point clouds. We discuss barcode isomorphisms and persistence modules that visualize and explain the stability of the detected features within a filtration to perturbations of the data. The chapter concludes with persistent chain complexes and the cohomology of these structures, which enhance our understanding of the evolution of the topology of filtrations on point clouds. We also investigate distances between persistence modules and the stability theorem to provide metrics for comparing topological features and ensure the statistical robustness of persistent homology.

#### 3.1.1 Filtrations of Complexes

Individual (co)homology groups are consistently defined with coefficients in a fixed field  $\mathbb{F}$ , organizing persistent (co)homology as a graded module over  $\mathbb{F}[x]$ . Replacing  $\mathbb{F}$  with the ring  $\mathbb{Z}$  leads to substantial issues, but most of the results still hold for principal ideal domains, as discussed in [11, §3.1].

We investigate the persistent topology of filtered topological spaces, with a primary focus on the prototypical example of a filtered cell complex. This structure is defined by a sequence  $\mathcal{X}$  of cell complexes:

$$\mathcal{X} : \emptyset \subset X_1 \subset X_2 \subset \cdots \subset X_n = X_\infty, \quad (3.1)$$

where  $X_1$  starts with a single vertex  $\sigma_1$ , and each subsequent complex  $X_i$  is

constructed by adding a single cell to the previous complex:

$$X_i := X_{i-1} \cup \sigma_i. \quad (3.2)$$

Here, the indexing set is  $\{1, 2, \dots, n\}$ . Additionally, associated real values  $a_i$  are assigned to these indices, satisfying  $a_1 \leq a_2 \leq \dots \leq a_n$ . This formulation clearly delineates the stepwise enlargement of the complex, illustrating the dynamic evolution of its topology as new cells are incrementally incorporated.

We will use the same running example as in [6, §2.2].

**Example 3.1.1** *The chosen illustrative example involves a cellular filtration of the 2-sphere, denoted by  $S^2$ . This process constructs a cell complex and introduces an ordering among the cells to facilitate differentiation between them. The development of the filtration can be articulated as follows [6, §2.2]:*

$$\begin{aligned} S^2 : & \emptyset \\ & \subset S_1 = \{\sigma_1\} \\ & \subset S_2 = \{\sigma_1, \sigma_2\} \\ & \subset S_3 = \{\sigma_1, \sigma_2, \sigma_3 := (\sigma_1, \sigma_2)\} \\ & \subset S_4 = \{\sigma_1, \sigma_2, \sigma_3, \sigma_4 := (\sigma_2, \sigma_1)\} \\ & \subset S_5 = \{\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5 := (\sigma_3, \sigma_4)\} \\ & \subset S_6 = \{\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5, \sigma_6 := (\sigma_4, \sigma_3)\}. \end{aligned}$$

*In this sequence, the cell  $\sigma_1$  symbolizes the initial point. The set  $S_2$  incorporates two distinct points. In  $S_3$ , a path connecting  $\sigma_1$  and  $\sigma_2$  is introduced.  $S_4$  augments this path with its reverse from  $\sigma_2$  to  $\sigma_1$  clearly distinguishing it from  $\sigma_3$ . Finally,  $S_5$  and  $S_6$  progressively differentiate between cells that represent the upper and lower halves of the sphere, respectively.*

### 3.1.2 Persistence Modules

#### Interval-Decomposable Modules

#### Interval-Indecomposable Modules

### 3.1.3 Persistent Homology on Complexes

In the context of algebraic topology, applying the homology functor  $H$  to a filtration of a complex  $\mathcal{X}$  yields a sequence of algebraic structures:

$$H(\mathcal{X}) : H(X_1) \rightarrow H(X_2) \rightarrow \dots \rightarrow H(X_n), \quad (3.3)$$

where  $H(-)$  generally represents either the  $k$ -th dimensional homology, denoted by  $H_k(-; \mathbb{F})$ , or the total homology, expressed as  $H_\bullet(-; \mathbb{F})$ . Here, this diagram characterizes a sequence of abelian groups or finite-dimensional vector spaces, interconnected through vector space homomorphisms, forming what is known as a persistence module.

Persistence modules are central to understanding how the features of a space evolve over time. They can typically be decomposed into a direct sum of interval modules [8, §8]. Each interval module is associated with an ordered pair of integers  $(p, q)$  where  $1 \leq p \leq q \leq n$ , within a finite filtration. These pairs  $(p, q)$  signify topological features that persist over an index set  $I := \{p, \dots, q\}$ , where  $\inf\{I\} = p$  and  $\sup\{I\} = q$ . Conventionally, these tuples are interpreted as half-open intervals  $[a_p, a_{q+1})$ , with  $a_{n+1} = \infty$  being a customary notation when the sequence extends beyond the largest indexed space.

The decomposition of a persistence module into its constituent interval modules is represented in a persistence diagram or a barcode. This barcode is a multiset of ordered tuples  $(p, q)$  or, alternatively, a multiset of half-open intervals  $[a_p, a_{q+1})$ . This collection is formally expressed through the forgetful functor  $\mathbf{Pers}(-)$ :

$$\mathbf{Pers}(H(\mathcal{X})) = \{(p_1, q_1), \dots, (p_m, q_m)\} \quad (3.4)$$

$$\cong \{[a_{p_1}, a_{q_1+1}), \dots, [a_{p_m}, a_{q_m+1})\}. \quad (3.5)$$

In practical applications, intervals where  $a_p = a_{q+1}$  are usually omitted, as they represent ephemeral topological features.

**Example 3.1.2** *In the further elaboration of the example [6, §2.2], we consider the topological subspaces  $S_1, S_3, S_5$ , all of which are contractible. Meanwhile,  $S_2, S_4, S_6$  are homeomorphic to the 0-sphere, 1-sphere, and 2-sphere, respectively. This structural distinction leads to four distinct intervals in the persistence diagram of the total homology of a sphere, specifically  $\mathcal{S}^2$ :*

$$\mathbf{Pers}(H_\bullet(\mathcal{S}^2)) = \{(1, 6)_0, (2, 2)_0, (4, 4)_1, (6, 6)_2\} \quad (3.6)$$

$$= \{(1, \infty)_0, (2, 3)_0, (4, 5)_1, (6, \infty)_2\}. \quad (3.7)$$

Here, the subscript  $k$  in  $(p, q)_k$  or  $[a_p, a_{q+1})_k$  denotes a topological feature in the  $k$ -dimensional homology.

$$\begin{aligned}
H_\bullet(\mathcal{X}) : \quad & H_\bullet(X_1) \rightarrow \cdots \rightarrow H_\bullet(X_{n-1}) \rightarrow H_\bullet(X_n), \\
H^\bullet(\mathcal{X}) : \quad & H^\bullet(X_1) \leftarrow \cdots \leftarrow H^\bullet(X_{n-1}) \leftarrow H^\bullet(X_n), \\
H_\bullet(X_\infty, \mathcal{X}) : \quad & H_\bullet(X_n) \rightarrow H_\bullet(X_n, X_1) \rightarrow \cdots \rightarrow H_\bullet(X_n, X_{n-1}), \\
H^\bullet(X_\infty, \mathcal{X}) : \quad & H^\bullet(X_n) \leftarrow H^\bullet(X_n, X_1) \leftarrow \cdots \leftarrow H^\bullet(X_n, X_{n-1}).
\end{aligned}$$

Figure 3.1: The four standard persistence modules.

### 3.1.4 The Four Standard Persistence Modules

The standard module of persistent homology,  $H_\bullet(\mathcal{X})$ , illustrates how the absolute homology groups  $H_\bullet(X_i)$  relate to each other as the index  $i$  changes. Similar observations can be made by considering the absolute cohomology groups  $H^\bullet(X_i)$ , the relative homology groups  $H_\bullet(X_n, X_i)$ , and the relative cohomology groups  $H^\bullet(X_n, X_i)$  [6, §2.4].

The persistence diagram for absolute cohomology is represented as a multiset of integer pairs  $(p, q)$ , where  $1 \leq p \leq q \leq n$  for a finite filtration. For relative homology and cohomology, the persistence diagram consists of a multiset of tuples  $(p, q)$  where  $0 \leq p \leq q \leq n - 1$  for a finite filtration. In each case, we interpret  $(p, q)$  as a half-open interval  $[a_p, a_{q+1})$  with the convention that  $a_0 = -\infty$  and  $a_{n+1} = \infty$  [6, §2.4].

**Example 3.1.3** For  $S^2$  we yield

$$\text{Pers}(H_\bullet(S_6, S^2)) = \{(0, 0)_0, (2, 2)_1, (4, 4)_2, (0, 5)_2\} \quad (3.8)$$

$$= \{[-\infty, 1)_0, [2, 3)_1, [4, 5)_2, [-\infty, 6)_3\}. \quad (3.9)$$

At index 2 there is a nontrivial element of  $H_1(S_6, S_2)$  represented by any arc connecting the two points of  $S_2$  [6, §2.4] – the homology class is  $[\sigma_3] = [\sigma_4]$ . This class vanishes in  $H_1(S_6, S_3)$ , thus we yield the interval  $[2, 3)$ .

## 3.2 Persistent (Co)homology

Inverse problems primarily involve inferring geometric shapes from measurements like path integrals. Classical methods such as Fourier transforms provide extensive information but struggle with nonlinearity and ill-posed conditions, requiring substantial regularization. Topology, particularly

through persistent homology, offers alternative methods for deducing topological rather than geometric information. This approach is especially useful in high-dimensional, discrete sets of points, exemplified in the finite case by geological sonar to detect subterranean features based on density variations [6, §1]. Persistent homology identifies topological features represented as intervals in a barcode or persistence diagram, crucial for understanding the presence and persistence of features such as holes or voids in topological spaces. This method is statistically robust and can provide both qualitative and quantitative insights into point sets, which we suspect to lie on some compact topological object [4, 5].

In this section, we continue to examine the consequences of absolute and relative homology and cohomology groups for filtrations of cell complexes, such as the introduced simplicial complexes. In particular, we derive the theory in the context of filtered simplicial complexes upon sets of points, embedded into some metric space. While we can apply the entire theory to simplicial complexes and their (co)homology, cell complexes provide a much broader context and significantly simplify notation in many instances.

In particular, there are at least four naturally arising persistent objects that can be extracted from a filtration of any topological space. We follow the results of de Silva, Morozov, and Vejdemo-Johansson for our explanations [6, §1]. They are

$$\text{persistent} \begin{Bmatrix} \text{absolute} \\ \text{relative} \end{Bmatrix} \begin{Bmatrix} \text{homology} \\ \text{cohomology} \end{Bmatrix}.$$

We address the computation of barcodes for all four types of persistent objects. We demonstrate that both absolute and relative (co)homologies yield identical barcodes and that transitions between these states are facilitated by established duality principles. The duality between homology and cohomology is akin to the duality in vector spaces, whereas a global duality specific to persistent topology allows for a unique interchange:

$$\begin{aligned} \text{Absolute homology} &\rightleftharpoons \text{relative cohomology.} \\ \text{Absolute cohomology} &\rightleftharpoons \text{relative homology.} \end{aligned}$$

The main results from the literature suggest that a single calculation is sufficient to compute all four persistent objects due to the commutative nature of the global duality.

### 3.2.1 Barcode Isomorphisms

We characterise the multisets for persistence modules that are decomposable into interval modules. The persistence diagram partitions into  $\mathbf{Pers}_0$ , comprising finite intervals  $[a, b)$  as per [6, §2.3], and  $\mathbf{Pers}_\infty$ , consisting of intervals  $[a, \infty)$ . This leads to the decomposition  $\mathbf{Pers} = \mathbf{Pers}_0 \cup \mathbf{Pers}_\infty$ .

In this chapter, we establish that persistent homology and cohomology yield the same intervals, or barcodes, for both absolute and relative (co-)homology frameworks. This equivalence necessitates invoking the universal coefficient theorem from algebraic topology, which we will prove beforehand. The universal coefficient theorem for cohomology elegantly ties together the cohomology of a space with coefficients in any abelian group  $G$  to the homology of the space with integer coefficients.

Specifically, as notation for this proof,  $H_n(X; \mathbb{Z})$  and  $H^n(X; \mathbb{Z})$  denote the  $n$ -th singular homology and cohomology groups with coefficients in  $\mathbb{Z}$ , respectively. We further involve  $\text{Hom}(A, G)$ , denoting group homomorphisms from an abelian group  $A$  to another abelian group  $G$ , and  $\text{Ext}^1(A, G)$  [9, §4.3], which measures obstructions in the splitting of a short exact sequence of abelian groups. Further, we use the properties of derived functors [10, §2.7].

**Theorem 3.2.1 (Universal Coefficients for Cohomology)** [9] Let  $X$  be a topological space and  $G$  an abelian group. For any integer  $n \geq 0$ , there is a short exact sequence:

$$0 \rightarrow \text{Ext}^1(H_{n-1}(X; \mathbb{Z}), G) \rightarrow H^n(X; G) \rightarrow \cdots \quad (3.10)$$

$$\cdots \rightarrow \text{Hom}(H_n(X; \mathbb{Z}), G) \rightarrow 0, \quad (3.11)$$

which splits, though not canonically.

**Proof** Let  $C_\bullet(X)$  be the singular chain complex of a topological space  $X$  with integer coefficients. The homology groups  $H_n(X; \mathbb{Z})$  are defined as:

$$H_n(X; \mathbb{Z}) := \ker(\partial_n) / \text{im}(\partial_{n+1}), \quad (3.12)$$

where  $\partial_n$  are the boundary maps in  $C_\bullet(X)$ . The chain group  $C_n(X)$  consists of formal sums of singular  $n$ -simplices in  $X$  with integer coefficients. The

boundary maps  $\partial_n : C_n(X) \rightarrow C_{n-1}(X)$  are defined by

$$\partial_n(\sigma) = \sum_{i=0}^n (-1)^i \sigma|_{[v_0, \dots, \hat{v}_i, \dots, v_n]}, \quad (3.13)$$

where  $\sigma : \Delta^n \rightarrow X$  is a singular simplex, and  $\sigma|_{[v_0, \dots, \hat{v}_i, \dots, v_n]}$  denotes the restriction of  $\sigma$  to the  $i$ -th face of the simplex, omitting the  $i$ -th vertex.

The functor  $\text{Hom}(-, G)$  applied to  $C_n(X)$  yields a group  $\text{Hom}(C_n(X), G)$ , and the coboundary  $\delta^n$  for the cochain complex  $\text{Hom}(C_\bullet(X), G)$  is

$$\delta^n(f) = f \circ \partial_{n+1} \quad (3.14)$$

for  $f \in \text{Hom}(C_{n+1}(X), G)$ . This leads to the cohomology groups

$$H^n(X; G) := \ker(\delta^n) / \text{im}(\delta^{n-1}). \quad (3.15)$$

We consider the projective resolution of  $C_\bullet(X)$  to effectively apply the  $\text{Ext}$  functor. Recall that for any abelian group  $A$ ,

$$\text{Ext}^1(A, G) = R^1 \text{Hom}(A, G), \quad (3.16)$$

where  $R^1$  denotes the first right derived functor of  $\text{Hom}$  [10, §2.7]. To show this equality, we begin by taking a projective resolution of  $A$  [10, p.39f]:

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow A \rightarrow 0, \quad (3.17)$$

where each  $P_i$  is a projective abelian group. Then we can apply the functor  $\text{Hom}(-, G)$  to the projective resolution:

$$0 \rightarrow \text{Hom}(P_0, G) \rightarrow \text{Hom}(P_1, G) \rightarrow \text{Hom}(P_2, G) \rightarrow \cdots \quad (3.18)$$

This sequence is exact on the left because  $\text{Hom}(-, G)$  is left exact and each  $P_i$  is projective. The first right derived functor  $R^1 \text{Hom}(A, G)$  is defined as the cohomology of this sequence at the location corresponding to  $P_1$ :

$$R^1 \text{Hom}(A, G) = \frac{\ker(\text{Hom}(P_1, G) \rightarrow \text{Hom}(P_2, G))}{\text{im}(\text{Hom}(P_0, G) \rightarrow \text{Hom}(P_1, G))}. \quad (3.19)$$

The group  $\text{Ext}^1(A, G)$  classifies extensions of  $A$  by  $G$ , equivalent to the



kernel / image calculation in the cohomology of the Hom-sequence:

$$\text{Ext}^1(A, G) = R^1 \text{Hom}(A, G). \quad (3.20)$$

Given  $H_n(X; \mathbb{Z})$ , consider the short exact sequence obtained from the projective resolution of  $\mathbb{Z}$ :

$$0 \rightarrow \mathbb{Z} \rightarrow F \rightarrow H_n(X; \mathbb{Z}) \rightarrow 0, \quad (3.21)$$

where  $F$  is free. Applying  $\text{Hom}(-, G)$  gives

$$0 \rightarrow \text{Hom}(H_n(X; \mathbb{Z}), G) \rightarrow \text{Hom}(F, G) \rightarrow \cdots \quad (3.22)$$

$$\cdots \rightarrow \text{Hom}(\mathbb{Z}, G) \rightarrow \text{Ext}^1(H_n(X; \mathbb{Z}), G) \rightarrow 0. \quad (3.23)$$

We apply  $\text{Ext}^\bullet$ , which gives rise to the long exact sequence of Ext groups:

$$0 \rightarrow \text{Hom}(H_n(X; \mathbb{Z}), G) \rightarrow H^n(X; G) \rightarrow \cdots \quad (3.24)$$

$$\cdots \rightarrow \text{Ext}^1(H_{n-1}(X; \mathbb{Z}), G) \rightarrow 0. \quad (3.25)$$

Finally, we verify exactness and splitting. The term  $\text{Ext}^1(H_{n-1}(X; \mathbb{Z}), G)$  measures the non-trivial extensions of  $G$  by  $H_{n-1}(X; \mathbb{Z})$ , which corresponds to the obstructions to lifting  $H_{n-1}(X; \mathbb{Z})$  linearly over  $G$ . The term  $\text{Hom}(H_n(X; \mathbb{Z}), G)$  represents the group homomorphisms from  $H_n(X; \mathbb{Z})$  to  $G$ , which naturally includes in  $H^n(X; G)$ . The sequence is exact at each stage by the properties of derived functors and their application to the singular chain complex. The sequence ends with 0 because  $\text{Ext}^1$  of a projective (or free) module vanishes, and  $\mathbb{Z}$  is free. The sequence splits because the functor  $\text{Hom}(-, G)$  preserves products and coproducts. However, the way it splits is not canonical and depends on the choice of a splitting homomorphism, which is not unique.

In the generalization of the universal coefficient theorem to the case of modules over a principal ideal domain, the  $\text{Ext}^1$  terms vanish since  $\mathbb{F}$  is a field, thus we obtain  $H^n(X; \mathbb{F}) \cong \text{Hom}(H_n(X; \mathbb{F}), \mathbb{F})$  [9, p.198 §3.3.1].  $\square$

**Theorem 3.2.2** *For all integers  $d \geq 0$ , it holds that [6, §2.3]:*

$$\text{Pers}(H_d(\mathcal{X})) = \text{Pers}(H^d(\mathcal{X})), \quad (3.26)$$

$$\text{Pers}(H_d(X_\infty, \mathcal{X})) = \text{Pers}(H^d(X_\infty, \mathcal{X})). \quad (3.27)$$

**Proof** When considering coefficients in a field  $\mathbb{F}$  rather than in a ring, the universal coefficient theorem assures us of a natural isomorphism between the  $d$ -th cohomology group and the homomorphisms from the  $d$ -th homology group to the base field:

$$H^d(X; \mathbb{F}) \cong \text{Hom}(H_d(X; \mathbb{F}), \mathbb{F}). \quad (3.28)$$

Therefore, the associated maps

$$H_d(X_i; \mathbb{F}) \rightarrow H_d(X_j; \mathbb{F}) \quad \text{and} \quad H^d(X_i; \mathbb{F}) \leftarrow H^d(X_j; \mathbb{F}) \quad (3.29)$$

are adjoint and hence possess the same rank. Since the persistence intervals over a field are uniquely determined by the dimension and rank of the homology vector spaces, it follows that this holds for both homology and cohomology. Consequently, they share the same barcode.  $\square$

Consider a filtration  $X_1 \subseteq X_2 \subseteq \dots \subseteq X_n$  of a topological space  $X$ , extending to  $X_\infty$  where  $X_\infty$  is the direct limit of the filtration. The homology groups  $H_d(X_n)$  for some fixed dimension  $d$  serve as the initial terms for the relative homology groups  $H_d(X_\infty, X)$ . Since  $H_d(X)$  is consistent with  $H_d(X_n)$  as  $n$  approaches infinity, these sequences can be unified into a single sequence:

$$H_d(X) \rightarrow H_d(X_\infty, X). \quad (3.30)$$

For this concatenated sequence, the indices are denoted as

$$\{1, 2, \dots, n = 0^b, 1^b, 2^b, \dots, (n-1)^b\}, \quad (3.31)$$

using the  $b$  symbol to indicate relative homology part of the sequence.

This structure allows us to discuss the persistence diagram associated with this homological configuration. The persistence intervals in this diagram can generally be categorized into three types:

- $(p, q)$  where  $1 \leq p \leq q < n$ , denoted as  $[p, q+1)$  or  $[a_p, a_{q+1})$ .
- $(p^b, q^b)$  where  $0 < p \leq q \leq n-1$ , denoted as  $[p^b, q^b+1)$  or  $[a_{p^b}, a_{q^b+1})$ .
- $(p, q^b)$  where  $1 \leq p \leq n$  and  $0 \leq q \leq n-1$ , represented as  $[p, q^b+1)$  or  $[a_p, a_{q^b+1})$ .

**Corollary 3.2.3** *The barcode  $\text{Pers}(H_d(\mathcal{X}) \rightarrow H_d(X_\infty, \mathcal{X}))$  comprises the following collections of intervals [6, §2.5]:*

- An interval  $[a, b)$  for every interval  $[a, b)$  in  $\text{Pers}_0(H_d(\mathcal{X}))$ .

- An interval  $[a^b, b^b)$  for every interval  $[a, b)$  in  $\mathbf{Pers}_0(H_{d-1}(\mathcal{X}))$ .
- An interval  $[a, a^b)$  for every interval  $[a, \infty)$  in  $\mathbf{Pers}_\infty(H_d(\mathcal{X}))$ .

**Proof** We begin by analyzing the first two types of intervals in the persistence diagram  $\mathbf{Pers}(H_d(X) \rightarrow H_d(X_\infty, X))$ .

These intervals either do not intersect the intermediate term  $H_d(X_n)$  or terminate before it. Consequently, they correspond precisely to the finite intervals in  $\mathbf{Pers}(H_d(X))$  and  $\mathbf{Pers}(H_d(X_\infty, X))$ , clarifying the first two cases. The correspondence  $\mathbf{Pers}_0(H_d(X_\infty, X)) = \mathbf{Pers}_0(H_{d-1}(X))$  helps in mapping these relationships.

The third case requires examining intervals of the form  $[a, b^b)$  and proving that they are invariably of the form  $[a, a^b)$ , which means that the paired intervals  $[a, \infty)$  and  $[-\infty, a)$  in  $\mathbf{Pers}_\infty(H_d(\mathcal{X}))$  and  $\mathbf{Pers}_\infty(H_d(X_\infty, \mathcal{X}))$  are restrictions of a single interval  $[a, a^b)$  in the concatenated sequence [6, p.6]. To establish this, we compare the ascending filtration defined by the images of  $H_d(X_i)$  in  $H_d(X_n)$  for  $i = 1, 2, \dots, n-1$  (denoted as  $\text{Im}(H_d(X_i) \rightarrow H_d(X_n))$ ) with the descending filtration defined by the kernels of  $H_d(X_n)$  in  $H_d(X_\infty, X)$  for  $i = 1, 2, \dots, n-1$  (denoted as  $\text{Ker}(H_d(X_n) \rightarrow H_d(X_n, X_i))$ ). This examination revolves around the fundamental properties of the homology groups in a filtration setting.

For each index  $i$ , the image and kernel correspond to the same subspace of  $H_d(X_n)$ . This equivalence is guaranteed by the homology long exact sequence associated with the pair  $(X_n, X_i)$ , which links the relative and absolute homology groups. Specifically, the exact sequence implies that any cycle in  $\text{Im}(H_d(X_i) \rightarrow H_d(X_n))$  that becomes a boundary in  $H_d(X_n, X_i)$  must vanish, thus equating the image and kernel. As a result, both filtrations align perfectly, establishing that the third type of interval indeed maps to self-closing intervals of the form  $[a, a^b)$ .  $\square$

### 3.2.2 Persistent Chain Complexes

Alternatively, following [6, §2.6], the standard persistence module can be described via a filtered cell complex  $\mathcal{X} := \sigma_1 \cup \dots \cup \sigma_d$ , so that the persistence module can be expressed as the sequence

$$\mathcal{C} : C_1 \rightarrow C_2 \rightarrow \dots \rightarrow C_d,$$

for a finite filtration, where each  $C_i := \langle \sigma_1, \dots, \sigma_i \rangle$  is a vector space (or an abelian group) over a field  $\mathbb{F}$ , generated by the elements  $\sigma_1, \dots, \sigma_i$ . The boundary operator is defined by  $\partial\sigma_j = \sum_{i < j} \lambda_{ij} \sigma_i$  for  $\lambda_{ij} \in \mathbb{F}$  for all  $i, j \in \{1, \dots, d\}$ .

Then  $C_\bullet(X_i) = (C_i, \partial_i)$  is the chain complex for the absolute homology of  $X_i$ , and  $C_\bullet(\mathcal{X}) = (\mathcal{C}, \partial)$  represents the persistent version for  $\mathcal{X}$ . This leads to the definition of the persistent absolute homology of  $\mathcal{X}$  as

$$H_\bullet(\mathcal{X}) = H(\mathcal{C}, \partial) : \quad \frac{\ker(\partial_1)}{\text{im}(\partial_1)} \rightarrow \frac{\ker(\partial_2)}{\text{im}(\partial_2)} \rightarrow \dots \rightarrow \frac{\ker(\partial_d)}{\text{im}(\partial_d)}.$$

For the persistent absolute cohomology  $H^\bullet(\mathcal{X})$ , we define

$$\mathcal{C}^b : \quad C_1^b \leftarrow C_2^b \leftarrow \dots \leftarrow C_d^b,$$

where  $C_i^b = \text{Hom}(C_i, \mathbb{F}) = \langle \sigma_1^b, \sigma_2^b, \dots, \sigma_i^b \rangle$ , with  $\{\sigma_i^b\}$  as the dual basis of  $\{\sigma_i\}$ . The coboundary  $\delta = \partial^b$  is defined as the adjoint of  $\partial$ . Then  $\mathcal{C}^b(\mathcal{X}) = (\mathcal{C}^b, \delta)$  and  $H^\bullet(\mathcal{X}) = H(\mathcal{C}^b, \delta) = \ker(\delta)/\text{im}(\delta)$ . This too constitutes a persistence module, with morphisms in the reverse direction.

**Example 3.2.1** For  $S^2$ , the boundary operator is given by

$$\partial\sigma_1 = \partial\sigma_2 = 0, \tag{3.32}$$

$$\partial\sigma_3 = \partial\sigma_4 = \sigma_1 - \sigma_2, \tag{3.33}$$

$$\partial\sigma_5 = \partial\sigma_6 = \sigma_3 - \sigma_4. \tag{3.34}$$

This information can also be summarized in a matrix. Similarly, the coboundary operator is given by

$$\delta\sigma_1^b = -\delta\sigma_2^b = \sigma_3^b + \sigma_4^b, \tag{3.35}$$

$$\delta\sigma_3^b = -\delta\sigma_4^b = \sigma_5^b + \sigma_6^b, \tag{3.36}$$

$$\delta\sigma_5^b = \delta\sigma_6^b = 0. \tag{3.37}$$

This data can analogously be summarized in a transposed matrix [6, p.7].

The relative homology and cohomology persistence modules are then defined as the homology of the persistence modules [6, p.7]:

$$(C_d/\mathcal{C}) : \quad C_d \rightarrow (C_d/C_1) \rightarrow (C_d/C_2) \rightarrow \dots \rightarrow (C_d/C_{d-1}), \tag{3.38}$$

$$(C_d/\mathcal{C})^b : \quad C_d^b \leftarrow (C_d/C_1)^b \leftarrow (C_d/C_2)^b \leftarrow \dots \leftarrow (C_d/C_{d-1})^b. \tag{3.39}$$

$$\begin{array}{ccccccc}
& \vdots & & \vdots & & \vdots & & \vdots \\
& \downarrow \partial_{n+2} & & \downarrow \partial_{n+2} & & \downarrow \partial_{n+2} & & \downarrow \partial_{n+2} \\
\cdots & \xrightarrow{f_{n+1}^{i-2}} & C_{n+1}^{i-1} & \xrightarrow{f_{n+1}^{i-1}} & C_{n+1}^i & \xrightarrow{f_{n+1}^i} & C_{n+1}^{i+1} & \xrightarrow{f_{n+1}^{i+1}} & C_{n+1}^{i+2} & \xrightarrow{f_{n+1}^{i+2}} & \cdots \\
& \downarrow \partial_{n+1} & & \downarrow \partial_{n+1} & & \downarrow \partial_{n+1} & & \downarrow \partial_{n+1} \\
\cdots & \xrightarrow{f_n^{i-2}} & C_n^{i-1} & \xrightarrow{f_n^{i-1}} & C_n^i & \xrightarrow{f_n^i} & C_n^{i+1} & \xrightarrow{f_n^{i+1}} & C_n^{i+2} & \xrightarrow{f_n^{i+2}} & \cdots \\
& \downarrow \partial_n & & \downarrow \partial_n & & \downarrow \partial_n & & \downarrow \partial_n \\
\cdots & \xrightarrow{f_{n-1}^{i-2}} & C_{n-1}^{i-1} & \xrightarrow{f_{n-1}^{i-1}} & C_{n-1}^i & \xrightarrow{f_{n-1}^i} & C_{n-1}^{i+1} & \xrightarrow{f_{n-1}^{i+1}} & C_{n-1}^{i+2} & \xrightarrow{f_{n-1}^{i+2}} & \cdots \\
& \downarrow \partial_{n-1} & & \downarrow \partial_{n-1} & & \downarrow \partial_{n-1} & & \downarrow \partial_{n-1} \\
& \vdots & & \vdots & & \vdots & & \vdots
\end{array}$$

Figure 3.2: A persistence complex, where moving to the right increases the filtration index, while moving downwards decreases the dimension.

We note that the mappings  $\rightarrow$  from  $\mathcal{C}$  and the mappings  $\leftarrow$  from  $(C_d/\mathcal{C})^\flat$  are injective, while the mappings  $\leftarrow$  from  $\mathcal{C}^\flat$  and the mappings  $\rightarrow$  from  $(C_d/\mathcal{C})$  are surjective. Thus, absolute and relative cohomology are structurally similar but qualitatively different from absolute homology and relative homology.

**Theorem 3.2.4** (Persistence Partition) [6, §2.6] Let  $\mathcal{C}$  be given and  $\partial$  as above, then there exists a partition

$$\{1, 2, \dots, d\} = P^\times \sqcup P \sqcup N$$

with a bijective pairing  $P \leftrightarrow N$ , such that

$$p \text{ is paired with } n \iff [p, n) \in \mathbf{Pairs}(\mathcal{C}, \partial). \quad (3.40)$$

Furthermore, there is a basis  $\hat{\sigma}_1, \hat{\sigma}_2, \dots, \hat{\sigma}_d$  of  $C_d$  such that

1.  $C_i = \langle \hat{\sigma}_1, \dots, \hat{\sigma}_i \rangle$  for each  $i$ .
2.  $\partial \hat{\sigma}_p = 0$  for all  $p \in P^\times$ .
3.  $\partial \hat{\sigma}_n = \hat{\sigma}_p$ , and thus  $\partial \hat{\sigma}_p = 0$ , for all  $[p, n) \in \mathbf{Pairs}$ .

It follows that the persistence diagram  $\mathbf{Pers}(H(\mathcal{C}, \partial))$  consists of  $[a_p, \infty)$  for  $p \in P^\times$  together with intervals  $[a_p, a_n)$  for  $[p, n) \in \mathbf{Pairs}$ .

**Proof** Let  $\mathcal{C} = \{C_i\}_{i \in I}$  be a filtered chain complex associated with a topological space, structured over a field  $\mathbb{F}$  for any finite index set  $I$ . The filtration indices  $1, 2, \dots, d$ , where  $d = |I|$ , classify the stages at which elements are born or die in homology. The partition of  $\{1, 2, \dots, d\}$  into  $P^\infty \sqcup P \sqcup N$  is then constructed such that:

- $P^\infty$  contains indices corresponding to elements that persist indefinitely, i.e., those for which  $\partial\sigma_i = 0$  and  $\sigma_i$  does not become a boundary at any higher index.
- $P$  and  $N$  are paired bijectively, where each  $p \in P$  (births) corresponds uniquely to an  $n \in N$  (deaths), signifying the termination of the homological feature introduced at  $p$ . This pairing is determined through the matrix reduction of  $\partial$ , ensuring that every cycle born at stage  $p$  becomes a boundary at stage  $n$ .

We then construct a basis  $\hat{\sigma}_i$  such that for each  $i$ ,  $\hat{\sigma}_i$  is a generator of  $C_i$ . Specifically, if  $i \in P^\infty$ , then  $\partial\hat{\sigma}_i = 0$ , indicating these elements are cycles that persist indefinitely. For pairs  $[p, n] \in \mathbf{Pairs}(\mathcal{C}, \partial)$  with  $p \in P$  and  $n \in N$ , set  $\partial\hat{\sigma}_n = \hat{\sigma}_p$  and  $\partial\hat{\sigma}_p = 0$ , reflecting the fact that the cycle  $\hat{\sigma}_p$  born at  $p$  becomes a boundary at  $n$  and thus ceases to contribute to homology past  $n$ .

Hence, the persistence diagram  $\mathbf{Pers}(H(\mathcal{C}, \partial))$  contains intervals  $[a_p, \infty)$  for each  $p \in P^\infty$ , indicating persistent homological features and intervals  $[a_p, a_n)$  for each  $[p, n] \in \mathbf{Pairs}$ , describing features with finite lifespans, starting as a cycle at  $p$  and terminating as a boundary at  $n$ .  $\square$

In 3.2.4 the index sets are defined as follows [6, p.8]:

- $P$  identifies the positive simplices that remain unpaired,
- $P^\infty$  identifies the positive simplices that do become paired,
- $N$  identifies the negative simplices.

The vectors  $\hat{\sigma}_p$  and  $\hat{\sigma}_n$  are cycles characterized by their leading terms  $\sigma_p$  and  $\sigma_n$  respectively, while the vector  $\hat{\sigma}_n$  is a chain with leading term  $\sigma_n$ . This chain 'kills' the homology class of its paired cycle  $\hat{\sigma}_p$  through the boundary relation  $\partial\hat{\sigma}_n = \hat{\sigma}_p$ .

**Theorem 3.2.5** *For all integers  $d \geq 0$  it holds that [6, §2.4]:*

$$\begin{aligned} \mathbf{Pers}(H_d(\mathcal{X})) &\cong \mathbf{Pers}(H_{d+1}(\mathcal{X})), \\ \mathbf{Pers}(H_d(X_\infty, \mathcal{X})) &\cong \mathbf{Pers}(H_d(X_\infty, \mathcal{X})). \end{aligned}$$

**Proof** TBD  $\square$

**Remark 3.2.2** *In this case, we get an isomorphism of multisets. This is due to the identification of the intervals  $[a, \infty) \leftrightarrow [-\infty, a)$  for  $\mathbf{Pers}_\infty$ . Thus, persistent homology and relative homology barcodes carry the same information, with a dimension shift for the finite intervals [6, §2.4].*

### 3.2.3 Cohomology of Chain Complexes

## 3.3 Zigzag Persistence

The zigzag persistence framework is initiated by constructing a zigzag diagram of topological or vector spaces. This diagram consists of a sequence of spaces  $S_1, S_2, \dots, S_n$ , where each adjacent pair  $(S_i, S_{i+1})$  is connected by a map  $S_i \rightarrow S_{i+1}$  or  $S_i \leftarrow S_{i+1}$  [3, §1]. In contrast to the classical approach to persistent homology, the linking maps can now point in arbitrary directions. The major limitation of traditional persistence is its reliance on a nested family  $\{S_t\}_{t \in [0, \infty]}$ , where  $S_t \subseteq S_{t'}$  whenever  $t \leq t'$ . This constraint applies to both the theoretical framework and the algorithms. Zigzag persistence overcomes this limitation. When the variable  $t$  is discretized to a finite set, the family of simplicial complexes can be represented as a diagram of spaces:  $S_1 \rightarrow S_2 \rightarrow \dots \rightarrow S_n$ , where the arrows denote inclusion maps. Applying the  $k$ -dimensional homology functor  $H_k(-; \mathbb{F})$  with coefficients in a field  $\mathbb{F}$ , we obtain a diagram of vector spaces:  $V_1 \rightarrow V_2 \rightarrow \dots \rightarrow V_n$ , where  $V_i = H_k(S_i; \mathbb{F})$ , the persistence module. Persistence is effective because persistence modules can be classified up to isomorphism, with each barcode corresponding to an isomorphism type.

Zigzag persistence aims to achieve a similar classification for diagrams where the arrows may point in either direction, thereby generalizing the framework of traditional persistence. In this section, we follow the work of Carlsson and de Silva [3, §2.1].

### 3.3.1 Zigzag Modules

We consider a sequence of vector spaces and linear maps of length  $n$ , which point in both directions:

$$V : \quad V_1 \xleftarrow{p_1} V_2 \xleftarrow{p_2} \dots \xleftarrow{p_{n-1}} V_n. \quad (3.41)$$

Each  $\xleftarrow{p_i}$  represents either a forward map  $\xrightarrow{f_i}$  or a backward map  $\xleftarrow{g_i}$ . We call the sequence  $V$  a zigzag diagram of vector spaces or a zigzag module

over  $\mathbb{F}$ . The sequence of symbols  $f$  and  $g$  is called the type of  $V$ . For example, a diagram of type  $\tau = fgg$  would be  $V_1 \xrightarrow{f_1} V_2 \xleftarrow{g_2} V_3 \xleftarrow{g_3} V_4$ . The length of a type  $\tau$  is the length of the diagram of type  $\tau$ . Thus,  $fgg$  has length 4. If the length and type of a zigzag module are known, it is called a  $\tau$ -module of length  $n$  and is included in the class  $\tau\text{Mod}$  of  $\tau$ -modules.

### 3.3.2 Decompositions of Zigzag Modules

### 3.3.3 Zigzag Modules and Filtrations

### 3.3.4 Decomposition of Filtered Vector Spaces

### 3.3.5 Streamlined Modules



# Bibliography

1. Bendich, P., Cohen–Steiner, D., Edelsbrunner, H., Harer, J. & Morozov, D. *Inferring local homology from sampled stratified spaces* in *48th Annual IEEE Symposium on Foundations of Computer Science* (2007), 536–546.
2. Boissonnat, J.-D., Chazal, F. & Yvinec, M. *Geometric and topological inference* (Cambridge University Press, 2018).
3. Carlsson, G. & De Silva, V. Zigzag persistence. *Foundations of Computational Mathematics* **10**, 367–405 (2010).
4. Chazal, F., Cohen–Steiner, D., Glisse, M., Guibas, L. J. & Oudot, S. Proximity of persistence modules and their diagrams, 237–246 (2009).
5. Chazal, F., De Silva, V. & Oudot, S. Persistence stability for geometric complexes. *Geometriae Dedicata* **173**, 193–214 (2014).
6. De Silva, V., Morozov, D. & Vejdemo-Johansson, M. Dualities in persistent (co)homology. *Inverse Problems* **27** (2011).
7. Edelsbrunner, H. Triangulations and meshes in computational geometry. *Acta numerica* **9**, 133–213 (2000).
8. Edelsbrunner, H. & Harer, J. *Computational Topology: An Introduction* (American Mathematical Society, 2010).
9. Hatcher, A. *Algebraic topology* (Cambridge University Press, 2005).
10. Weibel, C. A. *An Introduction to Homological Algebra* (Cambridge University Press, Cambridge, 1994).
11. Zomorodian, A. & Carlsson, G. Computing persistent homology. *Symposium on Computational Geometry* **20**, 347–356 (2004).

# Index

$d$ -simplex, 3

abstract simplicial complex, 5

affine subspace, 3

affinely independent, 3

convex hull, 4

face, 3

geometric realization, 5

natural realization, 5

proper faces, 4

properly situated, 4

relative homology group, 25

short exact sequence, 24

simplicial complex, 4

vertex scheme, 5