

Universal Coefficient Theorem Moore-Mayer-Vietoris Sequence for Homology of Ample Groupoids

Luciano Melodia M.A., B.Sc., B.A.

Application to Newcastle University
School of Mathematics, Statistics and Physics

February 18, 2026

Topics

What are we going to talk about?

Ample groupoid homology.

Homology via the Moore chain complex of ample groupoids.

Universal coefficient theorem.

A universal coefficient theorem for discrete abelian groups.

Moore–Mayer–Vietoris sequence.

A Mayer–Vietoris type sequence for clopen saturated covers.

Ample Groupoid Homology

Standing Hypotheses

We investigate ample groupoids.

We consider $C_c(\mathcal{G}_n, A)$.

A is a topological abelian group.

\mathcal{G} is an ample groupoid.

\mathcal{G}_n is the space of n -multiplicable arrows.

Standing Hypotheses

We investigate ample groupoids.

We consider $C_c(\mathcal{G}_n, A)$.

A is a topological abelian group.

\mathcal{G} is an ample groupoid.

\mathcal{G}_n is the space of n -multiplicable arrows.

Why is this important?

\mathcal{G} étale: structure maps in the nerve, such as face maps d_i and degeneracies s_j , are local homeomorphisms, so pushforwards $(d_i)_*$ are defined by finite fibre sums on compact support.

Standing Hypotheses

We investigate ample groupoids.

We consider $C_c(\mathcal{G}_n, A)$.

A is a topological abelian group.

\mathcal{G} is an ample groupoid.

\mathcal{G}_n is the space of n -multiplicable arrows.

Why is this important?

\mathcal{G} étale: structure maps in the nerve, such as face maps d_i and degeneracies s_j , are local homeomorphisms, so pushforwards $(d_i)_*$ are defined by finite fibre sums on compact support.

\mathcal{G} ample: compact open bisections form a basis.

$C_c(\mathcal{G}, \mathbb{Z})$ is generated by χ_K for compact open sets K .

The Nerve

On what do we compute homology?

$\mathcal{G}_\bullet := (\mathcal{G}_n, (d_i)_{i=0}^n, (s_j)_{j=0}^n)_{n \geq 0}$ is a simplicial space .

The Nerve

On what do we compute homology?

$\mathcal{G}_\bullet := (\mathcal{G}_n, (d_i)_{i=0}^n, (s_j)_{j=0}^n)_{n \geq 0}$ is a simplicial space .

Face maps. $d_i: \mathcal{G}_n \rightarrow \mathcal{G}_{n-1}$, $n = 1$: $d_0 = s$, $d_1 = r$. For $n \geq 2$:

$$d_i(\mathbf{g}) := \begin{cases} (g_2, \dots, g_n), & i = 0, \\ (g_1, \dots, g_i \cdot g_{i+1}, \dots, g_n), & 1 \leq i \leq n-1, \\ (g_1, \dots, g_{n-1}), & i = n. \end{cases}$$

The Nerve

On what do we compute homology?

$\mathcal{G}_\bullet := (\mathcal{G}_n, (d_i)_{i=0}^n, (s_j)_{j=0}^n)_{n \geq 0}$ is a simplicial space .

Face maps. $d_i: \mathcal{G}_n \rightarrow \mathcal{G}_{n-1}$, $n = 1$: $d_0 = s$, $d_1 = r$. For $n \geq 2$:

$$d_i(\mathbf{g}) := \begin{cases} (g_2, \dots, g_n), & i = 0, \\ (g_1, \dots, g_i \cdot g_{i+1}, \dots, g_n), & 1 \leq i \leq n-1, \\ (g_1, \dots, g_{n-1}), & i = n. \end{cases}$$

Degeneracy maps. $s_j: \mathcal{G}_n \rightarrow \mathcal{G}_{n+1}$, $n \geq 0$:

$$s_j(\mathbf{g}) := \begin{cases} u(x), & n = 0, x \in \mathcal{G}_0, \\ (u(r(g_1)), g_1, \dots, g_n), & n \geq 1, j = 0, \\ (g_1, \dots, g_j, u(r(g_{j+1})), g_{j+1}, \dots, g_n), & n \geq 2, 1 \leq j \leq n-1, \\ (g_1, \dots, g_n, u(s(g_n))), & n \geq 1, j = n. \end{cases}$$

$$\mathcal{G}_n := \begin{cases} \mathcal{G}_0, & n = 0, \\ \{\mathbf{g} \in \mathcal{G}^n \mid s(g_i) = r(g_{i+1}) \text{ for } 1 \leq i < n\}, & n \geq 1. \end{cases}$$

Moore Chains and Boundary

Compactly supported chains on the nerve.

Chains. $C_c(\mathcal{G}_n, A)$ denotes the abelian group of continuous maps $f: \mathcal{G}_n \rightarrow A$ with compact support. If A is discrete, then every $f \in C_c(\mathcal{G}_n, A)$ is locally constant.

Moore Chains and Boundary

Compactly supported chains on the nerve.

Chains. $C_c(\mathcal{G}_n, A)$ denotes the abelian group of continuous maps $f: \mathcal{G}_n \rightarrow A$ with compact support. If A is discrete, then every $f \in C_c(\mathcal{G}_n, A)$ is locally constant.

Boundary. Since \mathcal{G} is étale, each face map $d_i: \mathcal{G}_n \rightarrow \mathcal{G}_{n-1}$ is a local homeomorphism, hence the pushforward is well-defined:

$$(d_i)_* : C_c(\mathcal{G}_n, A) \rightarrow C_c(\mathcal{G}_{n-1}, A), \quad (d_i)_* f(y) := \sum_{x \in d_i^{-1}(y) \cap \text{supp}(f)} f(x),$$
$$\partial_n := \sum_{i=0}^n (-1)^i (d_i)_* : C_c(\mathcal{G}_n, A) \rightarrow C_c(\mathcal{G}_{n-1}, A).$$

Universal Coefficient Theorem

Universal Coefficient Theorem

Is \mathbb{Z} enough to recover homology through A ?

\mathcal{G} ample groupoid.

A discrete abelian group.

Universal Coefficient Theorem

Is \mathbb{Z} enough to recover homology through A ?

\mathcal{G} ample groupoid.

A discrete abelian group.

For all $n \geq 0$ there is a natural short exact sequence in **Ab**:

$$0 \rightarrow H_n(\mathcal{G}) \otimes_{\mathbb{Z}} A \xrightarrow{\iota_n^{\mathcal{G}}} H_n(\mathcal{G}; A) \xrightarrow{\pi_n^{\mathcal{G}}} \mathrm{Tor}_1^{\mathbb{Z}}(H_{n-1}(\mathcal{G}), A) \rightarrow 0.$$

The sequence splits, though not canonically:

$$H_n(\mathcal{G}; A) \cong (H_n(\mathcal{G}) \otimes_{\mathbb{Z}} A) \oplus \mathrm{Tor}_1^{\mathbb{Z}}(H_{n-1}(\mathcal{G}), A).$$

Universal Coefficient Theorem

If A is discrete, then $C_c(\mathcal{G}_n, A)$ is a free A -module.

Let $f \in C_c(\mathcal{G}_n, A)$. Since A is discrete and $\text{supp}(f)$ is compact, $\text{im}(f) = \{a_1, \dots, a_m\}$ is finite. Set $K_i := f^{-1}(\{a_i\})$ for $1 \leq i \leq m$.

Universal Coefficient Theorem

If A is discrete, then $C_c(\mathcal{G}_n, A)$ is a free A -module.

Let $f \in C_c(\mathcal{G}_n, A)$. Since A is discrete and $\text{supp}(f)$ is compact, $\text{im}(f) = \{a_1, \dots, a_m\}$ is finite. Set $K_i := f^{-1}(\{a_i\})$ for $1 \leq i \leq m$.

Each K_i is clopen in \mathcal{G}_n , the sets K_i are pairwise disjoint, and $\text{supp}(f) = \bigsqcup_{i=1}^m K_i$.

$$f = \sum_{i=1}^m a_i \chi_{K_i} \quad \text{in } C_c(\mathcal{G}_n, A).$$

Universal Coefficient Theorem

If A is discrete, then $C_c(\mathcal{G}_n, A)$ is a free A -module.

Let $f \in C_c(\mathcal{G}_n, A)$. Since A is discrete and $\text{supp}(f)$ is compact, $\text{im}(f) = \{a_1, \dots, a_m\}$ is finite. Set $K_i := f^{-1}(\{a_i\})$ for $1 \leq i \leq m$.

Each K_i is clopen in \mathcal{G}_n , the sets K_i are pairwise disjoint, and $\text{supp}(f) = \bigsqcup_{i=1}^m K_i$.

$$f = \sum_{i=1}^m a_i \chi_{K_i} \quad \text{in } C_c(\mathcal{G}_n, A).$$

Therefore the canonical map

$$\bigoplus_{K \in \mathcal{K}(\mathcal{G}_n)} A \longrightarrow C_c(\mathcal{G}_n, A), \quad (a_K)_{K \in \mathcal{K}(\mathcal{G}_n)} \longmapsto \sum_{K \in \mathcal{K}(\mathcal{G}_n)} a_K \chi_K,$$

is surjective. It is injective: if $\sum_{i=1}^m a_i \chi_{K_i} = 0$ with K_i pairwise disjoint compact open, then evaluating at any $g \in K_i$ gives $a_i = 0$. Thus

$$C_c(\mathcal{G}_n, A) \cong \bigoplus_{K \in \mathcal{K}(\mathcal{G}_n)} A, \quad \text{free as an } A\text{-module}.$$

Non-discrete Coefficients: What Fails

The general result.

For X locally compact, totally disconnected, Hausdorff with a basis of compact open sets and an abelian group A , consider the canonical map

$$\Phi_X: C_c(X, \mathbb{Z}) \otimes_{\mathbb{Z}} A \rightarrow C_c(X, A), \quad \chi_U \otimes a \mapsto a \chi_U.$$

Then

$$\text{im}(\Phi_X) \subseteq \{\xi \in C_c(X, A) \mid \xi(X) \text{ is finite}\}.$$

Non-discrete Coefficients: What Fails

The general result.

For X locally compact, totally disconnected, Hausdorff with a basis of compact open sets and an abelian group A , consider the canonical map

$$\Phi_X: C_c(X, \mathbb{Z}) \otimes_{\mathbb{Z}} A \rightarrow C_c(X, A), \quad \chi_U \otimes a \mapsto a \chi_U.$$

Then

$$\text{im}(\Phi_X) \subseteq \{\xi \in C_c(X, A) \mid \xi(X) \text{ is finite}\}.$$

In particular, Φ_X may fail to be surjective for non-discrete A . Moreover,

$$\Phi_X \text{ surjective} \Leftrightarrow \forall \xi \in C_c(X, A) : \xi(X) \text{ finite} \Leftrightarrow \Phi_X \text{ is an isomorphism.}$$

Non-discrete Coefficients: What Fails

The general result.

For X locally compact, totally disconnected, Hausdorff with a basis of compact open sets and an abelian group A , consider the canonical map

$$\Phi_X: C_c(X, \mathbb{Z}) \otimes_{\mathbb{Z}} A \rightarrow C_c(X, A), \quad \chi_U \otimes a \mapsto a \chi_U.$$

Then

$$\text{im}(\Phi_X) \subseteq \{\xi \in C_c(X, A) \mid \xi(X) \text{ is finite}\}.$$

In particular, Φ_X may fail to be surjective for non-discrete A . Moreover,

$$\Phi_X \text{ surjective} \Leftrightarrow \forall \xi \in C_c(X, A) : \xi(X) \text{ finite} \Leftrightarrow \Phi_X \text{ is an isomorphism.}$$

If A is discrete, then Φ_X is an isomorphism. The converse can fail.

Moore-Mayer-Vietoris Sequence

Mayer-Vietoris vs. Moore-Mayer-Vietoris

From saturated covers to homology.

Mayer-Vietoris:

$$X = U_1 \cup U_2,$$

$U_1, U_2 \subseteq X$ open.

Good cover:

$U_1, U_2, U_1 \cap U_2$ contractible.

In general: any nonempty finite intersection is contractible.

Compute $H_\bullet(X)$ from :

$H_\bullet(U_1), H_\bullet(U_2), H_\bullet(U_1 \cap U_2)$.

Moore-Mayer-Vietoris:

Mayer-Vietoris vs. Moore-Mayer-Vietoris

From saturated covers to homology.

Mayer-Vietoris:

$$X = U_1 \cup U_2,$$

$U_1, U_2 \subseteq X$ open.

Good cover:

$U_1, U_2, U_1 \cap U_2$ contractible.

In general: any nonempty finite intersection is contractible.

Compute $H_\bullet(X)$ from :

$H_\bullet(U_1), H_\bullet(U_2), H_\bullet(U_1 \cap U_2)$.

Moore-Mayer-Vietoris:

$$\mathcal{G}_0 = U_1 \cup U_2,$$

$U_1, U_2 \subseteq \mathcal{G}_0$ clopen.

Mayer-Vietoris vs. Moore-Mayer-Vietoris

From saturated covers to homology.

Mayer-Vietoris:

$$X = U_1 \cup U_2,$$

$$U_1, U_2 \subseteq X \text{ open.}$$

Good cover:

$$U_1, U_2, U_1 \cap U_2 \text{ contractible.}$$

In general: any nonempty finite intersection is contractible.

Compute $H_\bullet(X)$ from :

$$H_\bullet(U_1), H_\bullet(U_2), H_\bullet(U_1 \cap U_2).$$

Moore-Mayer-Vietoris:

$$\mathcal{G}_0 = U_1 \cup U_2,$$

$$U_1, U_2 \subseteq \mathcal{G}_0 \text{ clopen.}$$

Saturated cover:

$$\forall x \in U : s(g) = x \Rightarrow r(g) \in U.$$

$$\mathcal{G}|_U := \{g \in \mathcal{G} \mid s(g), r(g) \in U\},$$

$$\mathcal{G}|_U \hookrightarrow \mathcal{G} \text{ open.}$$

Mayer-Vietoris vs. Moore-Mayer-Vietoris

From saturated covers to homology.

Mayer-Vietoris:

$$X = U_1 \cup U_2,$$

$$U_1, U_2 \subseteq X \text{ open.}$$

Good cover:

$$U_1, U_2, U_1 \cap U_2 \text{ contractible.}$$

In general: any nonempty finite intersection is contractible.

Compute $H_\bullet(X)$ from :

$$H_\bullet(U_1), H_\bullet(U_2), H_\bullet(U_1 \cap U_2).$$

Moore-Mayer-Vietoris:

$$\mathcal{G}_0 = U_1 \cup U_2,$$

$$U_1, U_2 \subseteq \mathcal{G}_0 \text{ clopen.}$$

Saturated cover:

$$\forall x \in U : s(g) = x \Rightarrow r(g) \in U.$$

$$\mathcal{G}|_U := \{g \in \mathcal{G} \mid s(g), r(g) \in U\},$$

$$\mathcal{G}|_U \hookrightarrow \mathcal{G} \text{ open.}$$

Compute $H_\bullet(\mathcal{G})$ from:

$$H_\bullet(\mathcal{G}|_{U_1}), H_\bullet(\mathcal{G}|_{U_2}), H_\bullet(\mathcal{G}|_{U_1 \cap U_2}).$$

Reductions and Moore–Mayer–Vietoris

Long Natural Moore-Mayer-Vietoris Sequence for Homology.

For $\mathcal{U} \subseteq \mathcal{G}_0$ define the reduction

$$\mathcal{G}|_{\mathcal{U}} := \{g \in \mathcal{G} \mid s(g) \in \mathcal{U}, r(g) \in \mathcal{U}\}, \quad (\mathcal{G}|_{\mathcal{U}})_0 = \mathcal{U},$$

with structure maps the restrictions of $u, m, s, r, -^{-1}$ to $\mathcal{G}|_{\mathcal{U}}$. For $i \in \{1, 2\}$ and $\mathcal{U}_{12} := \mathcal{U}_1 \cap \mathcal{U}_2$ we write $\mathcal{G}|_{\mathcal{U}_i}$ and $\mathcal{G}|_{\mathcal{U}_{12}}$.

Reductions and Moore–Mayer–Vietoris

Long Natural Moore–Mayer–Vietoris Sequence for Homology.

For $\mathcal{U} \subseteq \mathcal{G}_0$ define the reduction

$$\mathcal{G}|_{\mathcal{U}} := \{g \in \mathcal{G} \mid s(g) \in \mathcal{U}, r(g) \in \mathcal{U}\}, \quad (\mathcal{G}|_{\mathcal{U}})_0 = \mathcal{U},$$

with structure maps the restrictions of $u, m, s, r, -^1$ to $\mathcal{G}|_{\mathcal{U}}$. For $i \in \{1, 2\}$ and $\mathcal{U}_{12} := \mathcal{U}_1 \cap \mathcal{U}_2$ we write $\mathcal{G}|_{\mathcal{U}_i}$ and $\mathcal{G}|_{\mathcal{U}_{12}}$.

Moore–Mayer–Vietoris long exact homology sequence:

$$\begin{array}{ccccccc} \cdots & \longleftarrow & H_{n-1}(\mathcal{G}|_{\mathcal{U}_1}; \mathcal{A}) \oplus H_{n-1}(\mathcal{G}|_{\mathcal{U}_2}; \mathcal{A}) & \xleftarrow{H_{n-1}(\alpha_\bullet)} & H_{n-1}(\mathcal{G}|_{\mathcal{U}_1 \cap \mathcal{U}_2}; \mathcal{A}) & \longleftarrow & \\ & & \partial_n & & & & \\ & \longleftarrow & H_n(\mathcal{G}; \mathcal{A}) & \xleftarrow{H_n(\beta_\bullet)} & H_n(\mathcal{G}|_{\mathcal{U}_1}; \mathcal{A}) \oplus H_n(\mathcal{G}|_{\mathcal{U}_2}; \mathcal{A}) & \xleftarrow{H_n(\alpha_\bullet)} & H_n(\mathcal{G}|_{\mathcal{U}_1 \cap \mathcal{U}_2}; \mathcal{A}) & \longleftarrow & \\ & & \partial_{n+1} & & & & \\ & \longleftarrow & H_{n+1}(\mathcal{G}; \mathcal{A}) & \xleftarrow{H_{n+1}(\beta_\bullet)} & H_{n+1}(\mathcal{G}|_{\mathcal{U}_1}; \mathcal{A}) \oplus H_{n+1}(\mathcal{G}|_{\mathcal{U}_2}; \mathcal{A}) & \longleftarrow & \cdots \end{array}$$

Takeaways

What you get and how to use it.

Setting: \mathcal{G} ample étale, A a topological abelian group,

Moore chains $C_c(\mathcal{G}_n, A)$ with boundary $\partial = \sum_{i=0}^n (-1)^i (d_i)_*$.

Two structural tools:

Takeaways

What you get and how to use it.

Setting: \mathcal{G} ample étale, A a topological abelian group,

Moore chains $C_c(\mathcal{G}_n, A)$ with boundary $\partial = \sum_{i=0}^n (-1)^i (d_i)_*$.

Two structural tools:

UCT for discrete coefficients:

$$0 \rightarrow H_n(\mathcal{G}) \otimes_{\mathbb{Z}} A \xrightarrow{\iota_n^{\mathcal{G}}} H_n(\mathcal{G}; A) \xrightarrow{\pi_n^{\mathcal{G}}} \mathrm{Tor}_1^{\mathbb{Z}}(H_{n-1}(\mathcal{G}), A) \rightarrow 0.$$

Takeaways

What you get and how to use it.

Setting: \mathcal{G} ample étale, A a topological abelian group,
Moore chains $C_c(\mathcal{G}_n, A)$ with boundary $\partial = \sum_{i=0}^n (-1)^i (d_i)_*$.

Two structural tools:

UCT for discrete coefficients:

$$0 \rightarrow H_n(\mathcal{G}) \otimes_{\mathbb{Z}} A \xrightarrow{\iota_n^{\mathcal{G}}} H_n(\mathcal{G}; A) \xrightarrow{\pi_n^{\mathcal{G}}} \mathrm{Tor}_1^{\mathbb{Z}}(H_{n-1}(\mathcal{G}), A) \rightarrow 0.$$

Moore–Mayer–Vietoris

for a clopen saturated cover $U_1 \cup U_2 = \mathcal{G}_0$.

There is a natural long exact sequence relating

$$H_{\bullet}(\mathcal{G}; A), H_{\bullet}(\mathcal{G}|_{U_1}; A), H_{\bullet}(\mathcal{G}|_{U_2}; A), H_{\bullet}(\mathcal{G}|_{U_{12}}; A).$$

Takeaways

What you get and how to use it.

Why discreteness matters:

For non-discrete A , the canonical comparison

$$\Phi_X : C_c(X, \mathbb{Z}) \otimes_{\mathbb{Z}} A \rightarrow C_c(X, A), \quad \chi_u \otimes a \mapsto a \chi_u,$$

need not be surjective. Tensor-level reduction in UCT can fail.

Takeaways

What you get and how to use it.

Why discreteness matters:

For non-discrete A , the canonical comparison

$$\Phi_X : C_c(X, \mathbb{Z}) \otimes_{\mathbb{Z}} A \rightarrow C_c(X, A), \quad \chi_u \otimes a \mapsto a \chi_u,$$

need not be surjective. Tensor-level reduction in UCT can fail.

How to use in practice:

Choose a clopen saturated cover.

$U_1, U_2 \subseteq \mathcal{G}_0$ so that reductions $\mathcal{G}|_{U_1}$, $\mathcal{G}|_{U_2}$, $\mathcal{G}|_{U_{12}}$ are computable.

Compute integral homology.

$H_{\bullet}(\mathcal{G}|_{U_i})$, $H_{\bullet}(\mathcal{G}|_{U_{12}})$, then glue to $H_{\bullet}(\mathcal{G})$ via MMV.

Thank you.

Homology of SFT Groupoids

Example: Diaconu-Renault Groupoid

Computing Homology with Moore–Mayer–Vietoris + UCT.

$A \in \text{Mat}(\mathbb{N} \times \mathbb{N}, \mathbb{N}_0)$ with no zero row and no zero column.

E_A a finite directed graph whose adjacency matrix is A .

The infinite path space is given by Sims 2021, 2.5:

$E_A^\infty = \{(e_n)_{n \geq 1} \in E_A^{\mathbb{N}} \mid r(e_n) = s(e_{n+1}) \text{ for all } n \geq 1\}$ with

$r(x, n, y) = x$, $s(x, n, y) = y$, $1_x = (x, 0, x)$,

$(x, n, y)^{-1} = (y, -n, x)$, $(x, n, y) \cdot (y, m, z) = (x, n + m, z)$ if $s(x, n, y) = r(y, m, z)$.

$\sigma: E_A^\infty \rightarrow E_A^\infty, (e_0, e_1, e_2, \dots) \mapsto (e_1, e_2, e_3, \dots)$.

$(\mathcal{G}_A)_0 = E_A^\infty$.

$(\mathcal{G}_A)_1 = \{(x, n, y) \in E_A^\infty \times \mathbb{Z} \times E_A^\infty \mid \exists k, \ell \in \mathbb{N}_0: n = k - \ell, \sigma^k(x) = \sigma^\ell(y)\}$.

Example: Diaconu-Renault Groupoid

Homology of SFT-Groupoids is well known.

$1 - A^T$ acts on \mathbb{Z}^N and we have by Matui 2012, 4.14:

$$H_0(\mathcal{G}_A) \cong \operatorname{coker}(1 - A^T),$$

$$H_1(\mathcal{G}_A) \cong \operatorname{ker}(1 - A^T),$$

$$H_n(\mathcal{G}_A) = 0 \text{ for } n \geq 2.$$

Consider now:

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}, \quad C = (3).$$

We compute the integral homology of \mathcal{G}_A , \mathcal{G}_B , and \mathcal{G}_C .

Example: Diaconu-Renault Groupoid

Computing Homology for \mathcal{G}_A .

For A we have

$$\mathbf{1} - A^T = \begin{pmatrix} -1 & -1 \\ -1 & 1 \end{pmatrix}, \quad \det(\mathbf{1} - A^T) = -2.$$

Hence $\mathbf{1} - A^T$ has full rank over \mathbb{Z} and $\ker(\mathbf{1} - A^T) = 0$.

Moreover, the Smith normal form is

$$\begin{pmatrix} -1 & -1 \\ -1 & 1 \end{pmatrix} \xrightarrow{R_1 \leftarrow -R_1} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \xrightarrow{R_2 \leftarrow R_2 + R_1} \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix} \xrightarrow{C_2 \leftarrow C_2 - C_1} \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix},$$

so $\operatorname{coker}(\mathbf{1} - A^T) \cong \mathbb{Z}/2\mathbb{Z}$.

We get $H_0(\mathcal{G}_A) \cong \mathbb{Z}/2\mathbb{Z}$, $H_1(\mathcal{G}_A) = 0$, $H_n(\mathcal{G}_A) = 0$ for $n \geq 2$.

Example: Diaconu-Renault Groupoid

Computing Homology for \mathcal{G}_B .

For B we have

$$\mathbf{1} - B^T = \begin{pmatrix} -1 & -1 \\ -1 & -1 \end{pmatrix}.$$

$(\mathbf{1} - B^T)(x, y)^T = 0 \Leftrightarrow -x - y = 0$, hence $\ker(\mathbf{1} - B^T) \cong \mathbb{Z}$.

The image is generated by $(1, 1)$, which is primitive in \mathbb{Z}^2 , so

$$\operatorname{coker}(\mathbf{1} - B^T) \cong \mathbb{Z}^2 / \langle (1, 1) \rangle_{\mathbb{Z}} \cong \mathbb{Z}.$$

Thus, we have for homology $H_0(\mathcal{G}_B) \cong \mathbb{Z}$, $H_1(\mathcal{G}_B) \cong \mathbb{Z}$, $H_n(\mathcal{G}_B) = 0$ for $n \geq 2$.

Example: Diaconu-Renault Groupoid

Computing Homology for \mathcal{G}_C .

For C we have $1 - C^T = -2$, so

$$\ker(1 - C^T) = 0 \quad \text{and} \quad \operatorname{coker}(1 - C^T) \cong \mathbb{Z}/2\mathbb{Z}.$$

Hence $H_0(\mathcal{G}_C) \cong \mathbb{Z}/2\mathbb{Z}$, $H_1(\mathcal{G}_C) = 0$, $H_n(\mathcal{G}_C) = 0$ for $n \geq 2$.

Example: Diaconu-Renault Groupoid

The disjoint union groupoid.

We have $\mathcal{G} = \mathcal{G}_A \sqcup \mathcal{G}_B \sqcup \mathcal{G}_C$, the disjoint union groupoid.

The nerve decomposes levelwise to $\mathcal{G}_n = (\mathcal{G}_A)_n \sqcup (\mathcal{G}_B)_n \sqcup (\mathcal{G}_C)_n$.

The Moore chain complex splits as a direct sum, thus

$$H_n(\mathcal{G}) \cong H_n(\mathcal{G}_A) \oplus H_n(\mathcal{G}_B) \oplus H_n(\mathcal{G}_C) \text{ for } n \geq 0.$$

In particular

$$H_0(\mathcal{G}) \cong \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})^2, \quad H_1(\mathcal{G}) \cong \mathbb{Z}, \quad H_n(\mathcal{G}) = 0 \text{ for } n \geq 2.$$

Define $\mathcal{U}_1 := (\mathcal{G}_A)_0 \sqcup (\mathcal{G}_B)_0$, $\mathcal{U}_2 := (\mathcal{G}_B)_0 \sqcup (\mathcal{G}_C)_0$.

Example: Diaconu-Renault Groupoid

The reduction groupoids.

The reductions are

$$\mathcal{G}|_{U_1} = \mathcal{G}_A \sqcup \mathcal{G}_B, \quad \mathcal{G}|_{U_2} = \mathcal{G}_B \sqcup \mathcal{G}_C, \quad \mathcal{G}|_{U_1 \cap U_2} = \mathcal{G}_B.$$

This yields the long exact sequence

$$\begin{array}{ccccccc} \cdots \rightarrow H_n(\mathcal{G}_B) & \xrightarrow{\alpha_n} & H_n(\mathcal{G}_A \sqcup \mathcal{G}_B) \oplus H_n(\mathcal{G}_B \sqcup \mathcal{G}_C) & \xrightarrow{\beta_n} & & & \\ & & \xrightarrow{\beta_n} & H_n(\mathcal{G}) & \xrightarrow{\partial_n} & H_{n-1}(\mathcal{G}_B) & \rightarrow \cdots \end{array}$$

Example: Diaconu-Renault Groupoid

Explicit formulas for α_n , β_n , ∂_n

$$\alpha_n : H_n(\mathcal{G}_B) \rightarrow H_n(\mathcal{G}_A) \oplus H_n(\mathcal{G}_B) \oplus H_n(\mathcal{G}_B) \oplus H_n(\mathcal{G}_C),$$

$$[b] \mapsto ([0], [b], [-b], [0]),$$

$$\beta_n : H_n(\mathcal{G}_A) \oplus H_n(\mathcal{G}_B) \oplus H_n(\mathcal{G}_B) \oplus H_n(\mathcal{G}_C) \rightarrow H_n(\mathcal{G}_A) \oplus H_n(\mathcal{G}_B) \oplus H_n(\mathcal{G}_C),$$

$$([a], [b_1], [b_2], [c]) \mapsto ([a], [b_1 + b_2], [c]).$$

∂_n vanishes in this example by exactness $\partial_n = 0: H_n(\mathcal{G}) \rightarrow H_{n-1}(\mathcal{G}_B)$, since β_n is surjective and $\ker(\beta_n) = \{([0], [b], [-b], [0]) \mid b \in H_n(\mathcal{G}_B)\} = \text{im}(\alpha_n)$.

Finite coefficients via UCT

Final homology groups for $\mathbb{Z}/p\mathbb{Z}$.

Fix a prime p . Assume $H_2(\mathcal{G}) = 0$, $H_1(\mathcal{G}) \cong \mathbb{Z}$, $H_0(\mathcal{G}) \cong \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})^2$.

Vanishing in higher degrees: $H_n(\mathcal{G}; \mathbb{Z}/p\mathbb{Z}) = 0$ for all $n \geq 2$.

Degree 0:

$$H_0(\mathcal{G}; \mathbb{Z}/p\mathbb{Z}) \cong H_0(\mathcal{G}) \otimes_{\mathbb{Z}} \mathbb{Z}/p\mathbb{Z} \cong \begin{cases} \mathbb{Z}/p\mathbb{Z}, & \text{for } p \text{ odd,} \\ (\mathbb{Z}/2\mathbb{Z})^3, & \text{for } p = 2. \end{cases}$$

Degree 1 via UCT:

$$0 \rightarrow H_1(\mathcal{G}) \otimes_{\mathbb{Z}} \mathbb{Z}/p\mathbb{Z} \rightarrow H_1(\mathcal{G}; \mathbb{Z}/p\mathbb{Z}) \rightarrow \operatorname{Tor}_1^{\mathbb{Z}}(H_0(\mathcal{G}), \mathbb{Z}/p\mathbb{Z}) \rightarrow 0,$$

hence

$$H_1(\mathcal{G}; \mathbb{Z}/p\mathbb{Z}) \cong \begin{cases} \mathbb{Z}/p\mathbb{Z}, & \text{for } p \text{ odd,} \\ (\mathbb{Z}/2\mathbb{Z})^3, & \text{for } p = 2. \end{cases}$$

Proof of the UCT

Proof of the UCT

Step 1: Chain-level identification.

Let $f \in C_c(\mathcal{G}_n, A)$ and write $\text{im}(f) = \{a_1, \dots, a_m\}$. Set $K_i := f^{-1}(\{a_i\})$. Then $\text{supp}(f) = \bigsqcup_{i=1}^m K_i$ with K_i clopen and $f|_{K_i} \equiv a_i$.

Extension by 0: $\chi_{K_i} \in C_c(\mathcal{G}_n, \mathbb{Z})$ and

$$f = \sum_{i=1}^m a_i \chi_{K_i} \quad \text{in } C_c(\mathcal{G}_n, A).$$

Proof of the UCT

Step 1: Chain-level identification.

Let $f \in C_c(\mathcal{G}_n, A)$ and write $\text{im}(f) = \{a_1, \dots, a_m\}$. Set $K_i := f^{-1}(\{a_i\})$. Then $\text{supp}(f) = \bigsqcup_{i=1}^m K_i$ with K_i clopen and $f|_{K_i} \equiv a_i$.

Extension by 0: $\chi_{K_i} \in C_c(\mathcal{G}_n, \mathbb{Z})$ and

$$f = \sum_{i=1}^m a_i \chi_{K_i} \quad \text{in } C_c(\mathcal{G}_n, A).$$

Define the canonical \mathbb{Z} -bilinear map

$$\Phi_{\mathcal{G}_n} : C_c(\mathcal{G}_n, \mathbb{Z}) \otimes_{\mathbb{Z}} A \longrightarrow C_c(\mathcal{G}_n, A), \quad \xi \otimes a \longmapsto a \cdot \xi.$$

Proof of the UCT

Step 1: Chain-level identification.

Let $f \in C_c(\mathcal{G}_n, A)$ and write $\text{im}(f) = \{a_1, \dots, a_m\}$. Set $K_i := f^{-1}(\{a_i\})$. Then $\text{supp}(f) = \bigsqcup_{i=1}^m K_i$ with K_i clopen and $f|_{K_i} \equiv a_i$.

Extension by 0: $\chi_{K_i} \in C_c(\mathcal{G}_n, \mathbb{Z})$ and

$$f = \sum_{i=1}^m a_i \chi_{K_i} \quad \text{in } C_c(\mathcal{G}_n, A).$$

Define the canonical \mathbb{Z} -bilinear map

$$\Phi_{\mathcal{G}_n} : C_c(\mathcal{G}_n, \mathbb{Z}) \otimes_{\mathbb{Z}} A \longrightarrow C_c(\mathcal{G}_n, A), \quad \xi \otimes a \longmapsto a \cdot \xi.$$

$\Phi_{\mathcal{G}_n}$ is surjective, and injective since $C_c(\mathcal{G}_n, \mathbb{Z})$ is free on $\{\chi_K \mid K \in \mathcal{K}(\mathcal{G}_n)\}$.

Proof of the UCT

Step 1: Chain-level identification.

Let $f \in C_c(\mathcal{G}_n, A)$ and write $\text{im}(f) = \{a_1, \dots, a_m\}$. Set $K_i := f^{-1}(\{a_i\})$. Then $\text{supp}(f) = \bigsqcup_{i=1}^m K_i$ with K_i clopen and $f|_{K_i} \equiv a_i$.

Extension by 0: $\chi_{K_i} \in C_c(\mathcal{G}_n, \mathbb{Z})$ and

$$f = \sum_{i=1}^m a_i \chi_{K_i} \quad \text{in } C_c(\mathcal{G}_n, A).$$

Define the canonical \mathbb{Z} -bilinear map

$$\Phi_{\mathcal{G}_n} : C_c(\mathcal{G}_n, \mathbb{Z}) \otimes_{\mathbb{Z}} A \longrightarrow C_c(\mathcal{G}_n, A), \quad \xi \otimes a \longmapsto a \cdot \xi.$$

$\Phi_{\mathcal{G}_n}$ is surjective, and injective since $C_c(\mathcal{G}_n, \mathbb{Z})$ is free on $\{\chi_K \mid K \in \mathcal{K}(\mathcal{G}_n)\}$.

$$C_c(\mathcal{G}_n, \mathbb{Z}) \otimes_{\mathbb{Z}} A \cong C_c(\mathcal{G}_n, A) \quad \text{for discrete } A.$$

Proof of the UCT

Step 2: Compatibility with the boundary.

For each face map $d_i: \mathcal{G}_n \rightarrow \mathcal{G}_{n-1}$, the pushforward $(d_i)_*$ is \mathbb{Z} -linear and satisfies

$$(d_i)_*(\xi \cdot a) = ((d_i)_*\xi) \cdot a \quad \text{for } \xi \in C_c(\mathcal{G}_n, \mathbb{Z}), a \in A.$$

Proof of the UCT

Step 2: Compatibility with the boundary.

For each face map $d_i: \mathcal{G}_n \rightarrow \mathcal{G}_{n-1}$, the pushforward $(d_i)_*$ is \mathbb{Z} -linear and satisfies

$$(d_i)_*(\xi \cdot a) = ((d_i)_*\xi) \cdot a \quad \text{for } \xi \in C_c(\mathcal{G}_n, \mathbb{Z}), a \in A.$$

Hence $\Phi_{\mathcal{G}_\bullet}$ intertwines the Moore boundary:

$$\Phi_{\mathcal{G}_{n-1}} \circ (\partial_n \otimes \text{id}_A) = \partial_n \circ \Phi_{\mathcal{G}_n}.$$

Proof of the UCT

Step 2: Compatibility with the boundary.

For each face map $d_i: \mathcal{G}_n \rightarrow \mathcal{G}_{n-1}$, the pushforward $(d_i)_*$ is \mathbb{Z} -linear and satisfies

$$(d_i)_*(\xi \cdot a) = ((d_i)_*\xi) \cdot a \quad \text{for } \xi \in C_c(\mathcal{G}_n, \mathbb{Z}), a \in A.$$

Hence $\Phi_{\mathcal{G}_\bullet}$ intertwines the Moore boundary:

$$\Phi_{\mathcal{G}_{n-1}} \circ (\partial_n \otimes \text{id}_A) = \partial_n \circ \Phi_{\mathcal{G}_n}.$$

Therefore $\Phi_{\mathcal{G}_\bullet}$ is an isomorphism of chain complexes

$$C_c(\mathcal{G}_\bullet, \mathbb{Z}) \otimes_{\mathbb{Z}} A \cong C_c(\mathcal{G}_\bullet, A).$$

Proof of the UCT

Step 3: Apply the classical algebraic UCT.

The Moore complex $C_c(\mathcal{G}_\bullet, \mathbb{Z})$ is a chain complex of free abelian groups. Applying the classical algebraic UCT to $C_c(\mathcal{G}_\bullet, \mathbb{Z})$ and transporting across the chain isomorphism from Steps 1–2 yields, for all $n \geq 0$, a short exact sequence

$$0 \rightarrow H_n(\mathcal{G}) \otimes_{\mathbb{Z}} A \xrightarrow{\iota_n^{\mathcal{G}}} H_n(\mathcal{G}; A) \xrightarrow{\pi_n^{\mathcal{G}}} \operatorname{Tor}_1^{\mathbb{Z}}(H_{n-1}(\mathcal{G}), A) \rightarrow 0.$$

Proof of the UCT

Step 3: Apply the classical algebraic UCT.

The Moore complex $C_c(\mathcal{G}_\bullet, \mathbb{Z})$ is a chain complex of free abelian groups. Applying the classical algebraic UCT to $C_c(\mathcal{G}_\bullet, \mathbb{Z})$ and transporting across the chain isomorphism from Steps 1–2 yields, for all $n \geq 0$, a short exact sequence

$$0 \rightarrow H_n(\mathcal{G}) \otimes_{\mathbb{Z}} A \xrightarrow{\iota_n^{\mathcal{G}}} H_n(\mathcal{G}; A) \xrightarrow{\pi_n^{\mathcal{G}}} \operatorname{Tor}_1^{\mathbb{Z}}(H_{n-1}(\mathcal{G}), A) \rightarrow 0.$$

This sequence is natural in \mathcal{G} and in discrete A . In general, for non-discrete topological abelian groups A , Moore homology need not satisfy such a UCT.

Failure of Isomorphism

Non-discrete Coefficients: What Fails

Failure of the tensor comparison.

If A is non-discrete and $0 \in A$ is not isolated, then surjectivity of Φ_X can fail even for compact, totally disconnected spaces with a basis of clopen subsets. Set

$$X := \left\{ \sum_{n=1}^{\infty} \frac{a_n}{3^n} \mid a_n \in \{0, 2\} \right\} \subset [0, 1],$$

$$A := (\mathbb{R}, \mathcal{O}_{\text{std}}),$$

$$\xi: X \rightarrow A, \quad x \mapsto x.$$

Then X is compact, Hausdorff, totally disconnected, and has a basis of clopen subsets. Hence $\xi \in C_c(X, A)$ and $\xi(X) = X$ is infinite. Therefore $\xi \notin \text{im}(\Phi_X)$, so Φ_X is not surjective.

Non-discrete Coefficients: What Fails

Isomorphism without discreteness.

$$A := (\mathbb{R}, \mathcal{O}_{\text{std}}), \quad (\{\bullet\}, \mathcal{O}_{\{\bullet\}} := \{\emptyset, \{\bullet\}\}).$$

Then $\{\bullet\}$ is locally compact, totally disconnected, Hausdorff, and X is compact open.

$$C_c(\{\bullet\}, \mathbb{Z}) \cong \mathbb{Z},$$

$$C_c(\{\bullet\}, A) \cong A,$$

$$C_c(\{\bullet\}, \mathbb{Z}) \otimes_{\mathbb{Z}} A \cong \mathbb{Z} \otimes_{\mathbb{Z}} A \cong A.$$

Under these identifications the canonical map

$$\Phi_{\{\bullet\}} : C_c(\{\bullet\}, \mathbb{Z}) \otimes_{\mathbb{Z}} A \rightarrow C_c(\{\bullet\}, A), \quad \chi_{\{\bullet\}} \otimes a \mapsto a \cdot \chi_{\{\bullet\}},$$

is the standard isomorphism $\mathbb{Z} \otimes_{\mathbb{Z}} A \rightarrow A$, $1 \otimes a \mapsto a$.

Proof of Moore–Mayer–Vietoris

Proof of Moore–Mayer–Vietoris

Proof idea for $H_n(\alpha_\bullet)$.

$(\iota_i)_n: (\mathcal{G}|_{U_{12}})_n \hookrightarrow (\mathcal{G}|_{U_i})_n$ is an open embedding, hence a local homeomorphism.
Therefore the functorial pushforward on Moore chains

$$((\iota_i)_n)_*: C_c((\mathcal{G}|_{U_{12}})_n, A) \rightarrow C_c((\mathcal{G}|_{U_i})_n, A)$$

is given by a **finite fibre sum**. Since $(\iota_i)_n$ is injective, it is extension by zero:

$$((\iota_i)_n)_* f(x) := \begin{cases} f(x), & x \in (\mathcal{G}|_{U_{12}})_n, \\ 0, & x \in (\mathcal{G}|_{U_i})_n \setminus (\mathcal{G}|_{U_{12}})_n. \end{cases}$$

Proof of Moore–Mayer–Vietoris

Proof idea for $H_n(\alpha_\bullet)$.

$(\iota_i)_n: (\mathcal{G}|_{U_{12}})_n \hookrightarrow (\mathcal{G}|_{U_i})_n$ is an open embedding, hence a local homeomorphism. Therefore the functorial pushforward on Moore chains

$$((\iota_i)_n)_*: C_c((\mathcal{G}|_{U_{12}})_n, A) \rightarrow C_c((\mathcal{G}|_{U_i})_n, A)$$

is given by a **finite fibre sum**. Since $(\iota_i)_n$ is injective, it is extension by zero:

$$((\iota_i)_n)_* f(x) := \begin{cases} f(x), & x \in (\mathcal{G}|_{U_{12}})_n, \\ 0, & x \in (\mathcal{G}|_{U_i})_n \setminus (\mathcal{G}|_{U_{12}})_n. \end{cases}$$

Define the chain map

$$\begin{aligned} \alpha_n: C_c((\mathcal{G}|_{U_{12}})_n, A) &\rightarrow C_c((\mathcal{G}|_{U_1})_n, A) \oplus C_c((\mathcal{G}|_{U_2})_n, A), \\ f &\mapsto ((\iota_1)_n)_* f, -((\iota_2)_n)_* f. \end{aligned}$$

Compatibility: $\partial((\iota_i)_n)_* = ((\iota_i)_{n-1})_* \partial$, $\partial \alpha_n = \alpha_{n-1} \partial$. Hence α_\bullet induces $H_n(\alpha_\bullet)$.

Proof of Moore–Mayer–Vietoris

Proof idea for $H_n(\beta_\bullet)$.

Let $\kappa_i: \mathcal{G}|_{U_i} \hookrightarrow \mathcal{G}$ be the inclusion of reductions.

$(\kappa_i)_n: (\mathcal{G}|_{U_i})_n \hookrightarrow \mathcal{G}_n$ is an open embedding, hence a local homeomorphism. Therefore the pushforward on Moore chains extends by zero:

$$((\kappa_i)_n)_* g(x) := \begin{cases} g(x), & x \in (\mathcal{G}|_{U_i})_n, \\ 0, & x \in \mathcal{G}_n \setminus (\mathcal{G}|_{U_i})_n. \end{cases}$$

Define

$$\begin{aligned} \beta_n: C_c((\mathcal{G}|_{U_1})_n, A) \oplus C_c((\mathcal{G}|_{U_2})_n, A) &\rightarrow C_c(\mathcal{G}_n, A), \\ (g_1, g_2) &\mapsto ((\kappa_1)_n)_* g_1 + ((\kappa_2)_n)_* g_2. \end{aligned}$$

Compatibility: $\partial((\kappa_i)_n)_* = ((\kappa_i)_{n-1})_* \partial$, $\partial \beta_n = \beta_{n-1} \partial$. Hence β_\bullet induces $H_n(\beta_\bullet)$.

Proof of Moore–Mayer–Vietoris

Proof idea for ∂_n .

Assume a SES of Moore chain complexes

$$\begin{aligned} 0 \rightarrow C_c((\mathcal{G}|_{U_{12}})_\bullet, A) &\xrightarrow{\alpha_\bullet} C_c((\mathcal{G}|_{U_1})_\bullet, A) \oplus C_c((\mathcal{G}|_{U_2})_\bullet, A) \\ &\xrightarrow{\beta_\bullet} C_c(\mathcal{G}_\bullet, A) \rightarrow 0. \end{aligned}$$

Here ∂ denotes the Moore boundary.

Let $[c] \in H_n(\mathcal{G}; A)$ with $\partial c = 0$ and choose b with $\beta_n(b) = c$.

Then $\beta_{n-1}(\partial b) = \partial(\beta_n(b)) = \partial c = 0$, hence

$$\partial b \in \ker(\beta_{n-1}) = \operatorname{im}(\alpha_{n-1}).$$

Choose $a \in C_c((\mathcal{G}|_{U_{12}})_{n-1}, A)$ with $\alpha_{n-1}(a) = \partial b$ and define

$$\partial_n([c]) := [a] \in H_{n-1}(\mathcal{G}|_{U_{12}}; A).$$

Standard homological algebra: ∂_n is well-defined, independent of choices, and yields exactness at $H_n(\mathcal{G}; A)$.

References I