

Problem Set: Waves

Karl Keshavarzi – ECE 105, Classical Mechanics

Part A: Wave Fundamentals and Mathematical Description

Problem 1: Wave Function Analysis

Consider a transverse wave on a string described by:

$$y(x, t) = A \sin(kx - \omega t + \phi)$$

where A is the amplitude, k is the wave number, ω is the angular frequency, and ϕ is the phase constant.

- (a) Derive the wave velocity v in terms of ω and k .
- (b) Show that the wave function satisfies the one-dimensional wave equation:

$$\frac{\partial^2 y}{\partial t^2} = v^2 \frac{\partial^2 y}{\partial x^2}$$

- (c) If a point on the string oscillates with maximum speed v_{max} , express v_{max} in terms of A and ω .
- (d) For $A = 0.05$ m, wavelength $\lambda = 2.0$ m, and period $T = 0.5$ s, calculate:
 - The wave number k
 - The angular frequency ω
 - The wave speed v
 - The maximum transverse speed of a point on the string

Problem 2: Wave Properties from Graphs

A travelling wave at $t = 0$ has the profile shown by the function:

$$y(x, 0) = A \sin(kx)$$

After time $t = t_1$, the wave has travelled a distance d to the right.

- (a) Write the wave function at time t_1 .

- (b) If the wave completes n full cycles in time t_0 , derive expressions for:
- The period T
 - The frequency f
 - The angular frequency ω
- (c) Show that the relationship between wave speed, wavelength, and frequency is $v = f\lambda$.
- (d) If $\lambda = 3.0$ m and the wave travels 15 m in 2.5 s, find the frequency.

Part B: Wave Velocity and Intensity

Problem 3: Waves on Strings

A transverse wave travels along a string with linear mass density μ under tension F_T .

- (a) Using dimensional analysis, show that the wave velocity must have the form:

$$v = C \sqrt{\frac{F_T}{\mu}}$$

where C is a dimensionless constant.

- (b) A string has mass m and length L . Express the wave velocity in terms of m , L , and F_T .
- (c) For a guitar string with $m = 2.0$ g, $L = 65$ cm, and tension $F_T = 80$ N, calculate:
- The linear mass density
 - The wave speed
 - The time for a wave to travel the length of the string

Problem 4: Sound Intensity and Decibels

A point source emits sound power P uniformly in all directions.

- (a) Show that the intensity I at distance r obeys the inverse square law:

$$I = \frac{P}{4\pi r^2}$$

- (b) The sound level in decibels (dB) is defined as $\beta = 10 \log_{10}(I/I_0)$, where $I_0 = 10^{-12}$ W/m² is the threshold of hearing. Show that doubling the distance from the source decreases the sound level by approximately 6 dB.
- (c) A speaker emits 50 W of acoustic power. Calculate the intensity and sound level at:
- $r = 1.0$ m
 - $r = 10.0$ m

Part C: Energy in Waves

Problem 5: Energy Transport

A sinusoidal wave $y(x, t) = A \sin(kx - \omega t)$ travels along a string with linear mass density μ .

- (a) Show that the kinetic energy per unit length is:

$$\frac{dK}{dx} = \frac{1}{2}\mu \left(\frac{\partial y}{\partial t} \right)^2$$

- (b) Derive the potential energy per unit length in terms of the string tension F_T and the slope $\partial y / \partial x$.

- (c) Show that the average total energy per unit length is:

$$\left\langle \frac{dE}{dx} \right\rangle = \frac{1}{2}\mu\omega^2 A^2$$

- (d) Find the power transmitted by the wave:

$$P = v \left\langle \frac{dE}{dx} \right\rangle$$

- (e) For a wave with $A = 2.0$ mm, $f = 100$ Hz, on a string with $\mu = 5.0$ g/m under tension $F_T = 50$ N, calculate:

- The wave speed
- The average energy per unit length
- The power transmitted

Part D: Wave Interference and Superposition

Problem 6: Two-Wave Interference

Two sinusoidal waves with the same amplitude A , wavelength λ , and frequency f travel in the same direction:

$$\begin{aligned} y_1(x, t) &= A \sin(kx - \omega t) \\ y_2(x, t) &= A \sin(kx - \omega t + \phi) \end{aligned}$$

- (a) Use the principle of superposition to find the resultant wave $y_T(x, t)$.
- (b) Show that the resultant can be written as:

$$y_T(x, t) = 2A \cos\left(\frac{\phi}{2}\right) \sin\left(kx - \omega t + \frac{\phi}{2}\right)$$

- (c) For what values of ϕ does:
- Constructive interference occur (maximum amplitude)?
 - Destructive interference occur (zero amplitude)?
- (d) If $\phi = \pi/3$, $A = 0.10$ m, $\lambda = 2.0$ m, and $v = 10$ m/s, find:
- The amplitude of the resultant wave
 - The frequency of oscillation
 - Write the complete wave function

Problem 7: Standing Waves on a String

Two waves of equal amplitude travel in opposite directions:

$$y_1(x, t) = A \sin(kx - \omega t)$$

$$y_2(x, t) = A \sin(kx + \omega t)$$

- (a) Show that the superposition creates a standing wave:

$$y(x, t) = 2A \sin(kx) \cos(\omega t)$$

- (b) Find the positions of nodes (where $y = 0$ always) and antinodes (maximum amplitude points).
- (c) A string of length L is fixed at both ends. Show that standing waves can only exist for wavelengths:

$$\lambda_n = \frac{2L}{n}, \quad n = 1, 2, 3, \dots$$

- (d) For a guitar string with $L = 65$ cm and wave speed $v = 400$ m/s:
- Find the fundamental frequency ($n = 1$)
 - Find the frequencies of the first three harmonics
 - Calculate the wavelength of the third harmonic

Problem 8: Temporal Interference

Two sound waves with slightly different frequencies f_1 and f_2 arrive at the same point.

- (a) Let $y_1 = A \cos(2\pi f_1 t)$ and $y_2 = A \cos(2\pi f_2 t)$. Use trigonometric identities to show:

$$y_{total} = 2A \cos(2\pi f_{mod} t) \cos(2\pi f_{avg} t)$$

where $f_{mod} = (f_1 - f_2)/2$ and $f_{avg} = (f_1 + f_2)/2$.

- (b) Explain why the "beat frequency" (frequency of amplitude maxima) is $f_{beat} = |f_1 - f_2|$, not f_{mod} .
- (c) Two tuning forks are struck. One has a frequency of 440 Hz. A beat frequency of 3 Hz is heard. What are the possible frequencies of the second fork?

Part E: Wave Reflection and Transmission

Problem 9: Boundary Conditions

A wave pulse travels along a string and encounters a boundary where the string properties change.

- (a) State the two boundary conditions that must be satisfied at the junction between two strings with different linear mass densities μ_1 and μ_2 .
- (b) For an incident wave $y_i = A_i \sin(k_1 x - \omega t)$ in medium 1, write expressions for:
- The reflected wave y_r
 - The transmitted wave y_t
- (c) Derive the amplitude reflection coefficient:

$$R = \frac{A_r}{A_i} = \frac{v_1 - v_2}{v_1 + v_2}$$

- (d) Show that the power is conserved: $P_i = P_r + P_t$
- (e) A wave travels from a light string ($\mu_1 = 2.0$ g/m) to a heavy string ($\mu_2 = 8.0$ g/m) under the same tension. Calculate:
- The ratio of wave speeds v_1/v_2
 - The reflection coefficient
 - The percentage of incident power transmitted

Problem 10: Standing Waves in Air

Sound waves can form standing waves in pipes.

- (a) **Open-Open Pipe:** A pipe of length L is open at both ends.
- Explain why there must be displacement antinodes at both ends.
 - Show that this implies the length must be an integer number of half-wavelengths: $L = n \frac{\lambda}{2}$.
 - Derive the frequency formula: $f_n = n \frac{v}{2L}$ for $n = 1, 2, 3 \dots$
- (b) **Open-Closed Pipe:** A pipe is open at one end and closed at the other.
- Explain why there is an antinode at the open end and a node at the closed end.
 - Show that this implies the length must be an odd integer number of quarter-wavelengths: $L = m \frac{\lambda}{4}$ (where m is odd).
 - Derive the frequency formula: $f_n = n \frac{v}{4L}$ for $n = 1, 3, 5 \dots$
- (c) An organ pipe is 3.0 m long. Calculate the fundamental frequency if:
- Both ends are open ($v_{\text{sound}} = 343$ m/s).
 - One end is closed.

Part F: Advanced and Optional Topics

Problem 11: Wave Packets and Group Velocity

A wave packet is formed by superposing waves with slightly different frequencies:

$$y(x, t) = A \sin(k_1 x - \omega_1 t) + A \sin(k_2 x - \omega_2 t)$$

(a) Show that this can be written as:

$$y(x, t) = 2A \cos\left(\frac{\Delta k}{2}x - \frac{\Delta \omega}{2}t\right) \sin(\bar{k}x - \bar{\omega}t)$$

where $\Delta k = k_2 - k_1$, $\Delta \omega = \omega_2 - \omega_1$, $\bar{k} = (k_1 + k_2)/2$, $\bar{\omega} = (\omega_1 + \omega_2)/2$.

(b) Identify the envelope (modulation) and carrier wave components.

(c) Define and derive the group velocity $v_g = \frac{d\omega}{dk}$.

(d) For waves in deep water where $\omega = \sqrt{gk}$ (where g is gravitational acceleration):

- Find the phase velocity $v_p = \omega/k$
- Find the group velocity v_g
- Show that $v_g = v_p/2$

(e) For $g = 9.8 \text{ m/s}^2$ and $\lambda = 100 \text{ m}$, calculate both velocities.

Problem 12: Doppler Effect

A source emitting waves of frequency f_s moves with velocity v_s toward a stationary observer. The wave speed in the medium is v .

(a) Derive the observed frequency:

$$f_o = f_s \left(\frac{v}{v - v_s} \right)$$

(b) What happens when v_s approaches v ?

(c) For a moving observer with velocity v_o toward a stationary source, show:

$$f_o = f_s \left(\frac{v + v_o}{v} \right)$$

(d) Derive the general Doppler formula for both source and observer in motion.

(e) An ambulance siren emits sound at $f_s = 1000 \text{ Hz}$. The ambulance approaches at 30 m/s . Given $v_{\text{sound}} = 343 \text{ m/s}$:

- Calculate the frequency heard as it approaches
- Calculate the frequency heard as it recedes

- Find the frequency shift

Problem 13: Sound Speed in Gases

Sound waves in a gas are longitudinal waves with speed:

$$v = \sqrt{\frac{\gamma P}{\rho}}$$

where γ is the heat capacity ratio, P is pressure, and ρ is density.

- Show that this can also be written as $v = \sqrt{\frac{\gamma RT}{M}}$ for an ideal gas.
- Calculate the speed of sound in air at 20°C given $\gamma = 1.40$ and $M = 28.97$ g/mol.

Problem 14: The Shock Wave (Sonic Boom)

Consider a source moving at velocity $v_s > v_{sound}$.

- Explain qualitatively the formation of a shock wave (Mach cone).
- Show that the half-angle of the Mach cone is given by $\sin \theta = \frac{v_{sound}}{v_s} = \frac{1}{\text{Mach Number}}$.
- A jet flies at Mach 2.0. Calculate the cone angle.

Numerical Answers

Problem 1(d): $k = \pi \text{ rad/m}$, $\omega = 4\pi \text{ rad/s}$, $v = 4.0 \text{ m/s}$, $v_{max} = 0.63 \text{ m/s}$

Problem 2(d): $f = 2.0 \text{ Hz}$

Problem 3(d): $\mu = 3.08 \times 10^{-3} \text{ kg/m}$, $v = 161 \text{ m/s}$, $t = 4.0 \text{ ms}$

Problem 4(c): $I_1 = 3.98 \text{ W/m}^2$ (126 dB); $I_{10} = 0.04 \text{ W/m}^2$ (106 dB)

Problem 5(e): $v = 100 \text{ m/s}$, $\langle dE/dx \rangle = 0.079 \text{ J/m}$, $P = 7.9 \text{ W}$

Problem 6(d): Amplitude = 0.173 m , $f = 5.0 \text{ Hz}$

Problem 7(e): $f_1 = 308 \text{ Hz}$, $f_2 = 615 \text{ Hz}$, $f_3 = 923 \text{ Hz}$, $\lambda_3 = 0.43 \text{ m}$

Problem 8(c): 437 Hz or 443 Hz

Problem 9(e): $v_1/v_2 = 2$, $R = 1/3$, $T = 4/3$, 89% transmitted

Problem 10(c): Open-Open: 57.2 Hz; Open-Closed: 28.6 Hz

Problem 11(e): $v_p = 12.5 \text{ m/s}$, $v_g = 6.25 \text{ m/s}$

Problem 12(e): $f_{approach} = 1096 \text{ Hz}$, $f_{recede} = 920 \text{ Hz}$, shift = 176 Hz