🗺️ Map of *The Inward Bow of a Real-Rooted Polynomial: Newton's Inequalities, Machine-Checked in Lean 4*

🏷️ **Inequalities for polynomials · real-rooted polynomials and log-concavity**

1 results | ✅ **1** formalized in Lean | ◐ **0** with a part in Lean | 💻 **1** checked by a program | written proof only (a result can carry several tags)

Dependencies in the written proof: an arrow A → B means that this proof of B uses A. Every result rests on its written proof; the tags record what was checked besides, and each statement's "Evidence & scope" block says how far. The colour of a box is its strongest tag, in the order above; the tags in the box give the rest. Boxes with a thick border are the main results; click a box to read the statement.

Q1: what does reality of the roots force on the coefficients?

Theorem 1
Newton's inequalities for a monic real-rooted polynomial, and log-concavity of the normalized means
✅ Lean · 💻 program

> \begin{itemize} \item A \texttt{sorry}-free Lean~4 proof of Newton's inequalities $e_k^2\binom{n}{k-1}\binom{n}{k+1}\ge e_{k-1}e_{k+1}\binom{n}{k}^2$ for real-rooted real polynomials (Theorem~1), and of the log-concavity of the normalized means $p_k=e_k/\binom nk$; the symmetric-function form assumes p monic, the coefficient form \texttt{newton_inequality_coeff} assumes nothing. \item The route is the classical one (Sylvester; Hardy--Littlewood--P\'olya): differentiate, reverse, differentiate again, down to a real-rooted quadratic, whose discriminant is the inequality once positive factorials are cleared. \item Four reusable blocks: real-rootedness under differentiation, iterated differentiation and reversal (the constant-term hypothesis dropped), and the coefficient-form discriminant for $\deg\le2$, which also absorbs the degenerate case $a_{i-1}=0$. \item The mathematics is classical; the paper claims the formalization (to the author's knowledge the first in any proof assistant, June 2026), not a new inequality. Equality cases and Maclaurin's chain are named, not proved. \end{itemize}

🚀 **Start here: Theorem 1**: Newton's inequalities for a monic real-rooted polynomial, and log-concavity of the normalized means.

Main results:

- **Theorem 1**: Newton's inequalities for a monic real-rooted polynomial, and log-concavity of the normalized means

> ## 🧭 Fields: where the paper lives and what it connects
>
> **Tools taken from other branches**
>
> - **Real-rootedness preservers** → Rolle's theorem for polynomials (real roots survive differentiation) and reversal x ↦ 1/x — Theorem 1
> - **Symmetric functions** → Vieta's relations between coefficients and the elementary symmetric functions of the roots (Mathlib's Multiset.esymm) — Theorem 1
> - **Formal verification** → Lean 4 + Mathlib: falling factorials of iterated derivatives, exact degrees in characteristic zero — Theorem 1
>
> **What the results serve**
>
> - **Log-concavity in combinatorics** ← real-rooted ⇒ log-concave coefficients, e.g. the matching numbers through Heilmann–Lieb (named in §4, not composed here) — Theorem 1
> - **Maclaurin's inequalities and the equality cases** ← the next targets the paper names; not proved