Every 4×4 biquadratic sum of squares is a sum of eleven squares

Real algebraic geometry · sums of squares on projective varieties

Carles Marín

Statements

Theorem 1 3
Lemma 2 4
Lemma 3 6
Lemma 4 — a regular point 7
Proposition 5 7
Proposition 6 8
Remark 7 9
Proposition 8 9

At a glance

> The exact value **BSR(4,4) = 11**: every biquadratic sum of squares in $x_{1..4}, y_{1..4}$ is a sum of eleven squares of bilinear forms. Qi and Cui gave a form that needs eleven; the previous upper bound was twelve. The proof classifies the Hankel multipliers of rank at most five whose image contains no product (Lemmas 2 and 3) and rules out the two shapes left with second variations built from the alternating forms (Propositions 6 and 8).

🚀 **Start here: Theorem 1**: BSR(4,4) = 11.

Main results:


- **Theorem 1**: BSR(4,4) = 11

8 results | ✅ **0** entirely in Lean | ◐ **5** partly in Lean

💻 **5** checked by a program | written proof only | (a result can carry several tags)

🗺️ Map of the results (click a box to go to it)

Dependencies in the written proof: an arrow A → B means that this proof of B uses A. Every result rests on its written proof; the tags record what was checked besides, and each statement's "Evidence & scope" block says how far. The colour of a box is its strongest tag, in the order above; the tags in the box give the rest. Boxes with a thick border are the main results; click a box to read the statement.



📐 Setting (copied from the paper, §§1–2)

Write $x = (x_1, \dots, x_4)$, $y = (y_1, \dots, y_4)$, $V = \mathbb{R}^4 \otimes \mathbb{R}^4$ (the bilinear forms) and R_2 the biquadratic forms, of dimension 100. $\mathrm{BSR}(m, n)$ is the least r such that every biquadratic sum of squares in $m + n$ variables is a sum of r squares of bilinear forms; $\Phi(h_1, \dots, h_{11}) = h_1^2 + \cdots + h_{11}^2$, $\Sigma_{11} = \Phi(V^{11})$ and Σ the cone of sums of squares. A *multiplier* is a symmetric bilinear form H on $A \otimes B$ with the Hankel symmetry

$H(x \otimes b, y \otimes c) = H(x \otimes c, y \otimes b)$; its *supports* a, b_0 are the dimensions of the two factors modulo the radicals. Writing $H = T^{*}JT$ with T onto E, $\dim E = \operatorname{rk} H$, and $T_b x = T(x \otimes b)$, the symmetry reads $\langle T_b x, T_c y\rangle_J = \langle T_c x, T_b y\rangle_J$.

1 The result

Theorem 1. p. 1

> $\mathrm{BSR}(4,4) = 11$.

Isabelle · ours · part **show**

Only a part is in Isabelle: partly: the regular point, the support count of Lemma 3, the polar multiplier of Lemma 2 (iii) and the trace count of Proposition 8; Lemmas 2 and 3 themselves, Proposition 5 and the second variations are not formalized

`regular_point_symmetric` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 25 — the regular point (Lemma 4)

```
theorem regular_point_symmetric:
  fixes H :: "nat \<Rightarrow> nat \<Rightarrow> nat \<Rightarrow> nat \
<Rightarrow> real" and n :: nat
  assumes sym: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c
< n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H c d a b"
    and hankel: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow>
c < n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H a d c b"
    and van: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c <
n \<Longrightarrow> d < n \<Longrightarrow> H a b c d + H b a c d = 0"
    and idx: "a < n" "b < n" "c < n" "d < n"
  shows "H a b c d = 0"
```

`support_triples` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 54 — the count closing Lemma 3

```
lemma support_triples:
  fixes a b r :: nat
  assumes "2 \<le> a" "a \<le> 4" "2 \<le> b" "b \<le> 4" "r \<le> 5" "r \<le>
(a - 1) * (b - 1)" "a + 1 \<le> r" "b + 1 \<le> r"
  shows "(a, b, r) \<in> {(4, 4, 5), (4, 3, 5), (3, 4, 5), (3, 3, 4)}"
```

`trace_count` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 92 — the trace count of Proposition 8

```
lemma trace_count:
  fixes lam :: "'i \<Rightarrow> real" and I :: "'i set"
  assumes fin: "finite I" and le1: "\<And>i. i \<in> I \<Longrightarrow> lam i \
<le> 1"
    and rk: "card {i \<in> I. lam i \<noteq> 0} \<le> 4" and neg: "j \<in> I"
"lam j < 0"
  shows "(\<Sum>i\<in>I. lam i) < 3"
```

Trust base: ⚠️ not checked. Written proof, audited (V244, V246); the theorem as a whole is not machine-checked: Isabelle/HOL checks, without sorry, four elementary steps (regular point, support count, polar, trace count)

> 🆕 **Classical and new:** the paper: the lower bound 11 is [QC, Theorem 3.1]; the previous upper bound 12 follows from BSR(m,3) = 2m [Mar, Corollary 7].

↳ Not used by another result of the paper.

Take a biquadratic form in $x_1, \dots, x_4$ and $y_1, \dots, y_4$ that is a sum of squares of bilinear forms, however many. The theorem says that eleven squares always suffice, and Qi and Cui gave a form for which ten do not. Before this note the best upper bound was twelve, from width three. The proof does not search for representations: it shows that the set of sums of eleven squares has no boundary inside the cone, because any boundary point would carry a Hankel multiplier of rank at most five that the classification of Lemmas 2–3 and the second variations of Propositions 6 and 8 rule out.

Evidence & scope

- **Program:** perturbations P + εb² of the Qi–Cui witness: approximate representations with eleven squares in all 312 cases. `Curiosity/research/cohete-sat/z2/bsr44/perturbar_qc.py`

2 Hankel multipliers of small rank

Lemma 2. p. 2

> *There is no multiplier on $\mathbb{C}^4 \otimes \mathbb{C}^4$ of rank 5, with supports $a = b_0 = 4$, whose image contains no simple tensor.*

Isabelle · ours · part **show**

Only a part is in Isabelle: only the last step of case (iii): the polar multiplier of (xᵀb)² is injective (polar_trace_injective)

`polar_trace_injective` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 67 — case (iii): X ↦ (tr(X) I + Xᵀ)/2 is injective, so the polar of (xᵀb)² has rank 16, not 5

```
lemma polar_trace_injective:
  fixes X :: "nat \<Rightarrow> nat \<Rightarrow> real" and n :: nat
  assumes zero: "\<And>i j. i < n \<Longrightarrow> j < n \<Longrightarrow> (if
i = j then (\<Sum>k<n. X k k) else 0) + X j i = 0"
    and ij: "i < n" "j < n"
  shows "X i j = 0"
```

↳ Used in: Theorem 1 (p. 3), Lemma 3 (p. 6), Proposition 6 (p. 8)

Proof

The three cases below are exhaustive: $s = 4$ with some $Q(b)$ invertible, $s = 4$ with every $Q(b)$ singular, and $s \le 3$.

(i) Some $Q(b)$ is invertible. Choosing bases by congruence, $E = \mathbb{C}^4 \oplus \mathbb{C}$ with the standard symmetric product and $T_b x = (x, 0)$. In a basis of a complement of b write $T_i x = (A_i x,\, u_i^{\top} x)$, $i = 1, 2, 3$. By (2),

$$A_i^{\top} = A_i, \qquad [A_i, A_j] = u_j u_i^{\top} - u_i u_j^{\top}. \tag{1}$$

With $v \odot w = vw^\top + wv^\top$ and $K_{ij} = A_i u_j - A_j u_i$, the Jacobi identity for A_1, A_2, A_3 and (4) give

$$K_{12} \odot u_3 + K_{23} \odot u_1 + K_{31} \odot u_2 = 0. \tag{2}$$

If the u_i span 0, T is not onto. If they span a line, the last coordinate of T is a simple covector of $\operatorname{im} H$. If they span a plane, after a change of basis of the complement $u_3 = 0$ with u_1, u_2 independent; then A_3 commutes with A_1, A_2, and (5) gives $(A_3 u_1) \odot u_2 = (A_3 u_2) \odot u_1$, whence $A_3 u_1 = t u_1$, $A_3 u_2 = t u_2$ for one t (compare coefficients in a basis starting with u_1, u_2). The space $P = \operatorname{im}(A_3 - tI)$ is non-zero (otherwise $T_3 = tT_b$ and B would have a radical, against $b_0 = 4$), stable under all A_i, and orthogonal to u_1, u_2 since A_3 is symmetric; by (4) the A_i commute on P, and a common eigenvector $x \in P$ has $T_c x \in \mathbb{C}(x, 0)$ for all c, so the Observation applies. If the u_i are independent, (5) forces $K_{ij} = t_i u_j - t_j u_i$ for three scalars t_i (complete u_1, u_2, u_3 to a basis: the terms with the fourth vector and the squares u_i^2 force $K_{12} \in \langle u_1, u_2 \rangle$, $K_{23} \in \langle u_2, u_3 \rangle$, $K_{31} \in \langle u_3, u_1 \rangle$, and the coefficients of $u_1 u_2$, $u_2 u_3$, $u_3 u_1$ give the signs). Then the symmetric matrices

$$\widehat{A}_i = \begin{pmatrix} A_i & u_i \\ u_i^\top & t_i \end{pmatrix}$$

commute (the upper-left block of their commutator vanishes by (4), the upper right one is $K_{ij} + t_j u_i - t_i u_j = 0$, and the rest follows by antisymmetry). A common eigenvector $z \neq 0$ exists over $\mathbb{C}$; if $\widehat{A}_i z = \lambda_i z$, then $\langle z, T_i x \rangle = \lambda_i \langle z, T_b x \rangle$, since $T_i x = \widehat{A}_i T_b x$, so $T^* J z$ is simple, and non-zero because T is onto.

(ii) Some T_b has rank 4 and every $Q(b)$ is singular. Then $U = \operatorname{im} T_b$ is a degenerate hyperplane with a radical of dimension one. In a Witt basis $E = \mathbb{C}^3 \oplus \langle e, f \rangle$, $\langle e, f \rangle = 1$, $\langle e, e \rangle = \langle f, f \rangle = 0$, with $T_b(w, z) = (w, z, 0)$, condition (2) gives each T_c the shape $\begin{pmatrix} A & p \\ l^\top & t \\ p^\top & q_0 \end{pmatrix}$ with $A = A^\top$. In $Q(b + \tau c)$ the upper 3×3 block is $I + O(\tau)$, the upper right block is $2\tau p + O(\tau^2)$ and the lower right entry is $2\tau q_0 + \tau^2 (p^\top p + 2 t q_0)$. Since the determinant vanishes identically, so does the Schur complement as a formal series; its terms of order one and two give $q_0 = 0$ and $-3 p^\top p = 0$. Thus $p(c)^\top p(c) = 0$ for every c, and $c \mapsto p(c)$ is linear; *by polarization* $p(c)^\top p(c') = 0$ for all c, c', so $\{p(c)\}$ spans a totally isotropic subspace of $\mathbb{C}^3$, of dimension at most one: $p(c) = \beta(c) p_0$. The covector $T^* J e$, $(w, z; c) \mapsto p(c)^\top w = \beta(c)\, p_0^\top w$, is simple and non-zero.

(iii) $s \leq 3$. By (3) and $Q(b) \neq 0$ generically, $s = 3$ and $\operatorname{rk} Q(b) = 1$ generically: U has a radical of dimension at least two and $Q(b) \neq 0$. So $q(x, b) = x^\top Q(b) x$ is, over $\mathbb{C}(b)$, a scalar times the square of a linear form in x (complete the square from a non-zero diagonal entry). By Gauss' lemma $q = g(b) L(x, b)^2$ in $\mathbb{C}[b][x]$ with L primitive, and the factors of a bihomogeneous polynomial are bihomogeneous. If L has degree 0 in b, the support in A is 1. Otherwise g is constant and $L = x^\top P b$, with P invertible by full support, so after changes of basis and scale $q = (x^\top b)^2$. Its polar multiplier acts on

matrices as $X \mapsto (\operatorname{tr}(X)I + X^\top)/2$, which is invertible ($X^\top = -\operatorname{tr}(X)I$ gives $5 \operatorname{tr} X = 0$ and $X = 0$): rank $16 \neq 5$.

Evidence & scope

- **Program:** the Jacobi identity (5), the Schur expansion of case (ii) and the rank 16 of the polar of $(x^{\mathrm{T}}b)^2$, in exact arithmetic. `Curiosity/research/cohete-sat/z2/bsr44/auditoria_V244/simbolico.py`

Lemma 3. p. 3

Let H be a nonzero multiplier on $\mathbb{C}^4 \otimes \mathbb{C}^4$ of rank $r \leq 5$ whose image contains no simple tensor. Up to exchanging the factors, (a, b_0, r) is either $(3, 4, 5)$ or $(3, 3, 4)$.

Isabelle · ours · part **show**

Only a part is in Isabelle: the closing count only: from 2 ≤ a, b₀ ≤ 4, r ≤ 5, r ≤ (a−1)(b₀−1) and r ≥ max(a,b₀) + 1 only four triples remain (support_triples)

`support_triples` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 54 — the arithmetic that closes Lemma 3

```
lemma support_triples:
  fixes a b r :: nat
  assumes "2 \<le> a" "a \<le> 4" "2 \<le> b" "b \<le> 4" "r \<le> 5" "r \<le>
(a - 1) * (b - 1)" "a + 1 \<le> r" "b + 1 \<le> r"
  shows "(a, b, r) \<in> {(4, 4, 5), (4, 3, 5), (3, 4, 5), (3, 3, 4)}"
```

↳ Used in: Theorem 1 (p. 3)

Proof

Restrict H to $A' \otimes B'$, with A', B' complements of the two radicals: the rank, the symmetry (1) and the absence of simple tensors are kept (a simple covector of the restricted image extends by zero on the radicals and stays simple and in $\operatorname{im} H$), and now both radicals are zero, with $\dim A' = a$, $\dim B' = b_0$. Since $\operatorname{im} H$ annihilates the radicals, $\mathbb{P}(\operatorname{im} H)$ is a subspace of dimension $r - 1$ avoiding the Segre variety of the effective factors, so $r \leq (a-1)(b_0-1)$; and $a, b_0 \geq 2$ since $H \neq 0$.

We show $r \geq a + 1$, and by the symmetry of (1) in the two factors also $r \geq b_0 + 1$. Suppose $r \leq a$ (≤ 4) and take b generic, with $s = \operatorname{rk} T_b$ maximal and $Q(b) \neq 0$. If $s<a$, then $\ker T_b \neq 0$, and the radical of $U = \operatorname{im} T_b$ has dimension at most $s - 1$ (as $Q(b) \neq 0$) and at most $r - s$ (as it lies in $U^{\perp_J}$), so at most $\min(s-1, r-s) \leq 1$ because $r \leq 4$; this contradicts (3) and the Observation. If $s = a$, then $s = a = r$, $T_b : A' \to E$ is an isomorphism, $Q(b)$ is non-degenerate, and the operators $T_b^{-1} T_c$ are self-adjoint for $Q(b)$ and, by (2), commute; a common eigenvector x has $T_c x \in \mathbb{C}\, T_b x$ for all c, and the Observation gives a simple tensor. So $r \geq \max(a, b_0) + 1$.

With $r \leq 5$ and $r \leq (a-1)(b_0-1)$ the only possibilities are $(a, b_0, r) = (4, 4, 5)$, $(4, 3, 5)$, $(3, 4, 5)$ and $(3, 3, 4)$, and Lemma 2 excludes $(4, 4, 5)$.

Evidence & scope

- **Program:** Newton search for rank-five Hankel multipliers with support 3×4 and product-free image: 14 numerical candidates. `Curiosity/research/cohete-sat/z2/bsr44/auditoria_V244/numerico.py`

3 The boundary multiplier

Lemma 4. (a regular point) p. 4

> *Identify both factors with $\mathbb{R}^4$. Let $S \subset V$ be the span of the four forms $x_i y_i$ and the six forms $x_i y_j + x_j y_i$ for $i<j$. If H is a Hankel multiplier and $S \subseteq \ker H$, then $H = 0$. Consequently $d\Phi$ is surjective at the tuple consisting of these ten forms and zero.*

Isabelle · ours · part **show**

Only a part is in Isabelle: the linear algebra; the duality from there to «dΦ onto» is on paper

`regular_point_symmetric` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 25 — a symmetric Hankel multiplier that vanishes on the symmetric tensors in its first argument is zero (in any size n)

```
theorem regular_point_symmetric:
  fixes H :: "nat \<Rightarrow> nat \<Rightarrow> nat \<Rightarrow> nat \
<Rightarrow> real" and n :: nat
  assumes sym: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c
< n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H c d a b"
    and hankel: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow>
c < n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H a d c b"
    and van: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c <
n \<Longrightarrow> d < n \<Longrightarrow> H a b c d + H b a c d = 0"
    and idx: "a < n" "b < n" "c < n" "d < n"
  shows "H a b c d = 0"
```

↳ Used in: Theorem 1 (p. 3), Proposition 5 (p. 7)

Proof

> If $S \subseteq \ker H$, then $H(s,v) = H(v,s) = 0$ for all $s \in S$ and $v \in V$, and S is the space of symmetric tensors, so $H(x \otimes y, x \otimes y) = -H(x \otimes y, y \otimes x) = -H(x \otimes x, y \otimes y) = 0$ for all x, y, and $H = 0$ by polarization. Hence the products of the ten summands with V span R_2 (we also checked the rank 100 exactly), and adding an eleventh summand, even zero, gives a point where $d\Phi$ is onto.

Evidence & scope

- **Program:** rank 100 of the products of the ten summands with V, in exact arithmetic; control with the four diagonal summands only: 58. `Curiosity/research/cohete-sat/z2/bsr44/auditoria_V244/punto_regular.py`

Proposition 5. p. 4

> *If $\Sigma_{11} \neq \Sigma$, there are $F \in \operatorname{int}\Sigma \cap \partial\Sigma_{11}$, a representation $F = \sum_{i=1}^{11} h_i^2$ and an indefinite real multiplier H on V such that, with $W = \operatorname{span}(h_i)$: $W \subseteq \ker H$, $\dim W = 11$ and hence $\operatorname{rk} H \leq 5$; the complex image of H contains no simple tensor; and $\sum_i h_i D_i = 0$ in R_2 implies $\sum_i H(D_i, D_i) \geq 0$.*

NEW **Classical and new:** the paper: the argument of [Mar, Proposition 3 and Corollary 4], after [BDS]; only the jet count $100 - 14 + 12 = 98 < 99$ depends on the size.

↳ Used in: Theorem 1 (p. 3), Proposition 6 (p. 8), Proposition 8 (p. 9)

Proof

We adapt the argument of [3 , Proposition 3 and Corollary 4] to $(m,n)=(4,4)$, using Lemma 4. The only count that depends on the size is the one excluding forms with a singular point with no real factor. Take a pair of conjugate points distinct in both factors and change coordinates over $\mathbb{C}$ to $(e_1,e_1),(e_2,e_2)$. A first jet has seven coordinates, the value and the six tangent derivatives, and the two first jets use disjoint sets of seven monomial coefficients, so the joint jet map is surjective over $\mathbb{C}$. Conjugation exchanges the two jets, and the restriction to real forms has real rank fourteen. The point varies in real dimension twelve. The incidence therefore has dimension at most $100-14+12=98$, and its projection cannot fill the boundary patch of dimension 99. A singular point with a real factor is impossible at an interior form, by definiteness of the quadratic form in the other factor. Independence of the h_i is as in [3 , Corollary 4] .

4 Support 3×4, rank five

Proposition 6. p. 4

No triple (F,h,H) satisfying the conclusions of Proposition 5 can have $(a,b_0,r)=(3,4,5)$ or $(4,3,5)$.

↳ Used in: Theorem 1 (p. 3), 7 (p. 9)

Proof

Here $\dim\ker H=16-5=11$, so $W=\ker H$. Let x_0 span the radical in the first factor and A' be a real complement, $\dim A'=3$. Then $x_0\otimes B\subseteq W$, and an orthogonal change of the eleven summands (an orthonormal basis of the preimage of $x_0\otimes B$, completed) gives $h_i=x_0\otimes y_i$, $1\le i\le 4$, for a real basis (y_i) of B. For $z_{ij}=-z_{ji}\in A'$ put

$$D_i=\sum_{j=1}^{4} z_{ij}\otimes y_j \quad (i\le 4), \qquad D_i=0 \quad (i{>}4).$$

Then $\sum_i h_i D_i=\sum_{i,j} x_0 z_{ij}\, y_i y_j=0$ by antisymmetry: these are syzygies, a space $A'\otimes\Lambda^2 B$ of dimension 18. With $H=T^{*}JT$ as in §2 (now over $\mathbb{R}$, $\dim E=5$), the second variation is $\sum_i H(D_i,D_i)=J^{\oplus 4}(\Psi z)$ for

$$\Psi : A'\otimes\Lambda^2 B\to E^4, \qquad \Psi(z)=\Big(\sum_j T_{y_j} z_{ij}\Big)_{i=1}^{4}.$$

By Proposition 5, $J^{\oplus 4}$ is non-negative on $\operatorname{im}\Psi$. Since H is indefinite, the positive index p of J is at most 4, and a subspace on which $J^{\oplus 4}$ is non-negative has dimension at most $4p\le 16$. Hence $\operatorname{rk}\Psi\le 16$.

On the other hand $\operatorname{rk}\Psi=17$. Complexify and compute the annihilator of $\operatorname{im}\Psi$ under $J^{\oplus 4}$: a tuple $(c_i)\in E^4$, i.e. a map $C:B\to E$ with $Cy_i=c_i$, annihilates $\operatorname{im}\Psi$ if and only if the matrix $\big(\langle Cy_i, T(a\otimes y_j)\rangle_J\big)_{ij}$ is symmetric for every $a\in A'$ (vary each z_{ij},

$i<j$, separately). The three maps $C = T(a \otimes \cdot)$, a in a basis of A', are independent and lie in the annihilator by (2). If there were a fourth independent C, it would extend T to $\widetilde{T} : \mathbb{C}^4 \otimes B \to E$ satisfying (2): the identities involving the new row are exactly the symmetry just written, and the one of the new row with itself is automatic. Then $\widetilde{T}^* J \widetilde{T}$ is a multiplier of rank 5 (as T is onto) with supports 4 and 4 (by independence of the new row, and since the second support was already full), and its image contains no simple tensor: a simple covector in it restricts to $A' \otimes B$ as a simple covector of im H, excluded by Proposition 5, or as zero, and then its representing vector z has $\langle z, T(A' \otimes B)\rangle_J = 0$, so $z = 0$. This contradicts Lemma 2. So the annihilator has dimension 3 and $\operatorname{rk} \Psi = 20 - 3 = 17 > 16$, a contradiction.

Evidence & scope

- **Program:** on the rank-five multiplier with support 3×4 of the supplement of [Mar]: rank $\Psi = 17 > 16$ and a negative value of the second variation; in the 14 numerical candidates rank $\Psi = 17$, and 15 in a control built from five product terms. `Curiosity/research/cohete-sat/z2/bsr44/v244_comprobar.py`

Remark 7. p. 5

One triple of quaternionic structures does not suffice: for $z_{ij} = \sum_a \eta^a_{ij} v_a$ with the 't Hooft symbols η^a, the second variation is $\sum_a v_a^\top (\sum_j H_{jj}) v_a$, because $(\eta^a)^\top \eta^b$ has symmetric part $\delta_{ab} I$ and $H_{jk} = H_{kj}$; it only says that $\sum_j H_{jj} \succeq 0$. The argument above uses the whole of $\Lambda^2 B$.

↳ Not used by another result of the paper.

5 Support 3×3, rank four

Proposition 8. p. 5

No triple (F, h, H) satisfying the conclusions of Proposition 5 can have $(a, b_0, r) = (3, 3, 4)$.

Isabelle · ours · part **show**

Only a part is in Isabelle: the final trace count of the case u ∉ W (trace_count); the two second variations are on paper

`trace_count` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 92 — eigenvalues at most 1, at most four non-zero, one negative: their sum is less than 3

```
lemma trace_count:
  fixes lam :: "'i \<Rightarrow> real" and I :: "'i set"
  assumes fin: "finite I" and le1: "\<And>i. i \<in> I \<Longrightarrow> lam i \
<le> 1"
    and rk: "card {i \<in> I. lam i \<noteq> 0} \<le> 4" and neg: "j \<in> I"
"lam j < 0"
  shows "(\<Sum>i\<in>I. lam i) < 3"
```

↳ Used in: Theorem 1 (p. 3)

Proof

Let x_0, y_0 span the real radicals of the two factors, A', B' complements and $u = x_0 \otimes y_0$. The space $U = x_0 \otimes B + A \otimes y_0$ has dimension 7 and lies in $K = \ker H$, $\dim K = 12$; W is a hyperplane of K, and $K = U \oplus K_0$ with $K_0 = K \cap (A' \otimes B')$.

Case $u \notin W$. Then $W \cap (x_0 \otimes B) = x_0 \otimes L$ with $\dim L = 3$ and L projecting isomorphically onto $B/\langle y_0 \rangle$. Orthogonalizing on the preimage of this intersection, $h_i = x_0 \otimes y_i$, $i \le 3$, for a basis of L. The restriction of H to $A' \otimes L$ keeps the indices, the rank four and the support three in the first factor. For $v_1, v_2, v_3 \in A'$ put $D_i = \sum_{j,p} \varepsilon_{ijp}\, v_p \otimes y_j$ for $i \le 3$ and $D_i = 0$ otherwise; they are syzygies by antisymmetry, and with H_{jl} the blocks of the restriction and $S = \sum_j H_{jj}$, Proposition 5 and the contraction of the alternating symbols give, as in [3 , § 5] ,

$$I_3 \otimes S - H|_{A' \otimes L} \succeq 0.$$

The diagonal blocks sum to $2S$, so $S \succeq 0$; if $Sx = 0$, the three diagonal values of this positive semidefinite form at $x \otimes y_j$ sum to zero, so each $x \otimes y_j$ is in its kernel, $H(x \otimes y_j, \cdot) = 0$, and $x = 0$ by full support in A'. So $S \succ 0$, and after the congruence by $S^{-1/2}$ in the three blocks the matrix of H has trace 3, all eigenvalues at most 1, rank four and at least one negative eigenvalue: its trace is less than 3, a contradiction.

Case $u \in W$. An orthogonal change makes h_1 a non-zero multiple of u. Let ℓ be a real linear functional on K with $W = \ker \ell$. For $v = \sum v_{ij} a_i \otimes b_j \in A' \otimes B'$ (bases a_i, b_j) the lift $P(v) = \sum_{ij} v_{ij} (a_i \otimes y_0)(x_0 \otimes b_j) \in \operatorname{Sym}^2 U$ has product uv in R_2. If $\ell|_U = 0$ all these products lie in WK; otherwise $\operatorname{Sym}^2 U / (U \cap W) U$ is one-dimensional and the class is measured by $(\ell|_U)^2$. So $uv \in WK$ whenever

$$\phi(v) = \sum_{ij} v_{ij}\, \ell(a_i \otimes y_0)\, \ell(x_0 \otimes b_j) = 0.$$

The functional ϕ on the active block is zero or simple. If it is non-zero it does not vanish on K_0, since otherwise it would lie in the image of the active part of H, which contains no simple tensor. Choose v in the active block with $H(v, v) < 0$; adding a real element of K_0 keeps this value and achieves $\phi(v) = 0$. Then $h_1 v \in WK$: write $-h_1 v = \sum_i h_i k_i$ with $k_i \in K$. The $D_i = \delta_{i1} v + k_i$ form a syzygy with second variation $H(v, v) < 0$, a contradiction.

6 Checks and status

No numbered statement in this section.

It is prose only (no statement, figure or table): read it in the paper (PDF).

The companion site

No numbered statement in this section.

It is prose only (no statement, figure or table): read it in the paper (PDF).

Acknowledgement

No numbered statement in this section.

It is prose only (no statement, figure or table): read it in the paper (PDF).

References

1

G. Blekherman, A. Dunbar, R. Sinn, Pythagoras numbers for ternary forms, arXiv:2410.17123v2.

2

L. Chen, D. Ž. Đoković, Properties and construction of extreme bipartite states having positive partial transpose, Comm. Math. Phys. 323 (2013), 241–284; arXiv:1203.1364.

3

C. Marín, For $m \geq 2$, every $m \times 3$ biquadratic sum of squares is a sum of $2m$ squares, Zenodo, doi:10.5281/zenodo.24051362, 2026.

4

L. Qi, C. Cui, The second-order augmented Zarankiewicz number, arXiv:2610.04448v1.