# Every 4×4 biquadratic sum of squares is a sum of eleven squares

Real algebraic geometry · sums of squares on projective varieties

Carles Marín

## Statements


Theorem 1 ..... 3
Lemma 2 ..... 4
Lemma 3 ..... 6
Lemma 4 — a regular point ..... 8
Proposition 5 ..... 9
Proposition 6 ..... 10
Remark 7 ..... 11
Proposition 8 ..... 11

# At a glance

> The exact value **BSR(4,4) = 11**: every biquadratic sum of squares in $x_{1..4}, y_{1..4}$ is a sum of eleven squares of bilinear forms. Qi and Cui gave a form that needs eleven; the previous upper bound was twelve. The proof classifies the Hankel multipliers of rank at most five whose image contains no product (Lemmas 2 and 3) and rules out the two shapes left with second variations built from the alternating forms (Propositions 6 and 8).

🚀 **Start here: Theorem 1**: BSR(4,4) = 11.

**Main results:**


- **Theorem 1**: BSR(4,4) = 11

**8** results | ✅ **0** entirely in Lean | ◐ **5** partly in Lean

💻 **5** checked by a program | written proof only | (a result can carry several tags)

### 🗺️ Map of the results (click a box to go to it)

Dependencies in the written proof: an arrow A → B means that this proof of B uses A. Every result rests on its written proof; the tags record what was checked besides, and each statement's "Evidence & scope" block says how far. The colour of a box is its strongest tag, in the order above; the tags in the box give the rest. Boxes with a thick border are the main results; click a box to read the statement.




### 📐 Setting (copied from the paper, §§1–2)

Write $x = (x_1, \dots, x_4)$, $y = (y_1, \dots, y_4)$, $V = \mathbb{R}^4 \otimes \mathbb{R}^4$ (the bilinear forms) and $R_2$ the biquadratic forms, of dimension 100. $\mathrm{BSR}(m, n)$ is the least $r$ such that every biquadratic sum of squares in $m + n$ variables is a sum of $r$ squares of bilinear forms; $\Phi(h_1, \dots, h_{11}) = h_1^2 + \cdots + h_{11}^2$, $\Sigma_{11} = \Phi(V^{11})$ and $\Sigma$ the cone of sums of squares. A *multiplier* is a symmetric bilinear form $H$ on $A \otimes B$ with the Hankel symmetry

$H(x \otimes b, y \otimes c) = H(x \otimes c, y \otimes b)$; its *supports* $a, b_0$ are the dimensions of the two factors modulo the radicals. Writing $H = T^{*}JT$ with $T$ onto $E$, $\dim E = \operatorname{rk} H$, and $T_b x = T(x \otimes b)$, the symmetry reads $\langle T_b x, T_c y\rangle_J = \langle T_c x, T_b y\rangle_J$.

# 1 The result

**Theorem 1.** p. 1

> $\mathrm{BSR}(4,4) = 11$.

**Isabelle · ours · part** **show**

**Only a part is in Isabelle:** partly: the regular point, the polar multiplier of Lemma 2 (iii) over the reals, the trace count of Proposition 8, and pieces of Lemma 3 (the support count, the factorisation in the case s = a = r, the case s < a under two hypotheses); outside the formal part: the passage to the surjectivity of dΦ, the extension of the polar multiplier to C, the identification of a simple tensor in im H, the Segre bound of Lemma 3, Lemma 2, Proposition 5 and the second variations

`regular_point_symmetric` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 25 — the regular point (Lemma 4)

```
theorem regular_point_symmetric:
  fixes H :: "nat \<Rightarrow> nat \<Rightarrow> nat \<Rightarrow> nat \
<Rightarrow> real" and n :: nat
  assumes sym: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c
< n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H c d a b"
    and hankel: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow>
c < n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H a d c b"
    and van: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c <
n \<Longrightarrow> d < n \<Longrightarrow> H a b c d + H b a c d = 0"
    and idx: "a < n" "b < n" "c < n" "d < n"
  shows "H a b c d = 0"
```

`support_triples` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 54 — the count closing Lemma 3

```
lemma support_triples:
  fixes a b r :: nat
  assumes "2 \<le> a" "a \<le> 4" "2 \<le> b" "b \<le> 4" "r \<le> 5" "r \<le>
(a - 1) * (b - 1)" "a + 1 \<le> r" "b + 1 \<le> r"
  shows "(a, b, r) \<in> {(4, 4, 5), (4, 3, 5), (3, 4, 5), (3, 3, 4)}"
```

`trace_count` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 92 — the trace count of Proposition 8

```
lemma trace_count:
  fixes lam :: "'i \<Rightarrow> real" and I :: "'i set"
  assumes fin: "finite I" and le1: "\<And>i. i \<in> I \<Longrightarrow> lam i \
<le> 1"
    and rk: "card {i \<in> I. lam i \<noteq> 0} \<le> 4" and neg: "j \<in> I"
"lam j < 0"
  shows "(\<Sum>i\<in>I. lam i) < 3"
```

**Trust base:** 🔒 kernel only — **only part of the statement is formal** (see the box). Isabelle: 3 results, no oracles

**Proof:** in this note · **Formalized:** Isabelle/HOL, by the author (public formalizations repository) · **Covers:** part: the regular point, the polar multiplier of Lemma 2 (iii) over the reals, the trace count of Proposition 8, and pieces of Lemma 3 (the support count, the factorisation in the case s = a = r, the case s < a under two hypotheses); outside the formal part: the passage to the surjectivity of dΦ, the extension of the polar multiplier to C, the identification of a simple tensor in im H, the Segre bound of Lemma 3, Lemma 2, Proposition 5 and the second variations · **Cite:** Theorem 1, p. 1, doi:10.5281/zenodo.24055900

> 🆕 **Classical and new:** the paper: the lower bound 11 is [QC, Theorem 3.1]; the previous upper bound 12 follows from BSR(m,3) = 2m [Mar, Corollary 7].

↳ Not used by another result of the paper.

Take a biquadratic form in $x_1, \dots, x_4$ and $y_1, \dots, y_4$ that is a sum of squares of bilinear forms, however many. The theorem says that eleven squares always suffice, and Qi and Cui gave a form for which ten do not. Before this note the best upper bound was twelve, from width three. The proof does not search for representations: it shows that the set of sums of eleven squares has no boundary inside the cone, because any boundary point would carry a Hankel multiplier of rank at most five that the classification of Lemmas 2–3 and the second variations of Propositions 6 and 8 rule out.

🛡️ **Evidence & scope**

- **Program:** perturbations P + εb² of the Qi–Cui witness: approximate representations with eleven squares in all 312 cases. working script `perturbar_qc.py`, not published

# 2 Hankel multipliers of small rank

**Lemma 2.** 📕 p. 2

> *There is no multiplier on $\mathbb{C}^4 \otimes \mathbb{C}^4$ of rank 5, with supports $a = b_0 = 4$, whose image contains no simple tensor.*

Isabelle · ours · part **show**

**Only a part is in Isabelle:** only the last step of case (iii), over the reals: X ↦ (tr(X)I + Xᵀ)/2 is injective on real matrices (polar_trace_injective); the paper needs complex matrices, which follows by applying it to the real and imaginary parts — that step is not formalized

`polar_trace_injective` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 67 — case (iii): X ↦ (tr(X) I + Xᵀ)/2 is injective, so the polar of (xᵀb)² has rank 16, not 5

```
lemma polar_trace_injective:
  fixes X :: "nat \<Rightarrow> nat \<Rightarrow> real" and n :: nat
  assumes zero: "\<And>i j. i < n \<Longrightarrow> j < n \<Longrightarrow> (if
i = j then (\<Sum>k<n. X k k) else 0) + X j i = 0"
    and ij: "i < n" "j < n"
  shows "X i j = 0"
```

**Trust base:** 🔒 kernel only — **only part of the statement is formal** (see the box). Isabelle: 1 result, no oracles

**Proof:** in this note · **Formalized:** Isabelle/HOL, by the author (public formalizations repository) · **Covers:** part: only the last step of case (iii), over the reals: X ↦ (tr(X)I + X$^{T}$)/2 is injective on real matrices (polar_trace_injective); the paper needs complex matrices, which follows by applying it to the real and imaginary parts — that step is not formalized · **Cite:** Lemma 2, p. 2, doi:10.5281/zenodo.24055900

↳ Used in: Theorem 1 (p. 3), Lemma 3 (p. 6), Proposition 6 (p. 10)

## Proof

The three cases below are exhaustive: $s = 4$ with some $Q(b)$ invertible, $s = 4$ with every $Q(b)$ singular, and $s \le 3$.

*(i) Some $Q(b)$ is invertible.* Choosing bases by congruence, $E = \mathbb{C}^4 \oplus \mathbb{C}$ with the standard symmetric product and $T_b x = (x, 0)$. In a basis of a complement of $b$ write $T_i x = (A_i x,\ u_i^{\top} x)$, $i = 1, 2, 3$. By (2),

$$A_i^{\top} = A_i, \qquad [A_i, A_j] = u_j u_i^{\top} - u_i u_j^{\top}. \tag{1}$$

With $v \odot w = v w^{\top} + w v^{\top}$ and $K_{ij} = A_i u_j - A_j u_i$, the Jacobi identity for $A_1, A_2, A_3$ and (4) give

$$K_{12} \odot u_3 + K_{23} \odot u_1 + K_{31} \odot u_2 = 0. \tag{2}$$

If the $u_i$ span $0$, $T$ is not onto. If they span a line, the last coordinate of $T$ is a simple covector of $\operatorname{im} H$. If they span a plane, after a change of basis of the complement $u_3 = 0$ with $u_1, u_2$ independent; then $A_3$ commutes with $A_1, A_2$, and (5) gives $(A_3 u_1) \odot u_2 = (A_3 u_2) \odot u_1$, whence $A_3 u_1 = t u_1$, $A_3 u_2 = t u_2$ for one $t$ (compare coefficients in a basis starting with $u_1, u_2$). The space $P = \operatorname{im}(A_3 - tI)$ is non-zero (otherwise $T_3 = t T_b$ and $B$ would have a radical, against $b_0 = 4$), stable under all $A_i$, and orthogonal to $u_1, u_2$ since $A_3$ is symmetric; by (4) the $A_i$ commute on $P$, and a common eigenvector $x \in P$ has $T_c x \in \mathbb{C}(x, 0)$ for all $c$, so the Observation applies. If the $u_i$ are independent, (5) forces $K_{ij} = t_i u_j - t_j u_i$ for three scalars $t_i$ (complete $u_1, u_2, u_3$ to a basis: the terms with the fourth vector and the squares $u_i^2$ force $K_{12} \in \langle u_1, u_2 \rangle$, $K_{23} \in \langle u_2, u_3 \rangle$, $K_{31} \in \langle u_3, u_1 \rangle$, and the coefficients of $u_1 u_2$, $u_2 u_3$, $u_3 u_1$ give the signs). Then the symmetric matrices

$$\widehat{A}_i = \begin{pmatrix} A_i & u_i \\ u_i^{\top} & t_i \end{pmatrix}$$

commute (the upper-left block of their commutator vanishes by (4), the upper right one is $K_{ij} + t_j u_i - t_i u_j = 0$, and the rest follows by antisymmetry). A common eigenvector $z \neq 0$ exists over $\mathbb{C}$; if $\widehat{A}_i z = \lambda_i z$, then $\langle z, T_i x \rangle = \lambda_i \langle z, T_b x \rangle$, since $T_i x = \widehat{A}_i T_b x$, so $T^{*} J z$ is simple, and non-zero because $T$ is onto.

*(ii) Some $T_b$ has rank $4$ and every $Q(b)$ is singular.* Then $U = \operatorname{im} T_b$ is a degenerate hyperplane with a radical of dimension one. In a Witt basis $E = \mathbb{C}^3 \oplus \langle e, f \rangle$, $\langle e, f \rangle = 1$

, $\langle e,e\rangle = \langle f,f\rangle = 0$, with $T_b(w,z) = (w,z,0)$, condition (2) gives each $T_c$ the shape $\begin{pmatrix} A & p \\ l^\top & t \\ p^\top & q_0 \end{pmatrix}$ with $A = A^\top$. In $Q(b+\tau c)$ the upper $3\times 3$ block is $I + O(\tau)$, the upper right block is $2\tau p + O(\tau^2)$ and the lower right entry is $2\tau q_0 + \tau^2(p^\top p + 2tq_0)$. Since the determinant vanishes identically, so does the Schur complement as a formal series; its terms of order one and two give $q_0 = 0$ and $-3p^\top p = 0$. Thus $p(c)^\top p(c) = 0$ for every $c$, and $c \mapsto p(c)$ is linear; *by polarization* $p(c)^\top p(c') = 0$ for all $c, c'$, so $\{p(c)\}$ spans a totally isotropic subspace of $\mathbb{C}^3$, of dimension at most one: $p(c) = \beta(c)p_0$. The covector $T^*Je$, $(w,z;c) \mapsto p(c)^\top w = \beta(c)\, p_0^\top w$, is simple and non-zero.

*(iii)* $s \le 3$. By (3) and $Q(b) \neq 0$ generically, $s = 3$ and $\operatorname{rk} Q(b) = 1$ generically: $U$ has a radical of dimension at least two and $Q(b) \neq 0$. So $q(x,b) = x^\top Q(b) x$ is, over $\mathbb{C}(b)$, a scalar times the square of a linear form in $x$ (complete the square from a non-zero diagonal entry). By Gauss' lemma $q = g(b)L(x,b)^2$ in $\mathbb{C}[b][x]$ with $L$ primitive, and the factors of a bihomogeneous polynomial are bihomogeneous. If $L$ has degree 0 in $b$, the support in $A$ is 1. Otherwise $g$ is constant and $L = x^\top Pb$, with $P$ invertible by full support, so after changes of basis and scale $q = (x^\top b)^2$. Its polar multiplier acts on matrices as $X \mapsto (\operatorname{tr}(X)I + X^\top)/2$, which is invertible ($X^\top = -\operatorname{tr}(X)I$ gives $5\operatorname{tr} X = 0$ and $X = 0$): rank $16 \neq 5$.

**Evidence & scope**

- **Program:** the Jacobi identity (5), the Schur expansion of case (ii) and the rank 16 of the polar of $(x^\top b)^2$, in exact arithmetic. `supplement.zip: check_symbolic_steps.py`

**Lemma 3.** p. 3

*Let $H$ be a nonzero multiplier on $\mathbb{C}^4 \otimes \mathbb{C}^4$ of rank $r \le 5$ whose image contains no simple tensor. Up to exchanging the factors, $(a, b_0, r)$ is either $(3,4,5)$ or $(3,3,4)$.*

**Isabelle · ours · part** show

**Only a part is in Isabelle:** pieces: the closing count (support_triples); the case s = a = r (case_s_eq_a), the case s < a with two hypotheses (the inclusion in im T_b of (3) and a one-dimensional radical; case_s_lt_a) and the Observation (observation), each concluding a nonzero FACTORISATION of the functional — in all three, the passage to a nonzero simple tensor α ⊗ β in im H uses the bilinearity of the paper and is not formalized. The Segre bound r ≤ (a−1)(b₀−1), the generic b and the assembly are not formalized

`support_triples` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 54 — the arithmetic that closes Lemma 3

```
lemma support_triples:
  fixes a b r :: nat
  assumes "2 \<le> a" "a \<le> 4" "2 \<le> b" "b \<le> 4" "r \<le> 5" "r \<le>
(a - 1) * (b - 1)" "a + 1 \<le> r" "b + 1 \<le> r"
  shows "(a, b, r) \<in> {(4, 4, 5), (4, 3, 5), (3, 4, 5), (3, 3, 4)}"
```

`case_s_eq_a` · `Isabelle/BSRFourFour/Supports_Lemma.thy`, line 286 — the case s = a = r: the operators T_b⁻¹T_c commute, a common eigenvector gives a simple covector in the image

```
theorem case_s_eq_a:
  fixes J :: "'e::ab_group_add \<Rightarrow> 'e \<Rightarrow> complex" and T ::
"'c \<Rightarrow> 'a::ab_group_add \<Rightarrow> 'e"
```

```
    and sm :: "complex \<Rightarrow> 'e \<Rightarrow> 'e" and smA :: "complex \
<Rightarrow> 'a \<Rightarrow> 'a"
  assumes fd: "FinDimVectorSpace smA (UNIV :: 'a set)" and nz: "\<exists>w::'a.
w \<noteq> 0"
    and sym: "\<And>b c x y. J (T b x) (T c y) = J (T c x) (T b y)"
    and Jadd: "\<And>u v w. J (u + v) w = J u w + J v w"
    and Jscal: "\<And>t u w. J (sm t u) w = t * J u w"
    and nondeg: "\<And>z. (\<And>w. J z w = 0) \<Longrightarrow> z = 0"
    and Tadd: "\<And>c u v. T c (u + v) = T c u + T c v"
    and Tscal: "\<And>c t u. T c (smA t u) = sm t (T c u)"
    and Tb_inj: "\<And>u. T b u = 0 \<Longrightarrow> u = 0"
    and Tb_onto: "\<And>w. \<exists>y. w = T b y"
  shows "\<exists>e \<beta>. e \<noteq> 0 \<and> \<beta> b \<noteq> 0 \<and> (\
<forall>c y. J e (T c y) = (\<beta> c / \<beta> b) * J e (T b y))
           \<and> (\<exists>y. J e (T b y) \<noteq> 0)"
```

`case_s_lt_a` · `Isabelle/BSRFourFour/Supports_Lemma.thy`, line 102 — the case s < a, with the inclusion T_c(ker T_b) ⊆ im T_b and a one-dimensional radical as hypotheses

```
theorem case_s_lt_a:
  fixes J :: "'e::ab_group_add \<Rightarrow> 'e \<Rightarrow> complex" and T ::
"'c \<Rightarrow> 'a \<Rightarrow> 'e"
    and sm :: "complex \<Rightarrow> 'e \<Rightarrow> 'e"
  assumes sym: "\<And>b c x y. J (T b x) (T c y) = J (T c x) (T b y)"
    and Jscal: "\<And>t u w. J (sm t u) w = t * J u w"
    and Jadd: "\<And>u v w. J (u + v) w = J u w + J v w"
    and nondeg: "\<And>z. (\<And>c y. J z (T c y) = 0) \<Longrightarrow> z = 0"
    and x: "T b x = 0" and xrad: "T d0 x \<noteq> 0"
    and inU: "\<And>c. T c x \<in> range (T b)"
    and line: "\<And>u. u \<in> range (T b) \<Longrightarrow> (\<forall>y. J u
(T b y) = 0) \<Longrightarrow> \<exists>t. u = sm t e"
  shows "\<exists>e' \<beta> d. e' \<noteq> 0 \<and> \<beta> d \<noteq> 0 \<and>
(\<forall>c y. J e' (T c y) = (\<beta> c / \<beta> d) * J e' (T d y))
           \<and> (\<exists>y. J e' (T d y) \<noteq> 0)"
```

`observation` · `Isabelle/BSRFourFour/Supports_Lemma.thy`, line 60 — the Observation of Section 2: T_c x = β(c) e gives a non-zero simple covector

```
theorem observation:
  fixes J :: "'e::ab_group_add \<Rightarrow> 'e \<Rightarrow> complex" and T ::
"'c \<Rightarrow> 'a \<Rightarrow> 'e"
    and sm :: "complex \<Rightarrow> 'e \<Rightarrow> 'e"
  assumes sym: "\<And>b c x y. J (T b x) (T c y) = J (T c x) (T b y)"
    and Jscal: "\<And>t u w. J (sm t u) w = t * J u w"
    and nondeg: "\<And>z. (\<And>c y. J z (T c y) = 0) \<Longrightarrow> z = 0"
    and Tx: "\<And>c. T c x = sm (\<beta> c) e" and d: "\<beta> d \<noteq> 0"
and e: "e \<noteq> 0"
  shows "(\<forall>c y. J e (T c y) = (\<beta> c / \<beta> d) * J e (T d y)) \
<and> (\<exists>y. J e (T d y) \<noteq> 0)"
```

`common_eigenvector` · `Isabelle/BSRFourFour/Supports_Lemma.thy`, line 154 — commuting endomorphisms over an algebraically closed field have a common eigenvector (classical; proved here)

```
theorem (in FinDimVectorSpace) common_eigenvector:
  assumes alg_closed: "\<And>p::'a poly. degree p > 0 \<Longrightarrow> \
<exists>c. poly p c = 0"
    and add: "\<And>T u v. T \<in> F \<Longrightarrow> u \<in> V \
<Longrightarrow> v \<in> V \<Longrightarrow> T (u + v) = T u + T v"
    and hom: "\<And>T a u. T \<in> F \<Longrightarrow> u \<in> V \
<Longrightarrow> T (a \<cdot> u) = a \<cdot> T u"
    and comm: "\<And>S T u. S \<in> F \<Longrightarrow> T \<in> F \
<Longrightarrow> u \<in> V \<Longrightarrow> S (T u) = T (S u)"
  shows "Subspace W \<Longrightarrow> (\<exists>w\<in>W. w \<noteq> 0) \
<Longrightarrow> (\<And>T. T \<in> F \<Longrightarrow> T ` W \<subseteq> W)
           \<Longrightarrow> \<exists>u\<in>W. u \<noteq> 0 \<and> (\<forall>T\
<in>F. \<exists>c. T u = c \<cdot> u)"
```

**Trust base:** 🔒 kernel only — **only part of the statement is formal** (see the box). Isabelle: 5 results, no oracles

**Proof:** in this note · **Formalized:** Isabelle/HOL, by the author (public formalizations repository) · **Covers:** part: pieces: the closing count (support_triples); the case s = a = r (case_s_eq_a), the case s < a with two hypotheses (the inclusion in im T_b of (3) and a one-dimensional radical; case_s_lt_a) and the Observation (observation), each concluding a nonzero FACTORISATION of the functional — in all three, the passage to a nonzero simple tensor α ⊗ β in im H uses the bilinearity of the paper and is not formalized. The Segre bound r ≤ (a−1)(b₀−1), the generic b and the assembly are not formalized · **Cite:** Lemma 3, p. 3, doi:10.5281/zenodo.24055900

↳ Used in: Theorem 1 (p. 3)

**Proof**

Restrict $H$ to $A' \otimes B'$, with $A', B'$ complements of the two radicals: the rank, the symmetry (1) and the absence of simple tensors are kept (a simple covector of the restricted image extends by zero on the radicals and stays simple and in $\operatorname{im} H$), and now both radicals are zero, with $\dim A' = a$, $\dim B' = b_0$. Since $\operatorname{im} H$ annihilates the radicals, $\mathbb{P}(\operatorname{im} H)$ is a subspace of dimension $r-1$ avoiding the Segre variety of the effective factors, so $r \le (a-1)(b_0-1)$; and $a, b_0 \ge 2$ since $H \neq 0$.

We show $r \ge a+1$, and by the symmetry of (1) in the two factors also $r \ge b_0+1$. Suppose $r \le a$ ($\le 4$) and take $b$ generic, with $s = \operatorname{rk} T_b$ maximal and $Q(b) \neq 0$. If $s<a$, then $\ker T_b \neq 0$, and the radical of $U = \operatorname{im} T_b$ has dimension at most $s-1$ (as $Q(b) \neq 0$) and at most $r-s$ (as it lies in $U^{\perp_J}$), so at most $\min(s-1, r-s) \le 1$ because $r \le 4$; this contradicts (3) and the Observation. If $s = a$, then $s = a = r$, $T_b : A' \to E$ is an isomorphism, $Q(b)$ is non-degenerate, and the operators $T_b^{-1}T_c$ are self-adjoint for $Q(b)$ and, by (2), commute; a common eigenvector $x$ has $T_c x \in \mathbb{C}\, T_b x$ for all $c$, and the Observation gives a simple tensor. So $r \ge \max(a, b_0)+1$.

With $r \le 5$ and $r \le (a-1)(b_0-1)$ the only possibilities are $(a, b_0, r) = (4,4,5)$, $(4,3,5)$, $(3,4,5)$ and $(3,3,4)$, and Lemma 2 excludes $(4,4,5)$.

**Evidence & scope**

- **Program:** Newton search for rank-five Hankel multipliers with support 3×4 and product-free image: 14 numerical candidates. `supplement.zip: search_rank_five_multipliers.py`

# 3 The boundary multiplier

**Lemma 4. (a regular point)** p. 4

*Identify both factors with $\mathbb{R}^4$. Let $S \subset V$ be the span of the four forms $x_i y_i$ and the six forms $x_i y_j + x_j y_i$ for $i<j$. If $H$ is a Hankel multiplier and $S \subseteq \ker H$, then $H = 0$. Consequently $d\Phi$ is surjective at the tuple consisting of these ten forms and zero.*

Isabelle · ours · part **show**

**Only a part is in Isabelle:** the linear algebra; the duality from there to «dΦ onto» is on paper

`regular_point_symmetric` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 25 — a symmetric Hankel multiplier that vanishes on the symmetric tensors in its first argument is zero (in any size n)

```
theorem regular_point_symmetric:
  fixes H :: "nat \<Rightarrow> nat \<Rightarrow> nat \<Rightarrow> nat \
<Rightarrow> real" and n :: nat
  assumes sym: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c
< n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H c d a b"
    and hankel: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow>
c < n \<Longrightarrow> d < n \<Longrightarrow> H a b c d = H a d c b"
    and van: "\<And>a b c d. a < n \<Longrightarrow> b < n \<Longrightarrow> c <
n \<Longrightarrow> d < n \<Longrightarrow> H a b c d + H b a c d = 0"
    and idx: "a < n" "b < n" "c < n" "d < n"
  shows "H a b c d = 0"
```

**Trust base:** 🔒 kernel only — **only part of the statement is formal** (see the box). Isabelle: 1 result, no oracles

**Proof:** in this note · **Formalized:** Isabelle/HOL, by the author (public formalizations repository) · **Covers:** part: the linear algebra; the duality from there to «dΦ onto» is on paper · **Cite:** Lemma 4, p. 4, doi:10.5281/zenodo.24055900

↳ Used in: Theorem 1 (p. 3), Proposition 5 (p. 9)

**Proof**

If $S \subseteq \ker H$, then $H(s,v) = H(v,s) = 0$ for all $s \in S$ and $v \in V$, and $S$ is the space of symmetric tensors, so $H(x \otimes y, x \otimes y) = -H(x \otimes y, y \otimes x) = -H(x \otimes x, y \otimes y) = 0$ for all $x, y$, and $H = 0$ by polarization. Hence the products of the ten summands with $V$ span $R_2$ (we also checked the rank 100 exactly), and adding an eleventh summand, even zero, gives a point where $d\Phi$ is onto.

**Evidence & scope**

- **Program:** rank 100 of the products of the ten summands with V, in exact arithmetic; control with the four diagonal summands only: 58. `supplement.zip: check_regular_point.py`

**Proposition 5.** p. 4

*If $\Sigma_{11} \neq \Sigma$, there are $F \in \operatorname{int}\Sigma \cap \partial\Sigma_{11}$, a representation $F = \sum_{i=1}^{11} h_i^2$ and an indefinite real multiplier $H$ on $V$ such that, with $W = \operatorname{span}(h_i)$: $W \subseteq \ker H$, $\dim W = 11$ and hence $\operatorname{rk} H \leq 5$; the complex image of $H$ contains no simple tensor; and $\sum_i h_i D_i = 0$ in $R_2$ implies $\sum_i H(D_i, D_i) \geq 0$.*

🆕 **Classical and new:** the paper: the argument of [Mar, Proposition 3 and Corollary 4], after [BDS]; only the jet count $100 - 14 + 12 = 98 < 99$ depends on the size.

↳ Used in: Theorem 1 (p. 3), Proposition 6 (p. 10), Proposition 8 (p. 11)

**Proof**

We adapt the argument of [ 3 , Proposition 3 and Corollary 4 ] to $(m,n) = (4,4)$, using Lemma 4. The only count that depends on the size is the one excluding forms with a singular point with no real factor. Take a pair of conjugate points distinct in both factors

and change coordinates over $\mathbb{C}$ to $(e_1, e_1), (e_2, e_2)$. A first jet has seven coordinates, the value and the six tangent derivatives, and the two first jets use disjoint sets of seven monomial coefficients, so the joint jet map is surjective over $\mathbb{C}$. Conjugation exchanges the two jets, and the restriction to real forms has real rank fourteen. The point varies in real dimension twelve. The incidence therefore has dimension at most $100 - 14 + 12 = 98$, and its projection cannot fill the boundary patch of dimension 99. A singular point with a real factor is impossible at an interior form, by definiteness of the quadratic form in the other factor. Independence of the $h_i$ is as in [ 3 , Corollary 4 ] .

# 4 Support $3 \times 4$, rank five

**Proposition 6.** p. 4

> *No triple $(F, h, H)$ satisfying the conclusions of Proposition 5 can have $(a, b_0, r) = (3, 4, 5)$ or $(4, 3, 5)$.*

↳ Used in: Theorem 1 (p. 3), 7 (p. 11)

**Proof**

Here $\dim \ker H = 16 - 5 = 11$, so $W = \ker H$. Let $x_0$ span the radical in the first factor and $A'$ be a real complement, $\dim A' = 3$. Then $x_0 \otimes B \subseteq W$, and an orthogonal change of the eleven summands (an orthonormal basis of the preimage of $x_0 \otimes B$, completed) gives $h_i = x_0 \otimes y_i$, $1 \le i \le 4$, for a real basis $(y_i)$ of $B$. For $z_{ij} = -z_{ji} \in A'$ put

$$D_i = \sum_{j=1}^{4} z_{ij} \otimes y_j \quad (i \le 4), \qquad D_i = 0 \quad (i{>}4).$$

Then $\sum_i h_i D_i = \sum_{i,j} x_0 z_{ij}\, y_i y_j = 0$ by antisymmetry: these are syzygies, a space $A' \otimes \Lambda^2 B$ of dimension 18. With $H = T^{*} J T$ as in §2 (now over $\mathbb{R}$, $\dim E = 5$), the second variation is $\sum_i H(D_i, D_i) = J^{\oplus 4}(\Psi z)$ for

$$\Psi : A' \otimes \Lambda^2 B \to E^4, \qquad \Psi(z) = \Big( \sum_j T_{y_j} z_{ij} \Big)_{i=1}^{4}.$$

By Proposition 5, $J^{\oplus 4}$ is non-negative on $\operatorname{im} \Psi$. Since $H$ is indefinite, the positive index $p$ of $J$ is at most 4, and a subspace on which $J^{\oplus 4}$ is non-negative has dimension at most $4p \le 16$. Hence $\operatorname{rk} \Psi \le 16$.

On the other hand $\operatorname{rk} \Psi = 17$. Complexify and compute the annihilator of $\operatorname{im} \Psi$ under $J^{\oplus 4}$: a tuple $(c_i) \in E^4$, i.e. a map $C : B \to E$ with $C y_i = c_i$, annihilates $\operatorname{im} \Psi$ if and only if the matrix $\big( \langle C y_i, T(a \otimes y_j) \rangle_J \big)_{ij}$ is symmetric for every $a \in A'$ (vary each $z_{ij}$, $i{<}j$, separately). The three maps $C = T(a \otimes \cdot)$, $a$ in a basis of $A'$, are independent and lie in the annihilator by (2). If there were a fourth independent $C$, it would extend $T$ to $\widetilde{T} : \mathbb{C}^4 \otimes B \to E$ satisfying (2): the identities involving the new row are exactly the symmetry just written, and the one of the new row with itself is automatic. Then $\widetilde{T}^{*} J \widetilde{T}$

is a multiplier of rank 5 (as $T$ is onto) with supports 4 and 4 (by independence of the new row, and since the second support was already full), and its image contains no simple tensor: a simple covector in it restricts to $A' \otimes B$ as a simple covector of $\operatorname{im} H$, excluded by Proposition 5, or as zero, and then its representing vector $z$ has $\langle z, T(A' \otimes B)\rangle_J = 0$, so $z = 0$. This contradicts Lemma 2. So the annihilator has dimension 3 and $\operatorname{rk} \Psi = 20 - 3 = 17 > 16$, a contradiction.

**Evidence & scope**

- **Program:** on the rank-five multiplier with support 3×4 of the supplement of [Mar]: rank $\Psi = 17 > 16$ and a negative value of the second variation; in the 14 numerical candidates rank $\Psi = 17$, and 15 in a control built from five product terms. `supplement.zip: check_proof_steps.py`

**Remark 7.** p. 5

One triple of quaternionic structures does not suffice: for $z_{ij} = \sum_a \eta^a_{ij} v_a$ with the 't Hooft symbols $\eta^a$, the second variation is $\sum_a v_a^\top (\sum_j H_{jj}) v_a$, because $(\eta^a)^\top \eta^b$ has symmetric part $\delta_{ab} I$ and $H_{jk} = H_{kj}$; it only says that $\sum_j H_{jj} \succeq 0$. The argument above uses the whole of $\Lambda^2 B$.

**Cite:** Remark 7, p. 5, doi:10.5281/zenodo.24055900

↳ Not used by another result of the paper.

# 5 Support $3 \times 3$, rank four

**Proposition 8.** p. 5

*No triple $(F, h, H)$ satisfying the conclusions of Proposition 5 can have $(a, b_0, r) = (3, 3, 4)$.*

**Isabelle · ours · part** **show**

**Only a part is in Isabelle:** the final trace count of the case u ∉ W (trace_count); the two second variations are on paper

`trace_count` · `Isabelle/BSRFourFour/BSR44_Pieces.thy`, line 92 — eigenvalues at most 1, at most four non-zero, one negative: their sum is less than 3

```
lemma trace_count:
  fixes lam :: "'i \<Rightarrow> real" and I :: "'i set"
  assumes fin: "finite I" and le1: "\<And>i. i \<in> I \<Longrightarrow> lam i \
<le> 1"
    and rk: "card {i \<in> I. lam i \<noteq> 0} \<le> 4" and neg: "j \<in> I"
"lam j < 0"
  shows "(\<Sum>i\<in>I. lam i) < 3"
```

**Trust base:** 🔒 kernel only — **only part of the statement is formal** (see the box). Isabelle: 1 result, no oracles

**Proof:** in this note · **Covers:** not formalized · **Cite:** Proposition 5, p. 4, doi:10.5281/zenodo.24055900

↳ Used in: Theorem 1 (p. 3)

**Proof**

Let $x_0, y_0$ span the real radicals of the two factors, $A', B'$ complements and $u = x_0 \otimes y_0$. The space $U = x_0 \otimes B + A \otimes y_0$ has dimension 7 and lies in $K = \ker H$, $\dim K = 12$; $W$ is a hyperplane of $K$, and $K = U \oplus K_0$ with $K_0 = K \cap (A' \otimes B')$.

*Case* $u \notin W$. Then $W \cap (x_0 \otimes B) = x_0 \otimes L$ with $\dim L = 3$ and $L$ projecting isomorphically onto $B/\langle y_0 \rangle$. Orthogonalizing on the preimage of this intersection, $h_i = x_0 \otimes y_i$, $i \le 3$, for a basis of $L$. The restriction of $H$ to $A' \otimes L$ keeps the indices, the rank four and the support three in the first factor. For $v_1, v_2, v_3 \in A'$ put $D_i = \sum_{j,p} \varepsilon_{ijp}\, v_p \otimes y_j$ for $i \le 3$ and $D_i = 0$ otherwise; they are syzygies by antisymmetry, and with $H_{jl}$ the blocks of the restriction and $S = \sum_j H_{jj}$, Proposition 5 and the contraction of the alternating symbols give, as in [ 3 , § 5 ] ,

$$I_3 \otimes S - H|_{A' \otimes L} \succeq 0.$$

The diagonal blocks sum to $2S$, so $S \succeq 0$; if $Sx = 0$, the three diagonal values of this positive semidefinite form at $x \otimes y_j$ sum to zero, so each $x \otimes y_j$ is in its kernel, $H(x \otimes y_j, \cdot) = 0$, and $x = 0$ by full support in $A'$. So $S \succ 0$, and after the congruence by $S^{-1/2}$ in the three blocks the matrix of $H$ has trace 3, all eigenvalues at most 1, rank four and at least one negative eigenvalue: its trace is less than 3, a contradiction.

*Case* $u \in W$. An orthogonal change makes $h_1$ a non-zero multiple of $u$. Let $\ell$ be a real linear functional on $K$ with $W = \ker \ell$. For $v = \sum v_{ij} a_i \otimes b_j \in A' \otimes B'$ (bases $a_i$, $b_j$) the lift $P(v) = \sum_{ij} v_{ij} (a_i \otimes y_0)(x_0 \otimes b_j) \in \operatorname{Sym}^2 U$ has product $uv$ in $R_2$. If $\ell|_U = 0$ all these products lie in $WK$; otherwise $\operatorname{Sym}^2 U / (U \cap W) U$ is one-dimensional and the class is measured by $(\ell|_U)^2$. So $uv \in WK$ whenever

$$\phi(v) = \sum_{ij} v_{ij}\, \ell(a_i \otimes y_0)\, \ell(x_0 \otimes b_j) = 0.$$

The functional $\phi$ on the active block is zero or simple. If it is non-zero it does not vanish on $K_0$, since otherwise it would lie in the image of the active part of $H$, which contains no simple tensor. Choose $v$ in the active block with $H(v, v) < 0$; adding a real element of $K_0$ keeps this value and achieves $\phi(v) = 0$. Then $h_1 v \in WK$: write $-h_1 v = \sum_i h_i k_i$ with $k_i \in K$. The $D_i = \delta_{i1} v + k_i$ form a syzygy with second variation $H(v, v) < 0$, a contradiction.

# 6 Checks and status

No numbered statement in this section.

It is prose only (no statement, figure or table): read it in the paper (PDF).

# The companion site

No numbered statement in this section.

It is prose only (no statement, figure or table): read it in the paper (PDF).

# Acknowledgement

No numbered statement in this section.

It is prose only (no statement, figure or table): read it in the paper (PDF).

# References

1

G. Blekherman, A. Dunbar, R. Sinn, Pythagoras numbers for ternary forms, arXiv:2410.17123v2.

2

L. Chen, D. Ž. Đoković, Properties and construction of extreme bipartite states having positive partial transpose, Comm. Math. Phys. 323 (2013), 241–284; arXiv:1203.1364.

3

C. Marín, For $m \geq 2$, every $m \times 3$ biquadratic sum of squares is a sum of $2m$ squares, Zenodo, doi:10.5281/zenodo.24051362, 2026.

4

L. Qi, C. Cui, The second-order augmented Zarankiewicz number, arXiv:2610.04448v1.