
Map of *Characters of Sp(2m) at the powers of Kostant's element*

Representation theory · characters of Lie groups | 05E10 | 20G05 | 17B22 | 05A15

12 results | ✅ **0** formalized in Lean | ◐ **0** with a part in Lean | 💻 **10** checked by a program | written proof only

(a result can carry several tags)

Dependencies in the written proof: an arrow A → B means that this proof of B uses A. Every result rests on its written proof; the tags record what was checked besides, and each statement's “Evidence & scope” block says how far. The colour of a box is its strongest tag, in the order above; the tags in the box give the rest. Boxes with a thick border are the main results; click a box to read the statement.



What the paper contributes

Kostant's element a_p is a point where every irreducible character of a compact simple group takes a value in $\{0, \pm 1\}$. The paper determines the irreducible characters of Sp(2m) at **every power** a_p^k.

- **When a character vanishes, and its value** (Theorem A): it vanishes exactly when $\lambda+\rho$ lies on more walls of the modular root arrangement than ρ does, equivalently when some folded residue class holds more entries than its capacity; otherwise the value is an explicit signed ratio of root products.
- **The absolute value is a product of dimensions** of representations of GL, Sp and odd SO, times $\kappa \in \{1, 2\}$ (Theorem B).
- **Every rational point of the same line:** a non-vanishing character is, up to sign and that factor, such a product times a character of Sp(2s) or SO(2s), $s \leq m$, at a regular element (Theorem 10.1); the points where the closed form holds for all weights are listed (Theorem C), and at the points where all characters are integers the remaining factor is 1, 2, 3, 4 or 6 (Theorem D).

Known and new, as the paper states it. For k prime to 2m+2, Theorem A is due to Cellini, Möseneder Frajria and Papi; for gcd(k, 2m+2) odd, Theorems A and B follow from Albion's factorisations of universal characters, with the sign added. New: for gcd(k, 2m+2) even and $q \geq 3$, the values on the classes not conjugate to that of ρ and the factor κ; the list of points in Theorem C and the exclusion of the others; in Theorem D, the constants for $q \geq 4$ even with $q \mid 2m$, $q \nmid m$, and for $q = 6$ with $m \equiv 4 \pmod 6$.

Key tool: an identity of residue multisets (Proposition 5.1), which takes the place of the conjugacy hypothesis in the work of Nadimpalli, Pattanayak and Prasad.

🚀 **Start here: Theorem A**: when the character vanishes, and its value.

Main results:

- **Theorem A**: when the character vanishes, and its value
- **Theorem B**: |value| = κ × product of dimensions
- **Theorem C**: where the closed form holds for all weights
- **Theorem D**: the constant $c \in \{1,2,3,4,6\}$

🧭 Fields: where the paper lives and what it connects

Tools taken from other branches

- **Root systems** → walls of the modular root arrangement: the count criterion, proved by the argument of Nadimpalli–Pattanayak–Prasad (the alcove argument of Cellini–Möseneder Frajria–Papi needs ρ regular and no longer applies) — Proposition 3.1, Theorem A
- **Elementary number theory** → residues modulo q in folded classes; an identity of residue multisets — Corollary 4.1, Proposition 5.1
- **Algebraic combinatorics** → partitions, Weyl dimension formulas; the factorisations of universal and classical characters (Albion, Ayyer–Kumari) — Theorem B
- **Algebraic number theory** → Galois action on the spectrum: which points give integer characters (a lemma of Gannon–Schopieray) — Proposition 8.1
- **Enumeration** → counting the non-vanishing residues — Theorem 9.1

What the results serve

- **Characters at elements of finite order** ← Kostant's $\{0, \pm 1\}$ theorem, extended to every power of a_p; the question of Nadimpalli, Pattanayak and Prasad on vanishing and values at torsion elements, addressed for these elements of Sp(2m) — Theorem A, Theorem B
- **Reeder's constants** ← their values on the classes not conjugate to that of ρ — Theorem A, Proposition 5.1
- **Factorisations of classical characters** ← a factor κ and a residual character of type C or D at every rational point of the line — Theorem B, Theorem 10.1, Theorem D